

1 **Clinical Practice Guideline:** **Occupational Therapy Medical Policy/Guidelines**

2

3 **Date of Implementation:** **October 18, 2012**

4

5 **Product:** **Specialty**

6

7

Related Policies:

- CPG 12: Medical Necessity Decision Assist Guideline for Rehabilitative Care
- CPG 30: Laser Therapy (LT)
- CPG 110: Medical Record Maintenance and Documentation Policies
- CPG 111: Patient Assessments: Medical Necessity Decision Assist Guideline for Evaluations and Re-evaluations
- CPG 112: Exercise Therapy for Treatment of Non-Specific Low Back Pain
- CPG 113: Exercise Therapy for Treatment of Neck Pain
- CPG 121: Passive Physiotherapy Modalities
- CPG 133: Techniques and Procedures Not Widely Supported As Evidence-Based
- CPG 135: Physical Therapy Medical Policy/Guideline
- CPG 143: Strapping and Taping
- CPG 144: Prosthetic Training and Evaluation
- CPG 146: Range of Motion Testing
- CPG 148: Wheelchair Management
- CPG 152: Orthotic Training and Evaluation
- CPG 165: Autism Spectrum Disorder (ASD) – Outpatient Rehabilitation Services (Speech, Physical, and Occupational Therapy)
- CPG 166: Speech-Language Pathology/Speech Therapy Guidelines
- CPG 175: Extra-Spinal Joint Manipulation/Mobilization for the Treatment of Upper Extremity Musculoskeletal Conditions
- CPG 178: Dry Needling
- CPG 257: Developmental Delay Screening and Testing
- CPG 269: H-Wave® Electrical Stimulation
- CPG 270: Cognitive Rehabilitation
- CPG 272: Electric Stimulation for Pain, Swelling and Function
- CPG 273: Superficial Heat and Cold
- CPG 274: Deep Heating Modalities (Therapeutic Ultrasound and Diathermy)
- CPG 276: MEDEK Therapy
- CPG 277: Non-invasive Interactive Neurostimulation (InterX®)
- CPG 286: Intensive Model of Therapy
- CPG 295: Physical Performance Testing or Measurement
- CPG 305: Virtual Physical Therapy and Rehabilitation Services

1 **TABLE OF CONTENTS**
 2 *(CTRL+Click on Section Heading to Follow Link)*

3 **1. PROVIDERS OF OCCUPATIONAL THERAPY SERVICES..... 3**
 4 **2. REHABILITATIVE OCCUPATIONAL THERAPY SERVICES 5**
 5 **3. MAINTENANCE OCCUPATIONAL THERAPY SERVICES..... 9**
 6 **4. HABILITATIVE OCCUPATIONAL THERAPY SERVICES 11**
 7 **5. REDUNDANT THERAPEUTIC EFFECTS AND REHABILITATIVE OR**
 8 **HABILITATIVE SERVICES 13**
 9 **6. THERAPEUTIC MODALITIES AND PROCEDURES..... 13**
 10 6.1 Passive Care and Active Care.....16
 11 6.2 Treatment Interventions.....17
 12 6.3 Precautions and Contraindications to Therapeutic Modalities and Procedures23
 13 **7. CLINICAL DOCUMENTATION 26**
 14 7.1 Evaluation and Re-evaluations27
 15 7.2 Treatment Sessions29
 16 7.3 Discharge/Discontinuation of Intervention.....30
 17 7.4 Duplicated / Insufficient Information32
 18 7.5 Centers for Medicare and Medicaid Services (CMS)32
 19 **8. CLINICAL REVIEW PROCESS 32**
 20 8.1 Definition of Key Terminology used in Clinical Reviews33
 21 8.2 Clinical Quality Evaluation34
 22 8.3 Critical Factors during Clinical Reviews.....37
 23 8.3.1 General Factors.....38
 24 8.3.2 Factors that Influence Adverse Determinations of Clinical Services (Partial Approvals/Denials).....45
 25 8.3.3 Referral / Coordination of Services50
 26 **9. LITERATURE REVIEW 50**
 27 9.1 Occupational Therapy for Conditions Considered Unproven.....50
 28 9.2 Specific Occupational Therapy Treatments Considered Unproven.....51
 29 **10. CODING/BILLING INFORMATION..... 55**

1

2 DESCRIPTION

3 This document addresses Occupational Therapy Services which may be delivered by an
 4 Occupational Therapist acting within the scope of a professional license. This document
 5 also addresses the processes associated with Medical Necessity Determinations performed
 6 by American Specialty Health (ASH) clinical quality evaluators on services submitted for
 7 review.

8

9 The availability of coverage for rehabilitative and/or habilitative services will vary by
 10 benefit design as well as by State and Federal regulatory requirements. Benefit plans may
 11 include a maximum allowable rehabilitation benefit, either in duration of treatment or in
 12 number of visits or in the conditions covered or type of services covered. When the
 13 maximum allowable benefit is exhausted or if the condition or service are not covered,
 14 coverage will no longer be provided even if the medical necessity criteria described below
 15 are met.

16

17 GUIDELINES

18

1. PROVIDERS OF OCCUPATIONAL THERAPY SERVICES

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

Aides and other nonqualified personnel are limited to provision of non-skilled services
 such as preparing the individual, treatment area, equipment, or supplies; assisting a
 qualified therapist or assistant; and transporting individuals.

According to the American Occupational Therapy Association, occupational therapists and
 occupational therapy assistants help people across their lifespan participate in the things
 they want and need to do through the therapeutic use of everyday activities (occupations).
 Occupational therapists provide services to patients who have impairments, functional
 limitations, disabilities, or changes in physical function and health status resulting from
 injury, disease, or other causes. OT addresses physical, cognitive, psychosocial, sensory,
 communication, and other areas of performance in various contexts and environments in

1 everyday life activities that affect health, well-being, and quality of life. The overarching
2 goal of occupational therapy is “to support [people’s] health and participation in life
3 through engagement in occupations.”

4 A service is not considered a skilled therapy service merely because it is furnished by a
5 therapist or by a therapist/therapy assistant under the direct or general supervision, as
6 applicable, of a therapist. If a service can be self-administered or safely and effectively
7 furnished by an unskilled person, without the direct or general supervision, as applicable,
8 of a therapist, the service cannot be regarded as a skilled therapy service even though a
9 therapist furnishes the service. Similarly, the unavailability of a competent person to
10 provide a non-skilled service, notwithstanding the importance of the service to the patient,
11 does not make it a skilled service when a therapist furnishes the service.

12
13 Services that do not require the professional skills of a therapist to perform or supervise
14 are not medically necessary, even if they are performed or supervised by a therapist,
15 physician or NPP. Therefore, if a patient’s therapy can proceed safely and effectively
16 through a home exercise program, self-management program, restorative nursing program
17 or caregiver assisted program, occupational therapy services are not indicated or medically
18 necessary. Occupational therapy is used for both rehabilitation and habilitation. Skilled
19 occupational therapy services may be necessary to improve a patient’s current condition,
20 to maintain the patient’s current condition, or to prevent or slow further deterioration of
21 the patient’s condition.

22
23 The plan of care for medically necessary occupational therapy services is established by a
24 licensed occupational therapist. The amount, frequency and duration of the occupational
25 therapy services must be reasonable (within regional norms and commonly accepted
26 practice patterns); the services must be considered appropriate and needed for the treatment
27 of the condition and must not be palliative in nature. Thus, once therapeutic benefit has
28 been achieved, or a home exercise program could be used for further gains without the
29 need for skilled occupational therapy, continuing supervised occupational therapy is not
30 considered medically necessary.

31
32 Rehabilitative services are intended to improve, adapt or restore functions which have been
33 impaired or permanently lost as a result of illness, injury, loss of a body part, or congenital
34 abnormality involving goals an individual can reach in a reasonable period of time. If no
35 improvement is documented after two weeks of treatment, an alternative treatment plan
36 should be attempted. Treatment is no longer medically necessary when the individual stops
37 progressing toward established goals.

38
39 Habilitative services are defined by the National Association of Insurance Commissioners
40 as “health care services that help a person keep, learn or improve skills and functioning for

1 daily living.” Habilitative services are intended to maintain, develop or improve skills
 2 needed to perform activities of daily living (ADLs) or instrumental activities of daily living
 3 (IADLs) which have not (but normally would have) developed or which are at risk of being
 4 lost as a result of illness, injury, loss of a body part, or congenital abnormality. Examples
 5 include therapy for a child who is not walking at the expected age.

6 **Note:** The availability of rehabilitative and/or habilitative benefits for occupational therapy
 7 services, state and federal mandates, and regulatory requirements should be verified and
 8 followed in addition to the benefit plan provisions and medical necessity criteria defined
 9 in this document.

10 **2. REHABILITATIVE OCCUPATIONAL THERAPY SERVICES**

11 **Medically Necessary**

12 (1) Rehabilitative occupational therapy (OT) services to improve, adapt, compensate, or
 13 restore functions which have been impaired or permanently lost as a result of illness,
 14 injury, loss of a body part, or congenital abnormality are considered **medically**
 15 **necessary** when **ALL** the following criteria are met:

- 16 1. The services are delivered by a qualified provider of occupational therapy services
 17 (i.e., appropriately trained and licensed by the state to perform occupational therapy
 18 services); and
- 19 2. Rehabilitative occupational therapy occurs when the judgment, knowledge, and
 20 skills of a qualified provider of occupational therapy services (as defined by the
 21 scope of practice for therapists in each state) are necessary to safely and effectively
 22 furnish a recognized therapy service due to the complexity and sophistication of the
 23 plan of care and the medical condition of the individual, with the goal of
 24 improvement of an impairment or functional limitation.
- 25 3. The patient demonstrates a physical and/or functional impairment as demonstrated
 26 by the inability to perform basic activities of daily living (ADLs) or instrumental
 27 activities of daily living (IADLs), or usual daily activities.
- 28 4. The patient demonstrates signs and symptoms of physical and/or functional
 29 impairment in one or more of the following areas:
 30
 - 31 a. Sensory and/or motor
 - 32 b. Cognitive/psychological
 - 33 c. Cardiopulmonary status and circulation
 - 34 d. Skin
- 35 5. The patient’s condition has the potential to improve or is improving in response to
 36 therapy, maximum improvement is yet to be attained; and there is an expectation
 37 that the anticipated improvement is attainable in a **reasonable and generally**
 38 **predictable period of time*** and will result in a clinically significant level of
 39 functional improvement; and

- 1 6. Improvement or restoration of function could not be reasonably expected as the
- 2 individual gradually resumes normal activities without the provision of skilled
- 3 rehabilitative services; and
- 4 7. The documentation objectively verifies progressive functional improvement over
- 5 specific time frames and clinically justifies the initiation of continuation of
- 6 rehabilitative services; and
- 7 8. The program is individualized, and there is documentation outlining quantifiable,
- 8 attainable treatment goals.

9
 10 ***Reasonable and predictable period of time:** The specific time frames in which one
 11 would expect practical functional improvement is dependent on various factors
 12 including whether the services are Rehabilitative or Habilitative services. A reasonable
 13 trial of care for rehabilitative services to determine the patient’s potential for
 14 improvement in or restoration of function is influenced by the diagnosis; clinical
 15 evaluation findings; stage of the condition (acute, sub-acute, chronic); severity of the
 16 condition; and patient-specific elements (age, gender, past and current medical history,
 17 family history, and any relevant psychosocial factors). Habilitative services may be
 18 prolonged and are primarily influenced by the type of ADLs or IADLs which have not
 19 developed, or which are at risk of being lost.

20
 21 (2) An occupational therapy evaluation is considered medically necessary for the
 22 assessment of a physical impairment.

23
 24 **Not Medically Necessary**

- 25 (1) Rehabilitative OT services are considered not medically necessary if any of the
 26 following is determined:
- 27 1. Rehabilitative services are NOT intended to improve, adapt, or restore functions
 - 28 which have been impaired or permanently lost as a result of illness, injury, loss of
 - 29 a body part, or congenital abnormality.
 - 30 2. Improvement or restoration of function could reasonably be expected to improve
 - 31 as the individual gradually resumes normal activities without the provision of
 - 32 skilled therapy services. For example:
 - 33 o A patient suffers a transient and easily reversible loss or reduction in function
 - 34 which could reasonably be expected to improve spontaneously as the patient
 - 35 gradually resumes normal activities.
 - 36 o A fully functional patient who develops temporary weakness from a brief period
 - 37 of bed rest following abdominal surgery.
 - 38 3. Therapy services that do not require the skills of a qualified provider of OT services.
 - 39 Examples include but are not limited to:

- 1 ○ General exercises (basic aerobic, strength, flexibility, or aquatic programs) to
2 promote overall fitness/conditioning.
- 3 ○ Services for the purpose of enhancing athletic or recreational sports
4 performance or for return to sport after injury or surgery.
- 5 ○ Massages and whirlpools for relaxation.
- 6 ○ General public education/instruction sessions.
- 7 ○ Repetitive gait or other activities and services that an individual can practice
8 independently and can be self-administered safely and effectively.
- 9 a) Activities that require only routine supervision and NOT the skilled services
10 of an occupational therapy provider.
- 11 b) When a home exercise program is sufficient and can be utilized to continue
12 therapy (examples of exceptions include but would not be limited to the
13 following: if patient has poor exercise technique that requires cueing and
14 feedback, lack of support at home if necessary for exercise program
15 completion, and/or cognitive impairment that doesn't allow the patient to
16 complete the exercise program).
- 17 4. The expectation does **not** exist that the service(s) will result in a clinically
18 significant improvement in the level of functioning within a reasonable and
19 predictable period of time (up to 4 weeks).
- 20 ○ If function could reasonably be expected to improve as the individual gradually
21 resumes normal activities, then the service is considered **not** medically
22 necessary.
- 23 ○ The patient's condition does not have the potential to improve or is not
24 improving in response to therapy; or would be insignificant relative to the extent
25 and duration of therapy required; and there is an expectation that further
26 improvement is NOT attainable.
- 27 ○ The documentation fails to objectively verify functional progress over a
28 reasonable period of time (up to 4 weeks).
- 29 ○ The patient has reached maximum therapeutic benefit.
- 30 5. A passive modality is **not** preparatory to other skilled treatment procedures or is
31 not necessary in order to provide other skilled treatment procedures safely and
32 effectively.
- 33 6. A passive modality has insufficient published evidence to support a clinically
34 meaningful physiologic effect on the target tissue or improve the potential for a
35 positive response to care for the condition being treated.
- 36 7. Reevaluations or assessments of a patient's status that are not separate and distinct
37 services from those work components included within occupational therapy
38 services provided.

- 1 8. Reevaluations or assessments of a patient’s status that are not necessary to continue
2 a course of therapy nor related to a new condition or exacerbation for which the
3 reevaluation will likely result in a change in the treatment plan.
- 4 9. The treatments/services are not supported by and are not performed in accordance
5 with peer-reviewed literature as documented in applicable ASH CPGs or other
6 literature accepted by ASH Clinical Quality committee.

1 (2) The following treatments/programs are not considered medically necessary because
 2 they are nonmedical, non-rehabilitative, educational, or training in nature. In addition,
 3 these treatments/programs, may be specifically excluded under many benefit plans:

- 4 • Back school
- 5 • Driving safety/driver training
- 6 • Vocational rehabilitation programs and any program or evaluation with the primary
 7 goal of returning an individual to work
- 8 • Work hardening programs
- 9 • Health and wellness intervention
- 10 • Education and achievement testing, including Intelligence Quotient (IQ) testing.
- 11 • Educational interventions (e.g., classroom environmental manipulation, academic
 12 skills training and parental training).
- 13 • Services provided within the school setting and duplicated in the rehabilitation
 14 setting.

15
 16 (3) Use of the any of the following treatments is unproven. Refer to *Techniques and*
 17 *Procedures Not Widely Supported as Evidence-Based (CPG 133)* and/or the specific
 18 guideline below for additional information.

- 19 1. Intensive model of constraint-induced movement therapy
- 20 2. Intensive Model of Therapy (IMOT) programs [*Intensive Model of Therapy (CPG*
 21 *286 - S)*]
- 22 3. Dry hydrotherapy/aqua massage/hydromassage
- 23 4. Non-invasive Interactive Neurostimulation (e.g., InterX®) [*Non-invasive*
 24 *Interactive Neurostimulation (e.g., InterX®) (CPG 277 – S)*]
- 25 5. Microcurrent Electrical Nerve Stimulation (MENS)
- 26 6. H-WAVE ® [*H-WAVE® Electrical Stimulation (CPG 269 – S)*]
- 27 7. Equestrian therapy (e.g., hippotherapy)
- 28 8. MEDEK Therapy [*MEDEK Therapy (CPG 276 – S)*]
- 29 9. The Interactive Metronome Program
- 30 10. Elastic therapeutic tape/taping (e.g., Kinesio™ tape, KT TAPE/KT TAPE PRO™,
 31 Spidertech™ tape) [*Strapping and Taping (CPG 143 – S)*]
- 32 11. Dry Needling [*Dry Needling (CPG 178 – S)*]
- 33 12. Laser therapy [*Laser Therapy (LT) (CPG 30 – S)*]

35 **3. MAINTENANCE OCCUPATIONAL THERAPY SERVICES**

36 According to the Centers for Medicare and Medicaid Services (CMS) guidelines, or when
 37 covered by private carriers, maintenance occupational therapy services are a covered
 38 benefit when skilled occupational therapy care is medically necessary to maintain
 39 functional status or to prevent or slow further deterioration in function. Unlike coverage

1 for rehabilitative therapy, coverage for maintenance therapy does not depend on the
 2 presence or absence of a patient's potential for improvement for therapy; the deciding
 3 factors are always whether the services are considered reasonable, effective treatments for
 4 the patient's condition and require the skills of a therapist.

5
 6 If the specialized skill, knowledge and judgment of a qualified occupational therapist are
 7 required to establish or design a maintenance program to maintain the patient's current
 8 condition or to prevent or slow further deterioration, **the establishment or design of a**
 9 **maintenance program is medically necessary.**

10
 11 If skilled occupational therapy services by a qualified occupational therapy therapist, or
 12 occupational therapy assistant under the supervision of a qualified therapist, are needed to
 13 instruct the patient or appropriate caregiver regarding the maintenance program, **such**
 14 **instruction is medically necessary.**

15
 16 If skilled occupational therapy services are needed for periodic reevaluations or
 17 reassessments of the maintenance program, such **periodic reevaluations or reassessments**
 18 **are medically necessary.**

19
 20 Once a maintenance program is designed or established, a maintenance program can
 21 generally be performed by the patient alone or with the assistance of family member,
 22 caregiver or unskilled personnel. In such situations, coverage is not medically necessary.
 23 The performance or delivery of the maintenance therapy program is considered medically
 24 necessary only when the documentation establishes that the following criteria has been
 25 met:

- 26
 27 1. The individualized assessment of a patient's clinical condition demonstrates
 28 that the specialized judgment, knowledge and skills of an occupational therapy
 29 practitioner (skilled care) are necessary for the performance of an effective
 30 maintenance program.
 31 2. When the needed therapy procedures required to maintain the patient's current
 32 function or to prevent or slow further deterioration are of such complexity and
 33 sophistication that the skills of a qualified occupational therapy practitioner (as
 34 defined by scope of practice in each state) are required to furnish the therapy
 35 procedure or
 36 3. The particular patient's special medical complications require the skills of a
 37 qualified occupational therapy practitioner to furnish a therapy service required
 38 to maintain the patient's current function or to prevent or slow further

deterioration, even if the skills of an occupational therapy practitioner are not ordinarily needed to perform such therapy procedures.

The plan of care must be developed by the physician, NPP (non-physician practitioner) or OT who will provide the OT services.

4. HABILITATIVE OCCUPATIONAL THERAPY SERVICES

Habilitative services may or may not be covered services. If the member’s contract excludes habilitative services, the contract prevails.

Medically Necessary

(1) Habilitative OT services are considered medically necessary when ALL the following criteria are met:

1. The therapy is intended to maintain or develop skills needed to perform Activities of Daily Living (ADLs) or Instrumental Activities of Daily Living (IADLs) which have not (but normally would have) developed or which are at risk of being lost as a result of illness (including developmental delay), injury, loss of a body part, or congenital abnormality.
2. The occupational therapy services are evidence-based and require the judgment, knowledge, and skills of a qualified provider of occupational therapy services due to the complexity and sophistication of the plan of care and the medical condition of the individual.
3. There is an expectation that the therapy will assist development of function or maintain an acceptable level of functioning.
4. An individual would either not be expected to develop the function or would be expected to permanently lose the function (not merely experience fluctuation in the function) without the habilitative service. If the undeveloped or impaired function is not the result of a loss of body part or injury, a physician experienced in the evaluation and management of the undeveloped or impaired function has confirmed that the function would not either be expected to develop or would be permanently lost without the habilitative service. This information also concurs with the written treatment plan, which is likely to result in meaningful development of function or prevention of the loss of function.
5. There is a written treatment plan documenting the short and long-term goals (including estimated time when goals will be met) of treatment, frequency and duration of treatment, and what quantitative outcome measures will be used to assess function objectively.
6. Documentation objectively verifies that, at a minimum, functional status is maintained or developed.
7. The services are delivered by a qualified provider of occupational therapy services.

1
2
3
4

Not Medically Necessary

(1) Habilitative OT services are considered not medically necessary if any of the criteria above are not met.

5. REDUNDANT THERAPEUTIC EFFECTS AND REHABILITATIVE OR HABILITATIVE SERVICES

1. Redundant rehabilitative or habilitative therapy services expected to achieve the same therapeutic goal are considered not medically necessary and it would be inappropriate to provide these services to the same body region during the same treatment session. This includes treatments, such as but not limited to:
 - multiple modalities procedures that have similar or overlapping physiologic effects (e.g., multiple forms of superficial or deep heating modalities).
 - massage therapy and myofascial release.
 - orthotics training and prosthetic training.
 - whirlpool and Hubbard tank.
2. Duplicative (same or similar) rehabilitative or habilitative therapy services provided as part of an authorized therapy program through another therapy discipline are not medically necessary and inappropriate in the provision of care for the same patient.
 - When individuals receive physical, occupational, or speech therapy, the therapists should provide different treatments that reflect each therapy discipline's unique perspective on the individual's impairments and functional deficits and not duplicate the same treatment. They must also have separate evaluations, treatment plans, and goals. This applies to chiropractic services as well.
 - As an example, when individuals receive manual therapy services from an occupational therapist and chiropractic or osteopathic manipulation, the services must be documented as separate and distinct, performed on different body parts, and must be justified and non-duplicative.

6. THERAPEUTIC MODALITIES AND PROCEDURES

In some states, occupational therapists are required to hold a specific certification to use modalities in practice. The CPT codebook defines a modality as "any physical agent applied to produce therapeutic changes to biologic tissue; includes but is not limited to thermal, acoustic, light, mechanical, or electric energy." Modalities may be supervised, which means that the application of the modality doesn't require direct one-on-one patient contact by the practitioner. Or modalities may involve constant attendance, which indicates that the modality requires direct one-on-one patient contact by the practitioner.

Supervised modalities are untimed therapies. Untimed therapies are usually reported only once for each date of service regardless of the number of minutes spent providing this service or the number of body areas to which they were applied. Untimed services billed

1 as more than one unit will require significant documentation to justify treatment greater
2 than one session per day. Examples of supervised modalities include application of:

- 3 • Hot or cold packs
- 4 • Mechanical traction
- 5 • Unattended electrical stimulation
- 6 • Vasopneumatic devices
- 7 • Whirlpool
- 8 • Paraffin bath
- 9 • Diathermy

10
11 Modalities that require constant attendance, are timed and reported in 15-minute
12 increments (one unit) regardless of the number of body areas to which they are applied.
13 Examples of modalities that require constant attendance include:

- 14 • Contrast baths
- 15 • Ultrasound
- 16 • Electrical stimulation
- 17 • Iontophoresis

18
19 The CPT codebook defines therapeutic procedures as "A manner of effecting change
20 through the application of clinical skills and/or services that attempt to improve function."
21 Except for Group Therapy (97150) and Work Hardening/Conditioning (97545-6),
22 therapeutic procedures require direct (one-on-one) patient contact by the Occupational
23 Therapist, are timed therapies, and must be reported in units of 15-minute increments. Only
24 the actual time that the Occupational Therapist is directly working with the patient
25 performing exercises/activities, instruction, or assessments is counted as treatment time.
26 The time that the patient spends not being treated because of a need for rest or equipment
27 set up is not considered treatment time. Any exercise/activity that does not require, or no
28 longer requires, the skilled assessment and intervention of a health care practitioner is not
29 considered a medically necessary therapeutic procedure. Exercises often can be taught to
30 the patient or a caregiver as part of a home/self-care program. Examples of therapeutic
31 procedures that require the Occupational Therapist to have direct (one-on-one) patient
32 contact include:

- 33 • therapeutic exercises
- 34 • neuromuscular reeducation
- 35 • gait training
- 36 • manual therapy (e.g., soft tissue mobilization)
- 37 • therapeutic activities
- 38 • sensory integrative techniques

- 1
 - wheelchair training

1 **Documentation Requirements to Substantiate Medical Necessity of Therapeutic** 2 **Modalities and Procedures**

3 Proper and sufficient documentation is essential to establish the clinical necessity and
4 effectiveness of each modality and procedure, aid in the determination of patient outcomes
5 management, and support continuity of patient care. At a minimum, documentation is
6 required for every treatment day and for each therapy performed. Each daily record should
7 include: the date of service, the name of each modality and/or procedure performed, the
8 parameters for each modality (e.g., amperage/voltage, location of pads/electrodes), area of
9 treatment, total treatment time spent for each therapy (mandatory for timed services), the
10 total treatment time for each date of service, and the identity of the person(s) providing the
11 services. Failure to properly identify and sufficiently document the parameters for each
12 therapy on a daily progress note may result in an adverse determination (partial approval
13 or denial).

14 15 **6.1 Passive Care and Active Care**

16 Generally, passive modalities are used to manage the acute inflammatory response, pain,
17 and/or muscle tightness or spasm in the early stages of musculoskeletal and related
18 condition management. They are most effective during the acute phase of treatment. The
19 use of passive modalities in the treatment of sub-acute or chronic conditions beyond the
20 acute inflammatory response time frame is generally considered not medically necessary
21 unless there is an exacerbation. Passive modalities are rarely beneficial alone and are most
22 effective when performed as part of a comprehensive treatment approach. Some
23 improvement with the use of passive modalities should be seen within three visits. If
24 passive therapy is not contributing to improvement, passive therapy should be discontinued
25 and other evidence supported interventions implemented. The use of passive modalities is
26 generally considered not medically necessary unless they are preparatory and essential to
27 the safe and effective delivery of other skilled treatment procedures (e.g., therapeutic
28 exercise training). Prolonged reliance on passive modalities is not supported by the clinical
29 literature.

30
31 A “passive therapy” is a procedure applied by a clinical practitioner without active
32 engagement of or movement by the patient (e.g., ultrasound, hot packs).

33
34 The selection of a passive modality should be based on an understanding of the known
35 physiologic effects of the modality, contraindications, the stage of injury and/or tissue
36 healing, anatomical location to be treated, patient specific conditions and the likelihood of
37 the therapy to enhance recovery or facilitate treatment with manual and active therapeutic
38 procedures. Use of more than two (2) modalities on each visit date is unusual and should
39 be justified in the documentation.

1 Transition from passive physiotherapy modalities to active treatment procedures should be
 2 timely and evidenced in the medical record, including instructions on self/home care.
 3 Active therapeutic procedures are typically started as swelling, pain, and inflammation are
 4 reduced. Active care elements include increasing range of motion, strengthening primary
 5 and secondary stabilizers of a given region, and increasing the endurance capability of the
 6 muscles. Care focuses on active participation of the patient in their exercise program.
 7 Activities of Daily Living (ADLs) training, muscle strengthening, movement retraining,
 8 and progressive resistive exercises are considered active procedures. Patients should
 9 progress from active procedures requiring the supervision of a skilled practitioner to a self-
 10 directed home activity program as soon as possible.

11 **6.2 Treatment Interventions**

12 Below are descriptions and medical necessity criteria, as applicable, for different treatment
 13 interventions, including specific modalities and therapeutic procedures associated with
 14 occupational therapy. This material is for informational purposes only and is not indicative
 15 of coverage, nor is it an exhaustive list of services provided.

16 **Hydrotherapy/Whirlpool/Hubbard Tank**

17 These modalities involve supervised use of agitated water in order to relieve muscle spasm,
 18 improve circulation, or cleanse wounds e.g., ulcers, skin conditions. Hydrotherapy may be
 19 considered medically necessary for pain relief, muscle relaxation and improvement of
 20 movement for persons with musculoskeletal conditions or for wound care (cleansing and
 21 debridement).
 22
 23
 24

25 **Fluidotherapy®**

26 This modality is used specifically for acute and subacute conditions of the extremities.
 27 Fluidotherapy® is a dry superficial thermal modality that transfers heat to soft tissues by
 28 agitation of heated air and Cellux particles. The indications for this modality are similar to
 29 paraffin baths and whirlpool and it is an acceptable alternative to other heat modalities for
 30 reducing pain, edema, and muscle spasm from acute or subacute traumatic or non-traumatic
 31 musculoskeletal disorders of the extremities, including complex regional pain syndrome
 32 (CRPS). A benefit of Fluidotherapy® is that patients can perform active range of motion
 33 (AROM) while undergoing treatment.
 34

35 **Vasopneumatic Devices**

36 These special devices apply pressure for swelling/edema reduction, either after an acute
 37 injury, following a surgical procedure, due to lymphedema, or due to pathology such as
 38 venous insufficiency. Education sessions for home use are considered medically necessary
 39 (up to two sessions). Cooling systems such as Game Ready® Systems, Cryocuff, Polar Care

1 Wave or any similar cold compression system devices are not considered vasopneumatic
2 devices and should not be billed as such.

3 **Hot/Cold Packs**

4 Hot packs increase blood flow, relieve pain and increase flexibility. Cold packs decrease
5 blood flow to an area for reduction of pain and swelling. They may be considered medically
6 necessary for musculoskeletal conditions that include significant pain and/or swelling.

8 **Paraffin Bath**

9 This modality uses hot wax for application of heat. It is indicated for use to relieve pain
10 and increase range of motion of extremities (typically wrists and hands) due to chronic
11 joint problems, post-injury, or post-surgical scenarios.

13 **Infrared Light Therapy**

14 Infrared light therapy is a form of heat therapy used to increase circulation to relieve muscle
15 spasm. Other heating modalities are considered superior to infrared lamps and should be
16 considered unless there is a contraindication to those other forms of heat. Utilization of the
17 Infrared Light Therapy CPT code is not appropriate for low level laser treatment. This also
18 does not refer to Anodyne® Therapy System.

20 **Electrical Stimulation**

21 Electrical stimulation is used in different variations to relieve pain, reduce swelling, heal
22 wounds, and improve muscle function. Functional electric stimulation is considered
23 medically necessary for muscle re-education (to improve muscle contraction) in the earlier
24 phases of rehabilitation.

26 **Iontophoresis**

27 Electric current is used to transfer certain chemicals (medications) into body tissues. Use
28 of iontophoresis may be considered medically necessary for the treatment of inflammatory
29 conditions, such as plantar fasciitis and lateral epicondylitis.

31 **Contrast Baths**

32 This modality is the application of alternative hot and cold baths and is typically used to
33 treat extremities with subacute swelling or chronic regional pain syndrome (CRPS).
34 Contrast baths may be considered medically necessary to reduce hypersensitivity and
35 swelling.

37 **Ultrasound**

38 This modality provides deep heating through high frequency sound wave application. Non-
39 thermal applications are also possible using the pulsed option. Ultrasound is commonly

- 1 used to treat many soft tissue conditions that require deep heating or micromassage to a
- 2 localized area to relieve pain and improve healing. Ultrasound may be considered
- 3 medically necessary to relieve pain and improve healing.

1 **Diathermy (e.g., shortwave)**

2 Shortwave diathermy utilizes high frequency magnetic and electrical current to provide
3 deep heating to larger joints and soft tissue, and may be considered medically necessary
4 for pain relief, increased circulation, and muscle spasm reduction. Microwave diathermy
5 presents an unacceptable risk profile and is considered not medically necessary.

6
7 **Therapeutic Exercises**

8 Therapeutic exercise includes instruction, feedback, and supervision of a person in an
9 exercise program specific to their condition. Therapeutic exercise may be considered
10 medically necessary to restore/develop strength, endurance, range of motion and flexibility
11 which has been lost or limited as a result of a disease or injury. Exercise performed by the
12 patient within a clinic facility or other location (e.g., home; gym) without a physician or
13 therapist present and supervising would be considered not medically necessary.

14
15 **Neuromuscular Reeducation (NMR)**

16 NMR generally refers to a treatment technique performed for the purpose of retraining the
17 connection of the brain and muscles, via the nervous system, to improve movement,
18 strength, balance, and function. The goal of NMR is to develop conscious control of
19 individual muscles and awareness of position of extremities. The procedure may be
20 considered medically necessary for impairments which affect the neuromuscular system
21 (e.g., poor static or dynamic sitting/standing balance, loss of gross and fine motor
22 coordination) that may result from musculoskeletal or neuromuscular disease or injury such
23 as severe trauma to nervous system, post orthopedic surgery, cerebral vascular accident
24 and systemic neurological disease. Example techniques may include proprioceptive
25 neuromuscular facilitation (PNF), BAP's boards, vestibular rehabilitation, desensitization
26 techniques. This does not include contract/relax or other soft tissue massage techniques.
27 NMR is typically used as the precursor to the implementation of Therapeutic Activities.

28
29 **Aquatic Therapy**

30 Pool therapy (aquatic therapy) is provided individually, in a pool, to debilitated or
31 neurologically impaired individuals. (The term is not intended to refer to relatively normal
32 functioning individuals who exercise, swim laps or relax in a hot tub or Jacuzzi.) The goal
33 is to develop and/or maintain muscle strength and range of motion by reducing forces of
34 gravity through total or partial body immersion (except for head). Aquatic therapy may be
35 considered medically necessary to develop and/or maintain muscle strength and range of
36 motion when it is necessary to reduce the force of gravity through partial body immersion.

37
38 **Soft Tissue Mobilization**

1 Soft tissue mobilization techniques are more specific in nature and include, but are not
2 limited to, myofascial release techniques, friction massage, and trigger point techniques.
3 Specifically, myofascial release is a soft tissue manual technique that involves
4 manipulation of the muscle, fascia, and skin. Skilled manual techniques (active and/or
5 passive) are applied to soft tissue to effect changes in the soft tissues, articular structures,
6 neural or vascular systems. Examples are facilitation of fluid exchange, restoration of
7 movement in acutely edematous muscles, or stretching of shortened connective tissue. This
8 procedure is considered medically necessary for treatment of pain and restricted motion of
9 soft tissues resulting in functional deficits.

10 **Joint Mobilization**

11 Joint mobilization is utilized to reduce pain and increase joint mobility. Most often
12 mobilizations are indicated for the upper extremity, especially the hand.

14 **Therapeutic Activities**

15 Therapeutic activities or functional activities (e.g., bending, lifting, carrying, reaching,
16 pushing, pulling, stooping, catching and overhead activities may be considered medically
17 necessary) to improve function when there has been a loss or restriction of mobility,
18 strength, balance or coordination. These dynamic activities must be part of an active
19 treatment plan and directed at a specific outcome. As an example, this intervention may be
20 considered medically necessary after a patient has completed exercises focused on
21 strengthening and range of motion but needs to improve function-based activities.
22

23 **Activities of Daily Living (ADL) Training**

24 This procedure is considered medically necessary to enable the patient to perform essential
25 activities of daily living, instrumental activities of daily living, and self-care including
26 bathing, feeding, preparing meals, toileting, dressing, walking, making a bed, and
27 transferring from bed to chair, wheelchair or walker. Services provided concurrently by
28 physical therapists and occupational therapists may be considered medically necessary if
29 there are separate and distinct functional goals.
30

31 **Cognitive Skills Development**

32 This procedure is considered medically necessary for persons with acquired cognitive
33 deficits resulting from head trauma, or acute neurologic events including cerebrovascular
34 accident, pediatric developmental condition, or other situations. It is not appropriate for
35 persons without potential for improvement. Occupational therapists and speech language
36 pathologists with specific training typically provide this care. This procedure should be
37 aimed at improving or restoring specific functions which were impaired by an identified
38 illness or injury.
39

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

Sensory Integration

Sensory integration involves perceiving, modulating, organizing, and interpreting these sensations to optimize occupational performance and participation. Sensory integration is mainly an intervention for children with developmental and behavioral disorders. The activities included in SI provide vestibular, proprioceptive, auditory, and tactile stimuli, which in turn organize the sensory system. See CPG 149 Sensory Integrative (SI) Therapy for medical necessity criteria.

Orthotic Management and Training

Orthotic management and training may be considered medically necessary when the documentation specifically demonstrates that the specific knowledge, skills, and judgment of an Occupational Therapist are required to train the patient in the proper use of braces and/or splints (orthotics). Many braces or splints do not require specific training by the Occupational Therapist in their use and can be safely procured and applied by the patient. Patients with cognitive, dexterity, or other significant deficits may need specific training where other patients do not.

Prosthetic Training

Prosthetic training may be considered medically necessary when the professional skills of the practitioner are required to train the patient in the proper fitting and use of a prosthetic (an artificial body part, such as a limb). Periodic return visits beyond the third month may be necessary.

Wheelchair Management Training

This procedure is considered medically necessary only when it is part of an active treatment plan directed at a specific goal. The member must have the capacity to learn from instructions. Typically, three (3) sessions are adequate.

Active Wound Care Management

The CPT codebook defines active wound care procedures as those procedures "performed to remove devitalized tissue and/or necrotic tissue and promote healing" (AMA, current year). The practitioner is required to have direct one-on-one contact with the patient. Examples of active wound care management include non-selective debridement of an open wound, including topical application; use of whirlpool or other modalities; and negative pressure wound therapy. Occupational therapy state rules and regulations will dictate if occupational therapists can perform wound care.

Lymphedema Management

For more information, see the *Lymphedema (CPG 157 – S)* clinical practice guideline.

6.3 Precautions and Contraindications to Therapeutic Modalities and Procedures

1. The use of thermotherapy is contraindicated for the following:

- Recent or potential hemorrhage
- Thrombophlebitis
- Impaired sensation
- Impaired mentation
- Malignant tumor
- IR irradiation of the eyes

Precautions for use of thermotherapy include:

- Acute injury or inflammation
- Pregnancy
- Impaired circulation
- Poor thermal regulation
- Edema
- Cardiac insufficiency
- Metal in the area
- Over an open wound
- Over areas where topical counterirritants have recently been applied
- Demyelinated nerve

2. The use of cryotherapy is contraindicated for the following:

- Cold hypersensitivity
- Cold intolerance
- Cryoglobulinemia
- Paroxysmal cold hemoglobinuria
- Raynaud disease or phenomenon
- Over regenerating peripheral nerves
- Over an area with circulatory compromise or peripheral vascular disease

Precautions for cryotherapy include:

- Over the superficial branch of a nerve
- Over an open wound
- Hypertension
- Poor sensation or mentation

1 3. The use of immersion hydrotherapy is contraindicated for the following:

- 2 • Cardiac instability
- 3 • Confusion or impaired cognition
- 4 • Maceration around a wound
- 5 • Bleeding
- 6 • Infection in the area to be immersed
- 7 • Bowel incontinence
- 8 • Severe epilepsy
- 9 • Suicidal patients

10
11 Precautions for full body immersion in hot or very warm water include:

- 12 • Pregnancy
- 13 • Multiple Sclerosis
- 14 • Poor thermal regulation

15
16 4. Contraindications for Traction include:

- 17 • Where motion is contraindicated
- 18 • Acute injury or inflammation
- 19 • Joint hypermobility or instability
- 20 • Peripheralization of symptoms with traction
- 21 • Uncontrolled hypertension

22
23 Precautions for Traction include:

- 24 • Structural diseases or conditions affecting the tissues in the area to be treated
25 (e.g., tumor, infection, osteoporosis, RA, prolonged systemic steroid use, local
26 radiation therapy)
- 27 • When pressure of the belts may be hazardous (e.g., with pregnancy, hiatal
28 hernia, vascular compromise, osteoporosis)
- 29 • Displaced annular fragment
- 30 • Medial disc protrusion
- 31 • When severe pain fully resolves with traction
- 32 • Claustrophobia or other psychological aversion to traction
- 33 • Inability to tolerate prone or supine position
- 34 • Disorientation

35
36 Additional precautions for cervical traction:

- 37 • TMJ problems
- 38 • Dentures

1 5. The use of thermal shortwave diathermy (SWD) is contraindicated for the following

- 2 • Any metal in the treatment area or on/in the body.
- 3 • Malignancy
- 4 • Eyes
- 5 • Testes
- 6 • Growing epiphyses

7
8 Contraindications for all forms of SWD:

- 9 • Implanted or transcutaneous neural stimulators including cardiac pacemakers
- 10 • Pregnancy

11
12 Precautions for all forms of SWD:

- 13 • Near electronic or magnetic equipment
- 14 • Obesity
- 15 • Copper-bearing intrauterine contraceptive devices

16
17 6. Contraindications for use of Electrical Currents:

- 18 • Demand pacemakers, implantable defibrillator, or unstable arrhythmia
- 19 • Placement of electrodes over carotid sinus
- 20 • Areas where venous or arterial thrombosis or thrombophlebitis is present
- 21 • Pregnancy – over or around the abdomen or low back

22
23 Precautions for electrical current use:

- 24 • Cardiac disease
- 25 • Impaired mentation
- 26 • Impaired sensation
- 27 • Malignant tumors
- 28 • Areas of skin irritation or open wounds

29
30 7. Contraindications to the use of ultrasound include:

- 31 • Malignant tumor
- 32 • Pregnancy
- 33 • Central Nervous Tissue
- 34 • Joint cement
- 35 • Plastic components
- 36 • Pacemaker or implantable cardiac rhythm device
- 37 • Thrombophlebitis
- 38 • Eyes

- 1 • Reproductive organs
2 Precautions for Ultrasound include:
3 • Acute inflammation
4 • Epiphyseal plates
5 • Fractures
6 • Breast implants
7

8 The use of electrical muscle stimulation, SWD, thermotherapy, cryotherapy, ultrasound,
9 laser/light therapy, immersion hydrotherapy, and mechanical traction with pediatric
10 patients is contraindicated if the patient cannot provide the proper feedback necessary for
11 safe application.
12

13 In addition to the contraindications listed above, there are a wide range of services which
14 are considered unproven, pose a significant health and safety risk, are scientifically
15 implausible and/or are not widely supported as evidence based. Such services would be
16 considered not medically necessary and include, but are not limited to:

- 17 • Axial/Spinal decompression
18 • Dry needling
19 • Laser therapy
20 • Manual muscle testing to diagnosis non-neuromusculoskeletal conditions
21 • Microcurrent Electrical Nerve Stimulation (MENS)
22 • Other unproven procedures (see the *Techniques and Procedures Not Widely*
23 *Supported as Evidence-Based (CPG 133 – S)* clinical practice guideline for
24 complete list)
25

26 7. CLINICAL DOCUMENTATION

27 Medical record keeping is an essential component of patient evaluation and management.
28 Medical records should be legible and should contain, at a minimum sufficient information
29 to identify the patient, support the diagnosis, justify the treatment, accurately document the
30 results, indicate advice and cautionary warnings provided to the patient and provide
31 sufficient information for another practitioner to assume continuity of the patient's care at
32 any point in the course of treatment. Good medical record keeping improves the likelihood
33 of a positive outcome and reduces the risk of treatment errors. It also provides a resource
34 to review cases for opportunities to improve care, provides evidence for legal records, and
35 offers necessary information for third parties who need to review and understand the
36 rationale and type of services rendered (e.g., medical billers and auditors/reviewers).
37

38 Outcome measures are important in determining effectiveness of a patient's care. The use
39 of standardized tests and measures early in an episode of care establishes the baseline status

1 of the patient, providing a means to quantify change in the patient's functioning. Outcome
 2 measures provide information about whether predicted outcomes are being realized. When
 3 comparison of follow-up with baseline outcome metrics does not demonstrate minimal
 4 clinically important difference (MCID) (minimal amount of change in a score of a valid
 5 outcome assessment tool) the treatment plan should be changed or be discontinued. Failure
 6 to use Functional Outcome Measures (FOMs) / Outcome Assessment Tools (OATs) may
 7 result in insufficient documentation of patient progress and may result in an adverse
 8 determination (partial approval or denial) of continued care.

10 **7.1 Evaluation and Re-evaluations**

11 The initial evaluation is usually completed in a single session. The initial evaluation should
 12 document the necessity of a course of therapy through objective findings and subjective
 13 patient/caregiver self-reporting. Initial evaluations are completed to determine the medical
 14 necessity of initiating rehabilitative therapy or skilled instruction in maintenance activities
 15 that the patient and/or caregiver can perform at home. The occupational therapist performs
 16 an initial examination and evaluation to establish a working diagnosis, prognosis, and plan
 17 of care prior to intervention. Determination of referral to another health care practitioner is
 18 also an essential part of an initial evaluation. An initial evaluation for a new condition by
 19 an Occupational Therapist is defined as the evaluation of a patient:

- 20 • For which this is their first encounter with the practitioner or practitioner group;
- 21 • Who presents with:
 - 22 ○ A new injury or new condition; or
 - 23 ○ The same or similar complaint after discharge from previous care.
- 24 • Choice of code is dependent upon the level of complexity.

25
 26 The evaluation codes reflect three (3) levels of patient presentation:

- 27 1. low complexity;
- 28 2. moderate complexity; and
- 29 3. high complexity.

30
 31 Four (4) components are used to select the appropriate occupational therapy evaluation
 32 CPT code. These include:

- 33 1. Occupational profile and client history (medical and therapy);
- 34 2. Assessments of occupational performance;
- 35 3. Clinical decision making;
- 36 4. Development of plan of care.

37
 38 Relevant CPT Codes: 97165, 97166, and 97167 – Occupational Therapy evaluation.

1 The occupational therapist evaluation:

- 2 • Is documented, dated, and appropriately authenticated by the occupational therapist
- 3 who performed it
- 4 • Identifies the occupational therapy needs of the patient
- 5 • Incorporates appropriate tests and measures to facilitate outcome measurement
- 6 • Produces data that are sufficient to allow evaluation, prognosis, and the
- 7 establishment of a plan of care

8
9 The written plan of care should be sufficient to determine the medical necessity of
10 treatment, including:

- 11 • The diagnosis along with the date of onset or exacerbation of the disorder/diagnosis
- 12 • A reasonable estimate of when the goals will be reached
- 13 • Long-term and short-term goals that are specific, quantitative and objective
- 14 • Occupational therapy evaluation pertinent findings
- 15 • The frequency and duration of treatment
- 16 • Rehabilitation or habilitation prognosis
- 17 • The specific treatment techniques and/or exercises to be used in treatment
- 18 • Signatures of the patient's occupational therapist

19
20 Re-evaluations are distinct from therapy assessments. There are several routine
21 reassessments that are not considered re-evaluations. These include ongoing reassessments
22 that are part of each skilled treatment session, progress reports, and discharge summaries.
23 Re-evaluation provides additional objective information not included in documentation of
24 ongoing assessments, treatment or progress notes. Assessments are considered a routine
25 aspect of intervention and are not billed separately from the intervention. Continuous
26 assessment of the patient's progress is a component of the ongoing therapy services and is
27 not payable as a re-evaluation.

28
29 Re-evaluation services are considered medically necessary when **ALL** of the following
30 conditions are met:

- 31 • Re-evaluation is not a recurring routine assessment of patient status;
- 32 • The documentation of the re-evaluation includes all of the following elements:
 - 33 ○ An evaluation of progress toward current goals;
 - 34 ○ Making a professional judgment about continued care;
 - 35 ○ Making a professional judgment about revising goals and/or treatment or
 - 36 terminating services.

37 **AND the following indication is documented:**

38 An exacerbation or significant change in patient/client status or condition.

39

1 Relevant CPT Codes: 97168 – Occupational Therapy re-evaluation
 2 In order to reflect that continued OT services are medically necessary, intermittent progress
 3 reports must demonstrate that the individual is making functional progress.
 4

5 **7.2 Treatment Sessions**

6 An occupational therapy intervention is the purposeful interaction of the occupational
 7 therapy practitioner (OT or OTA) with the patient and, when appropriate, with other
 8 individuals involved in patient care, using various occupational therapy procedures and
 9 techniques to produce changes in the condition that are consistent with the diagnosis and
 10 prognosis. Occupational therapy interventions consist of coordination, communication,
 11 and documentation; patient-related and family/caregiver instruction; and procedural
 12 interventions. Occupational therapists aim to alleviate impairment and functional limitation
 13 by designing, implementing, and modifying therapeutic interventions. An occupational
 14 therapy session can vary in duration; however, treatment sessions lasting more than one
 15 hour per day are infrequent in outpatient settings (payor medical or reimbursement
 16 coverage policy may limit unit or session duration per date of service). Treatment sessions
 17 for more than one hour per day may be medically appropriate but must be supported in the
 18 documented plan of care and based on a patient's medical condition. An occupational
 19 therapy session may include:

- 20 • Evaluation or reevaluation
- 21 • Therapeutic use of everyday life and other purposeful activities, and other
 22 interventions focusing on preparing patients for daily activities performed in life
 23 and work
- 24 • Basic and advanced functional training in daily living, self-care and home
 25 management including activities of daily living (ADL) and instrumental activities
 26 of daily living (IADL)
- 27 • Management of feeding, eating and swallowing to improve eating and feeding
 28 performance
- 29 • Cognitive, perceptual, safety and judgment evaluation and training
- 30 • Adaptive training in and modification of activities, processes and environments
 31 (home, work, school, or community), including ergonomic applications and
 32 performance improvement
- 33 • Assessment, design, fabrication, application, fitting, and training in assistive
 34 technology, adaptive devices, and orthotic devices
- 35 • Training in the use of prosthetic devices
- 36 • Higher level independent living skill instruction and community/work functional
 37 reintegration
- 38 • Functionally oriented upper extremity interventions

- 1 • Training of the patient, caregivers, and family/parents in home exercise and activity
- 2 programs
- 3 • Skilled reassessment of the individual's problems, plan, and goals as part of the
- 4 treatment session

5 Documentation of each treatment session should include at a minimum:

- 6 • Date of treatment;
- 7 • Subjective complaints and current status (including functional deficits and ADL
- 8 restrictions);
- 9 • Description/name of each specific treatment intervention provided that match the
- 10 CPT codes billed, including;
 - 11 ○ Treatment time for each modality or procedure performed
 - 12 ○ Parameters of any modality or procedure, (e.g., voltage/amperage,
 - 13 pad/electrode placement, area of treatment, types of exercises/activities, and
 - 14 intended goal of each therapy)
- 15 • The patient's response to each service and to the entire treatment session;
- 16 • Any progress toward the goals in objective, measurable terms using consistent and
- 17 comparable methods;
- 18 • Any changes to the plan of care;
- 19 • Recommendations for follow-up visit(s);
- 20 • Signature/electronic identifier, name and credentials of the treating clinician.

21 **7.3 Discharge/Discontinuation of Intervention**

22 The occupational therapist discharges the patient from occupational therapy services when

23 the anticipated goals or expected outcomes for the patient have been achieved. The

24 occupational therapist discontinues intervention when the patient is unable to continue to

25 progress toward goals or when the occupational therapist determines that the patient will

26 no longer benefit from occupational therapy.

27

28

29 The occupational therapy discharge documentation includes:

- 30 • The status of the patient at discharge and the goals and outcomes attained
- 31 • Appropriate date and authentication by the occupational therapist who performed
- 32 the discharge
- 33 • When a patient is discharged prior to attainment of goals and outcomes, the status
- 34 of the patient and the rationale for discontinuation
- 35 • Initial, subsequent, and final FOMs scores
- 36 • Proposed self-care recommendations, if applicable
- 37 • Referrals to other health care practitioners/referring physicians as appropriate

- 1 • If the patient self- discharges, documentation of final status and if known, the
- 2 reason for discontinuation of services.

1 **7.4 Duplicated / Insufficient Information**

2 (1) Entries in the medical record should be contemporaneous, individualized, appropriately
 3 comprehensive, and made in a chronological, systematic, and organized manner.
 4 Duplicated/nearly duplicated medical records (a.k.a. cloned records) are not acceptable. It
 5 is not clinically reasonable or physiologically feasible that a patient's condition will be
 6 identical on multiple encounters. (Should the finding be identical for encounters, it would
 7 be expected that treatment would end because patient is not making progress toward current
 8 goals.)

9
 10 This includes, but not limited to:

- 11 • duplication of information from one treatment session to another (for the same or
 12 different patient[s]);
- 13 • duplication of information from one evaluation to another (for the same or different
 14 patient[s]).

15
 16 Duplicated medical records do not meet professional standards of medical record keeping
 17 and may result in an adverse determination (partial approval or denial) of those services.

18
 19 (2) The use of a system of record keeping that does not provide sufficient information
 20 (e.g., checking boxes, circling items from lists, arrows, travel cards with only dates of visit
 21 and listings). These types of medical record keeping may result in an adverse determination
 22 (partial approval or denial) of those services.

23
 24 Effective and appropriate records keeping that meet professional standards of medical
 25 record keeping document with adequate detail a proper assessment of the patient's status,
 26 the nature and severity of his/her complaint(s) or condition(s), and/or other relevant clinical
 27 information (e.g., history, parameters of each therapy performed, objective findings,
 28 progress towards treatment goals, response to care, prognosis).

29 30 **7.5 Centers for Medicare and Medicaid Services (CMS)**

31 For Medicare and Medicaid services, medical records keeping must follow and be in
 32 accordance with Medicare and any additional state Medicaid required documentation
 33 guidelines.

34 35 **8. CLINICAL REVIEW PROCESS**

36 Medical necessity evaluations require approaching the clinical data and scientific evidence
 37 from a global perspective and synthesizing the various elements into a congruent picture
 38 of the patient's condition and need for skilled treatment intervention. Clinical review
 39 decisions made by the clinical quality evaluators are based upon the information provided

1 by the treating practitioner in the submitted documentation and other related findings and
 2 information. Failure to appropriately document pertinent clinical information may result in
 3 adverse determinations (partial approval or denial) of those services. Therefore, thorough
 4 documentation of all clinical information that established the diagnosis/diagnoses and
 5 supports the intended treatment is essential.

7 **8.1 Definition of Key Terminology used in Clinical Reviews**

8 **Elective/Convenience Services**

9 Examples of elective/convenience services include: (a) preventive services; (b) wellness
 10 services; (c) services not necessary to return the patient to pre-illness/pre-injury functional
 11 status and level of activity; (d) services provided after the patient has reached MTB.
 12 (Elective/convenience services may not be covered through specific client or ASH
 13 benefits.)

15 **Minimal Clinically Important Difference (MCID)**

16 The MCID is the minimal amount of change in a score of a valid outcome assessment tool
 17 that indicates an actual improvement in the patient's function or pain. Actual significance
 18 of outcome assessment tool findings requires correlation with the overall clinical
 19 presentation, including updated subjective and objective examination/evaluation findings.

21 **Maximum Therapeutic Benefit (MTB)**

22 MTB is the patient's health status when the application of skilled therapeutic services has
 23 achieved its full potential (which may or may not be the complete resolution of the patient's
 24 condition.) At the point of MTB, continuation of the same or similar skilled treatment
 25 approach will not significantly improve the patient's impairments and function during this
 26 episode of care.

28 If the patient continues to have significant complaints, impairments, and documented
 29 functional limitations, one should consider the following:

- 30 • Altering the treatment regimen such as utilizing a different physiological approach
 31 to the treatment of the condition, or decreasing the use of passive care (modalities,
 32 massage etc.) and increasing the active care (therapeutic exercise) aspects of
 33 treatment to attain greater functional gains;
- 34 • Reviewing self-management program including home exercise programs; and/or
- 35 • Referring the patient for consultation by another health care practitioner for
 36 possible co-management or a different therapeutic approach.

38 **Preventive Services**

1 Preventive services are designed to reduce the incidence or prevalence of illness,
 2 impairment, and risk factors, and to promote optimal health, wellness, and function. These
 3 services are not designed or performed to treat or manage a specific health condition.
 4 (Preventive services may or may not be covered under specific clients or through ASH
 5 benefits.)

6 **Acute**

7 The stage of an injury, illness, or disease, in which the presence of clinical signs and
 8 symptoms is less than six weeks in duration, typically characterized by the presence of one
 9 or more signs of inflammation or other adaptive response.

10
 11 **Sub-Acute**

12 The stage of an injury, illness, or disease, in which the presence of clinical signs and
 13 symptoms is greater than six weeks, but not greater than twelve weeks in duration.

14
 15 **Chronic**

16 The stage of an injury, illness, or disease, in which the presence of clinical signs and
 17 symptoms is greater than twelve weeks in duration.

18
 19 **Red Flag(s)**

20 Signs and symptoms presented through history or examination/assessment that warrant
 21 more detailed and immediate medical assessment and/or intervention.

22
 23 **Yellow Flag(s)**

24 Adverse prognostic indicators with a psychosocial predominance associated with chronic
 25 pain and disability. Yellow flags signal the potential need for more intensive and complex
 26 treatment and/or earlier specialist referral.

27
 28 **Co-Morbid Condition(s)**

29 The presence of a concomitant condition, that has an unrelated pathology or disease
 30 process, but may inhibit, lengthen, or alter in some way the expected response to care.

31
 32 **8.2 Clinical Quality Evaluation**

33 The goal of the clinical quality evaluators during the review and decision-making process
 34 is to approve, as appropriate, those clinical services necessary to return the patient to pre-
 35 clinical/pre-morbid health status or stabilize a chronic condition, as supported by the
 36 documentation presented. The clinical quality evaluator is to evaluate if the documentation
 37 and other clinical information presented by the treating provider has appropriately
 38 substantiated the patient's condition and appropriately justifies the treatment plan that is
 39 presented.

Approval

ASH clinical quality evaluators have the responsibility to approve appropriate care for all services that are medically necessary. The clinical quality evaluators assess the clinical data supplied by the practitioner in order to determine whether submitted services and/or the initiation or continuation of care has been documented as medically necessary. The practitioner is accountable to document the medical necessity of all services submitted/provided. It is the responsibility of the peer clinical quality evaluator to evaluate the documentation in accordance with their training, understanding of practice parameters, and review criteria adopted by ASH through its clinical committees.

The following items influence clinical service approvals:

- No evidence of contraindication(s) to services submitted for review;
- Complaints, exam findings, and diagnoses correlate with each other;
- Treatment Plan is supported by the nature and severity of complaints;
- Treatment Plan is supported by exam findings;
- Treatment Plan is expected to improve symptoms (e.g., pain, function) within a reasonable period of time;
- Maximum therapeutic benefit has not been reached;
- Treatment Plan requires the skills of the provider; and
- Demonstration of progression toward active home/self-care and discharge.

Partial Approval

Occurs when only a portion of the submitted services are determined to be medically necessary services. The partial approval may refer to a decrease in treatment frequency, treatment duration, number of Durable Medical Equipment (DME)/supplies/appliances, number of therapies, or other services from the original amount/length submitted for review. This decision may be due to any number of reasons, such as:

- the practitioner's documentation of the history and exam findings are inconsistent with the clinical conclusion(s)
- the treatment dosage (frequency/duration) submitted for review is not supported by the underlying diagnostic or clinical features
- the need to initiate only a limited episode of care in order to monitor the patient's response to care

Additional services may be submitted and reviewed for evaluation of the patient's response to the initial trial of care. If the practitioner or patient disagrees with the partial approval of services, they contact the clinical quality evaluator listed on their response form to discuss

1 the case, submit additional documentation through the Reopen process, or submit
 2 additional documentation to appeal the decision through the Provider Appeals and Member
 3 Grievances process.

4 **Non-approval / Denial**

6 Occurs when none of the services submitted for review are determined to be medically
 7 necessary services. The most common causes for a non-approval/denial of all services are
 8 administrative or contractual in nature (e.g., ineligibility, reached plan benefit limits, non-
 9 coverage). Clinically, it is appropriate to deny continued/ongoing care if the patient's
 10 condition(s) are not, or are no longer, responding favorably to the services being rendered
 11 by the treating practitioner, or the patient has reached maximum therapeutic benefit.

13 **Additional / Continued Care**

14 Approval of additional treatment/services requires submission of additional information,
 15 including the patient's response to care and updated clinical findings. In cases where an
 16 additional course of care is submitted, the decision to approve additional services will be
 17 based upon the following criteria:

- 18 • The patient has made clinically significant progress under the initial treatment
 19 plan/program based on a reliable and valid outcome tool or updated subjective and
 20 objective examination findings.
- 21 • Additional clinically significant progress can be reasonably expected by continued
 22 treatment (the patient has not reached MTB or maximum medical improvement).
- 23 • There is no indication that immediate care/evaluation is required by other health
 24 care professionals.

26 Any exacerbation or flare-up of the condition that contributes to the need for additional
 27 treatment/services must be clearly documented.

29 Ancillary diagnostic procedures should be selected based on clinical history and
 30 examination findings that suggest the necessity to rule out underlying pathology or to
 31 confirm a diagnosis that cannot be verified through less invasive methods.

- 32 • Information is expected to directly impact the treatment/services and course of care.
- 33 • The benefit of the procedure outweighs the risk to the patient's health (short and
 34 long term).
- 35 • The procedure is sensitive and specific for the condition being evaluated (e.g., an
 36 appropriate procedure is utilized to evaluate for pathology).

38 The clinical information that the clinical quality evaluator expects to see when evaluating
 39 the documentation in support of the medical necessity of submitted treatment/services

1 should be commensurate with the nature and severity of the presenting complaint(s) and
2 scope of the practitioner of services and may include but is not limited to:

- 3 • History
- 4 • Physical Examination/Evaluation
- 5 • Documented Treatment Plan and Goals
- 6 • Estimated time of Discharge

7
8 In general, the initiation of care is warranted if there are no contraindications to prescribed
9 care, there is reasonable evidence to suggest the efficacy of the prescribed intervention,
10 and the intervention is within the scope of services permitted by State or Federal law. The
11 treatment submission for a disorder is typically structured in time-limited increments
12 depending on clinical presentation. Dosage (frequency and duration of service) should be
13 appropriately correlated with clinical findings, potential complications/barriers to recovery
14 and clinical evidence. When the practitioner discovers that a patient is nonresponsive to
15 the applied interventions within a reasonable time frame, re-assessment and treatment
16 modification should be implemented and documented. If the patient's condition(s) worsen,
17 the practitioner should take immediate and appropriate action to discontinue or modify care
18 and/or make an appropriate healthcare referral.

19
20 Services that do not require the professional skills of a practitioner to perform or supervise
21 are not medically necessary., If a patient's recovery can proceed safely and effectively
22 through a home exercise program or self-management program, services are not indicated
23 or medically necessary.

24 **8.3 Critical Factors during Clinical Reviews**

25
26 The complexity and/or severity of historical factors, symptoms, examination findings, and
27 functional deficits play an essential role to help quantify the patient's clinical status and
28 assess the effectiveness of planned interventions over time. Clinical quality evaluators
29 consider patient-specific variables as part of the medical necessity verification process. The
30 entire clinical picture must be taken into consideration with each case evaluated based upon
31 unique patient and condition characteristics.

32
33 Such variables may include, but not be limited to co-morbid conditions and other barriers
34 to recovery, the stage(s) of the condition(s), mechanism of injury, severity of the
35 symptoms, functional deficits, and exam findings, as well as social and psychological status
36 of the patient and the available support systems for self-care. In addition, the patient's age,
37 symptom severity, and the extent of positive clinical findings may influence duration,
38 intensity, and frequency of services approved as medically necessary. For example:

- 1 • Severe symptomatology, exam findings, and/or functional deficits may require
2 more care overall (e.g., longer duration, more services per encounter, and frequency
3 of encounters that the average); these patients require a higher frequency; but may
4 require short-term trials of care initially to assess patient response to care.
- 5 • Less severe symptomatology, exam findings and/or functional deficits usually
6 require less care (e.g., shorter duration, fewer services per encounter, and frequency
7 of encounters that the average); overall but may allow for less oversight and a
8 longer initial trial of care.
- 9 • As patients age, they may have a slower response to care, and this may affect the
10 approval of a trial of care.
- 11 • Because pediatric patients (under the age of 12) have not reached musculoskeletal
12 maturity, it may be necessary to modify the types of therapies approved as well as
13 shorten the initial trial of care.
- 14 • Complicating and/or co-morbid condition factors vary depending upon individual
15 patient characteristics, the nature of the condition/complaints, historical and
16 examination elements, and may require appropriate coordination of care and/or
17 more timely re-evaluation.

18
19 The following are examples of the factors clinical quality evaluators consider when
20 verifying the medical necessity of rehabilitative services for musculoskeletal conditions
21 and pain disorders.

22 **8.3.1 General Factors**

23
24 Multiple patient-specific historical and clinical findings may influence clinical decisions,
25 such as but not limited to:

- 26 • Red Flags
- 27 • Yellow Flags (Psychosocial Factors)
- 28 • Co-morbid conditions (e.g., diabetes, inflammatory conditions, joint instability)
- 29 • Age (older or younger)
- 30 • Non-compliance with treatment and/or self-care recommendations
- 31 • Lack of response to appropriate care
- 32 • Lifestyle factors (e.g., smoking, diet, stress, deconditioning)
- 33 • Work and recreational activities
- 34 • Pre-operative/post-operative care
- 35 • Medication use (type and compliance)

36
37 Nature of Complaint(s)

- 38 • Acute and severe symptoms

- 1 • Functional testing results that display severe disability/dysfunction
- 2 • Pain that radiates below the knee or elbow (for spinal conditions)

3

4 History

- 5 • Trauma resulting in significant injury or functional deficits.
- 6 • Pre-existing pathologies/surgery(ies)
- 7 • Congenital anomalies (e.g., severe scoliosis)
- 8 • Recurring exacerbations
- 9 • Prior episodes (e.g., >3 for spinal conditions)
- 10 • Multiple new conditions which introduce concerns regarding the cause of these
- 11 conditions

1 Examination

- 2 • Severe signs/findings
- 3 • Results from diagnostic testing that are likely to impact coordination of care and
- 4 response to care (e.g., fracture, joint instability, neurological deficits)
- 5

6 **Assessment of Red Flags**

7 At any time the patient is under care, the practitioner is responsible for seeking and

8 recognizing signs and symptoms that require additional diagnostics, treatment/service,

9 and/or referral. A careful and adequately comprehensive history and evaluation in addition

10 to ongoing monitoring during the course of treatment is necessary to discover potential

11 serious underlying conditions that may need urgent attention. Red flags can present

12 themselves at several points during the patient encounter and can appear in many different

13 forms. If a red flag is identified during a medical necessity review, the clinical quality

14 evaluator should communicate with the provider of services as soon as possible by

15 telephone and/or through standardized communication methods. When a red flag is

16 identified, the clinical quality evaluator may not approve services and recommend

17 returning the patient back to the referring healthcare practitioner or referring the patient to

18 other appropriate health care practitioner/specialist with the measure of urgency as

19 warranted by the history and clinical findings.

20

21 Due to the rarity of actual red flag diagnoses in clinical practice, it is emphasized that the

22 practitioner does not need to perform expensive or invasive diagnostic procedures (e.g., x-

23 ray, advanced imaging, laboratory studies) in the absence of suspicious clinical

24 characteristics. Important red flags and events as well as the points during the clinical

25 encounter at which they are likely to appear include but may not be limited to:

26

27 **Past or Current History**

- 28 • Personal or family history of cancer.
- 29 • Current or recent urinary tract, respiratory tract, or other infection.
- 30 • Anticoagulant therapy or blood clotting disorder.
- 31 • Metabolic bone disorder (osteopenia and osteoporosis).
- 32 • Unintended weight loss.
- 33 • Unexplained dizziness or hearing loss.
- 34 • Trauma with skin penetration; and
- 35 • Immunosuppression (AIDS/ARC).

1 Present Complaint

- 2 • Writhing or cramping pain.
- 3 • Precipitation by significant trauma.
- 4 • Pain that is worse at night or not relieved by any position.
- 5 • Suspicion of cerebrovascular compromise.
- 6 • Symptom's indicative of progressive neurological disorder.

8 Physical Examination/Assessment

- 9 • Inability to reproduce symptoms of musculoskeletal diagnosis or complaints.
- 10 • Pulsing abdominal mass.
- 11 • Fever, chills, or sweats without other obvious source.
- 12 • New or recent neurologic deficit (special senses, sensory, language, and motor).
- 13 • Signs of carotid/vertebrobasilar insufficiency.
- 14 • Uncontrolled hypertension.
- 15 • Signs of nutritional deficiency.
- 16 • Signs of allergic reaction requiring immediate attention.
- 17 • Abuse/neglect.
- 18 • Psychological distress.

20 Pattern of Symptoms Not Consistent with Benign Disorder

- 21 • Chest tightness, difficulty breathing, chest pain.
- 22 • Headache of morbid proportion.
- 23 • Rapidly progressive neurological deficit.
- 24 • Significant, unexplained extremity weakness or clumsiness.
- 25 • Change in bladder or bowel function.
- 26 • New or worsening numbness or paresthesia.
- 27 • Saddle anesthesia.
- 28 • New or recent bilateral radiculopathy.

30 Lack of Response to Appropriate Care

- 31 • History of consultation/care from a series of practitioners or a variety of health care approaches without resolving the patient's complaint.
- 32 • Unsatisfactory clinical progress, especially when compared to apparently similar cases or natural progression of the condition.
- 33 • Signs and symptoms that do not fit the normal pattern and are not resolving.

37 **Assessment of Yellow Flags**

1 When yellow flags are present, clinicians need to be vigilant for deviations from the normal
 2 course of illness and recovery. Examples of yellow flags include depressive symptoms,
 3 injuries still in litigation, signs, and symptoms not consistent with pain severity, and
 4 behaviors incongruent with underlying anatomic and physiologic principles.

5
 6 If a yellow flag is identified during a medical necessity review, the reviewer should
 7 communicate with the provider of services as soon as possible by telephone and/or through
 8 standardized communication methods. The clinical quality evaluator may recommend
 9 returning the patient back to the referring healthcare practitioner or referring the patient to
 10 other health care practitioner/specialist as appropriate.

11 **Assessment of Historical Information**

12 The following factors are assessed in review and determination if the services are medically
 13 necessary:
 14

- 15 • The mechanism of onset and date of onset are congruent with the stated condition's
 16 etiology.
- 17 • The patient's past medical history and response to care do not pose
 18 contraindication(s) for the services submitted for review.
- 19 • The patient's past medical history of pertinent related and unrelated conditions does
 20 not pose contraindication(s) for the services submitted for review.
- 21 • The patient's complaint(s) have component(s) that are likely to respond favorably
 22 to services submitted for review.
- 23 • Provocative and palliative factors identified on examination indicate the presence
 24 of a musculoskeletal condition as expected per diagnosis(es) or complaints, or as
 25 consistent with other type of diagnosis(es).
- 26 • The patient's severity of limitations to activities of daily living (ADLs) are
 27 appropriate and commensurate for the presence of the condition(s) or disorder(s).
- 28 • The quality, radiation, severity, and timing of pain are congruent with the
 29 documented condition(s) or disorder(s).
- 30 • The patient's past medical history of having the same or similar condition(s)
 31 indicates a favorable response to care.
- 32 • The absence or presence of co-morbid condition(s) may or may not present absolute
 33 or relative contraindications to care.

34 **Assessment of Examination Findings**

- 35 • The exam procedures, level of complexity, and components are appropriate for the
 36 patient's complaint(s) and historical findings.
- 37 • Objective palpatory, orthopedic, neurologic, and other physical examination
 38 findings are current, clearly defined, qualified, and quantified, including the nature,
 39

- 1 extent, severity, character, professional interpretation, and significance of the
 2 finding(s) in relation to the patient’s complaint(s) and differential diagnosis(es).
- 3 • Exam findings provide evidence justifying the condition(s) is/are likely to respond
 4 favorably to services submitted for review.
 - 5 • Exam findings provide a reasonable and reliable basis for the stated diagnosis(es).
 - 6 • Exam findings provide a reasonable and reliable basis for treatment planning;
 7 accounting for variables such as age, sex, physical condition, occupational and
 8 recreational activities, co-morbid conditions, etc.
 - 9 • The patient’s progress is being appropriately monitored each visit (as noted within
 10 daily chart notes and during periodic re-exams) to ensure that acceptable clinical
 11 progress is realized.

13 **Assessment of Treatment / Treatment Planning**

- 14 • Treatment dosage (frequency and duration of service) is appropriately correlated
 15 with the nature and severity of the subjective complaints, potential
 16 complications/barriers to recovery, and objective clinical evidence.
- 17 • Services that do not require the professional skills of a practitioner to perform or
 18 supervise are not medically necessary, even if they are performed or supervised by
 19 an Occupational Therapist. Therefore, if the continuation of a patient’s care can
 20 proceed safely and effectively through a home exercise program or self-
 21 management program, services are not indicated or medically necessary.
- 22 • The use of passive modalities in the treatment of subacute or chronic conditions
 23 beyond the acute inflammatory response phase requires documentation of the
 24 anticipated benefit and condition-specific rationale in order to be considered
 25 medically necessary.
- 26 • The treatment plan includes the use of therapeutic procedures to address functional
 27 deficits and ADL restrictions.
- 28 • The set therapeutic goals are functionally oriented, realistic, measurable, and
 29 evidence based.
- 30 • The proposed date of release/discharge from treatment is clearly defined.
- 31 • The treatment/therapies are appropriately correlated with the nature and severity of
 32 the patient’s condition(s) and set treatment goals.
- 33 • Functional Outcome Measures (FOM) demonstrate minimal clinically important
 34 difference (MCID) from baseline results through periodic reevaluations during the
 35 course of care. This is important in order to determine the need for continued care,
 36 the appropriate frequency of visits, estimated date of release from care, and if a
 37 change in the treatment plan or a referral to an appropriate health care
 38 practitioners/specialist is indicated.
- 39 • Home care, self-care, and active-care instructions are documented.

- 1 • Durable Medical Equipment (DME), supplies, appliances, and supports are
- 2 provided when medically necessary and appropriately correlated with clinical
- 3 findings and clinical evidence.

1 **Assessment of Diagnostic Imaging / Special Studies**

- 2 • Laboratory tests are performed only when medically necessary to improve
3 diagnostic accuracy and treatment planning. Abnormal values are professionally
4 interpreted as they relate to the patient's complaint(s) or to unrelated co-morbid
5 conditions that may or may not impact the patient's prognosis and proposed
6 treatment.
- 7 • X-ray procedures are performed only when medically necessary to improve
8 diagnostic accuracy and treatment planning. (Indicators from history and physical
9 examination indicating the need for x-ray procedures are described in the *X-Ray
10 Guidelines (CPG 1-S)* clinical practice guideline).
- 11 • Advanced imaging studies, when medically necessary and/or available, are
12 evaluated for structural integrity and to rule out osseous, related soft tissue
13 pathology, or other pathology.
- 14 • EMG and NCV studies, when medically necessary and/or available, are evaluated
15 for objective evidence of neural deficit. For more information, see the
16 *Electrodiagnostic Testing (CPG 129-S)* clinical practice guideline.
- 17 • Imaging or special studies' findings are appropriate given the nature and severity
18 of the patient's condition(s) and the findings obtained are likely to influence the
19 basis for the proposed treatment.

21 **8.3.2 Factors that Influence Adverse Determinations of Clinical Services (Partial 22 Approvals/Denials)**

23 Factors that influence adverse determinations of clinical services may include but are not
24 limited to these specific considerations and other guidelines and factors identified
25 elsewhere in this policy.: Topics/factors covered elsewhere in this guideline are also
26 applicable in this section and may result in an adverse determination on medical necessity
27 review. To avoid redundancy, many of those factors have not been listed below.

29 **Additional Factors Considered in Determination of Medical Necessity 30 History / Complaints / Patient Reported Outcome Measures**

- 31 • The patient's complaint(s) and/or symptom(s) are not clearly described
- 32 • There is poor correlation and/or a significant discrepancy between the complaint(s)
33 and/or symptom(s) as documented by the treating practitioner and as described by
34 the patient
- 35 • The patient's complaint(s) and/or symptom(s) have not demonstrated clinically
36 significant improvement
- 37 • The nature and severity of the patient's complaint(s) and/or symptom(s) are
38 insufficient to substantiate the medical necessity of any/all submitted services
- 39 • The patient has little or no pain as measured on a valid pain scale

- 1 • The patient has little or no functional deficits using a valid functional outcome
- 2 measure or as otherwise documented by the practitioner

3 **Evaluation Findings**

- 4 • There is poor correlation and/or a significant discrepancy in any of the following:
 - 5 ○ patient's history
 - 6 ○ subjective complaints
 - 7 ○ objective findings
 - 8 ○ diagnosis
 - 9 ○ treatment plan
- 10 • The application of various exam findings to diagnostic or treatment decisions are
- 11 not clearly described or measured (e.g., severity, intensity, professional
- 12 interpretation of results, significance)
- 13 • The patient's objective findings have not demonstrated clinically significant
- 14 improvement
- 15 • The objective findings are essentially normal or are insufficient to support the
- 16 medical necessity of any/all submitted services
- 17 • The submitted objective findings are insufficient due to any of, but not limited to,
- 18 the following reasons:
 - 19 ○ old or outdated relative to the requested dates of service
 - 20 ○ do not properly describe the patient's current status
 - 21 ○ do not substantiate the medical necessity of the current treatment plan do
 - 22 not support the patient's diagnosis/diagnoses do not correlate with the
 - 23 patient's subjective complaint(s) and/or symptom(s)
- 24 • Not all of the patient's presenting complaints were properly examined
- 25 • The patient does not have any demonstrable functional deficits or impairments
- 26 • The patient has not made reasonable progress toward pre-clinical status or
- 27 functional outcomes under the initial treatment/services
- 28 • Clinically significant therapeutic progress is not evident through a review of the
- 29 submitted records. This may indicate that the patient has reached maximum
- 30 therapeutic benefit
- 31 • The patient is approaching or has reached maximum therapeutic benefit
- 32 • The patient's exam findings have returned to pre-injury status or prior level of
- 33 function
- 34 • There is inaccurate reporting of clinical findings
- 35 • The exam performed is for any of the following:
 - 36 ○ Wellness
 - 37 ○ pre-employment
 - 38 ○ sports pre-participation
- 39 • The exam performed is non-standard and solely technique/protocol based

1 **Diagnosis**

- 2 • The diagnosis is not supported by one or more of the following:
 - 3 ○ patient's history (e.g., date/mechanism of onset)
 - 4 ○ subjective complaints (e.g., nature and severity, location)
 - 5 ○ objective findings (e.g., not clearly defined and/or quantified, not
 - 6 professionally interpreted, significance not noted)

8 **Submitted Medical Records**

- 9 • The submitted records are insufficient to reliably verify pertinent clinical
 - 10 information, such as (but not limited to):
 - 11 ○ patient's clinical health status
 - 12 ○ the nature and severity of the patient's complaint(s) and/or symptom(s)
 - 13 ○ date/mechanism of onset
 - 14 ○ objective findings
 - 15 ○ diagnosis/diagnoses
 - 16 ○ response to care
 - 17 ○ functional deficits/limitations
 - 18 • There are daily notes submitted for the same dates of service with different/altere
 - 19 findings without an explanation
 - 20 • There is evidence of duplicated or nearly duplicated records for the same patient
 - 21 for different dates of service, or for different patients
 - 22 • There is poor correlation and/or a significant discrepancy between the information
 - 23 presented in the submitted records with the information presented during a verbal
 - 24 communication between the reviewing clinical quality evaluator and treating
 - 25 practitioner
 - 26 • The treatment time (in minutes) and/or the number of units used in the performance
 - 27 of a timed service (e.g., modality, procedure) during each encounter/office visit was
 - 28 not documented
 - 29 • Some or all of the service(s) submitted for review are not documented as having
 - 30 been performed in the daily treatment notes

32 **Treatment / Treatment Planning**

- 33 • The submitted records show that the nature and severity of the patient's
 - 34 complaint(s) and/or symptom(s) require a limited, short trial of care in order to
 - 35 monitor the patient's response to care and determine the efficacy of the current
 - 36 treatment plan. This may include, but not limited to, any of the following:
 - 37 ○ significant trauma affecting function
 - 38 ○ acute/sub-acute stage of condition
 - 39 ○ moderate-to-severe or severe subjective and objective findings

- 1 ○ possible neurological involvement
- 2 ○ presence of co-morbidities that may significantly affect the treatment plan
- 3 and/or the patient's response to care
- 4 • There is poor correlation of the treatment plan with the nature and severity of the
- 5 patient's complaint(s) and/or symptom(s), such as (but not limited to):
- 6 ○ use of acute care protocols for chronic condition(s)
- 7 ○ prolonged reliance on passive care
- 8 ○ active care and reduction of passive care are not included in the treatment
- 9 plan
- 10 ○ inappropriate use of passive modalities in the plan of care
- 11 ○ use of passive modalities as stand-alone treatments (which is rarely
- 12 therapeutic) or as the sole treatment approach to the patient's condition(s)
- 13 • There is evidence from the submitted records that the patient's treatment can
- 14 proceed safely and effectively through a home exercise program or self-
- 15 management program
- 16 • The patient's function has improved, complaints and symptoms have decreased,
- 17 and patient requires less treatment (e.g., lesser units of services per office visit,
- 18 lesser frequency, shorter total duration to discharge)
- 19 • The patient's symptoms and/or exam findings are mild and the patient's treatment
- 20 plan requires a lesser frequency (e.g., units of services, office visits per week)
- 21 and/or total duration
- 22 • Therapeutic goals have not been documented; goals should be measurable and
- 23 written in terms of function and include specific parameters
- 24 • Therapeutic goals have not been reassessed in a timely manner to determine if the
- 25 patient is making expected progress
- 26 • Failure to make progress or respond to care as documented within subjective
- 27 complaints, objective findings and/or functional outcome measures
- 28 • The patient's condition(s) is/are not amenable to the proposed treatment plan
- 29 • Additional significant improvement cannot be reasonably expected by continued
- 30 treatment and treatment must be changed or discontinued
- 31 • The patient has had ongoing care without any documented lasting therapeutic
- 32 benefits
- 33 • The condition requires an appropriate referral and/or coordination with other
- 34 appropriate health care services
- 35 • The patient is not complying with the treatment plan that includes lifestyle changes
- 36 to help reduce frequency and intensity of symptoms
- 37 • The patient is not adhering to treatment plan that includes medically necessary
- 38 frequency and intensity of services

- 1 • The use of multiple passive modalities with the same or similar physiologic effects
- 2 to the identical region is considered redundant and not reasonable or medically
- 3 necessary
- 4 • Home care, self-care, and/or active-care instructions are not implemented or
- 5 documented in the submitted records
- 6 • Uncomplicated diagnoses do not require services beyond the initial treatment plan
- 7 before discharging the patient to active home/self-care
- 8 • As symptoms and clinical findings improve the frequency of services (e.g., visits
- 9 per week/month) did not decrease. The submitted services do not or no longer
- 10 require the professional skills of the treating practitioner.
- 11 • The treatment plan is for any of the following:
- 12 o preventive care
- 13 o elective/convenience/wellness care
- 14 o back school
- 15 o vocational rehabilitation or return to work programs
- 16 o work hardening programs
- 17 o routine educational, training, conditioning, return to sport, or fitness.
- 18 o non-covered condition
- 19 • There is duplication of services with other healthcare practitioners/specialties.
- 20 • The treatment plan is not supported due to, but not limited to, any of the following
- 21 reasons:
- 22 o technique-/protocol-based instead of individualized and evidence based
- 23 o generic and not individualized for the patient's specific needs
- 24 o does not correlate with the set therapeutic goals
- 25 o not supported in the clinical literature (e.g., proprietary, unproven)
- 26 o not considered evidence-based and/or professionally accepted
- 27 • The treatment plan includes services that are considered not evidence-based, not
- 28 widely accepted, unproven and/or not reasonable or medically necessary,
- 29 inappropriate or unrelated to the patient's complaint(s) and/or diagnosis/diagnoses
- 30 (e.g., Low level laser therapy, axial/spinal decompression, select forms of EMS
- 31 such as microcurrent, H-wave. Also see the *Techniques and Procedures Not Widely*
- 32 *Supported as Evidence-Based (CPG 133 – S)* clinical practice guideline for
- 33 complete list)

34 **Health and Safety**

- 35 • There are signs, symptoms and/or other pertinent information presented through the
- 36 patient's history, exam findings, and/or response to care that require urgent
- 37 attention, further testing, and/or referral to and/or coordination with other
- 38 healthcare practitioners/specialists
- 39

- 1 • There is evidence of the presence of Yellow and/or Red Flags (See section on Red
- 2 and Yellow Flags above)
- 3 • There are historical, subjective, and/or objective findings which present as
- 4 contraindications for the plan of care

6 **8.3.3 Referral / Coordination of Services**

7 When a potential health and safety issue is identified, the clinical quality evaluator must
 8 communicate with the provider of services as soon as possible by telephone and/or through
 9 standardized communication methods to recommend returning the patient back to the
 10 referring health care practitioner or referring the patient to other appropriate health care
 11 practitioner/specialist with the measure of urgency as warranted by the history and clinical
 12 findings.

13
 14 Clinical factors that may require referral or coordination of services include, but not limited
 15 to:

- 16 • Symptoms worsening following treatment;
- 17 • Deteriorating condition (e.g., orthopedic or neurologic findings, function, etc.);
- 18 • Reoccurring exacerbations despite continued treatment;
- 19 • No progress despite treatment;
- 20 • Unexplained diagnostic findings (e.g., suspicion of fracture);
- 21 • Identification of Red Flags;
- 22 • Identification of co-morbid conditions that don't appear to have been addressed
- 23 previously that represent absolute contraindications to services;
- 24 • Constitutional signs and symptoms indicative of systemic condition (e.g.,
- 25 unintended weight loss of greater than 4.5 kg/10 lbs. over 6-month period);
- 26 • Inability to provoke symptoms with standard exam;
- 27 • Treatment needed outside of scope of practice.

28
 29 The Clinical Policy is reviewed and approved by the ASH Clinical Quality committees that
 30 are comprised of contracted network practitioners including practitioners of the same
 31 clinical discipline as the treating providers for whom compliance with the practices
 32 articulated in this this document is required. Guidelines are updated at least annually, or as
 33 new information is identified that result in material changes to one or more of these
 34 policies.

35 36 **9. LITERATURE REVIEW**

37 **9.1 Occupational Therapy for Conditions Considered Unproven**

38 There is a limited amount of evidence regarding individual occupational therapy
 39 interventions for specific conditions. There are several Cochrane systematic reviews and

1 other reviews that have been published regarding occupational therapy for various
 2 conditions (Steultjens, et al., 2004; Steultjens et al., 2005; Legg et al., 2006; Dixon et al.,
 3 2007; Hoffman et al., 2011; Hoare et al., 2021; Quinn et al., 2021; Legg et al., 2021; García-
 4 Pérez et al., 2021; Fields and Smallfield, 2022; Cunningham et al., 2022; Wood et al.,
 5 2022). The reviews in general found that that there is improvement seen with occupational
 6 therapy however, evidence with respect to specific interventions is limited. Passive
 7 modalities, such as ultrasound, electric stimulation, traction, laser, and hot and cold packs,
 8 are often used in combination with manual therapies and exercise despite insufficient
 9 and/or inconclusive evidence for many conditions. Often methodologic flaws and
 10 heterogeneity of studies result in an inability to draw confirmatory conclusions.

11 **9.2 Specific Occupational Therapy Treatments Considered Unproven**

12 **Constraint-Induced Movement Therapy (CIMT)**

13 Constraint-induced movement therapy (CIMT) is a multi-faceted intervention that has been
 14 proposed for neurological conditions that involve hemiparesis. CIMT is also referred to as
 15 constraint-induced therapy or forced use therapy and is primarily provided by physical
 16 therapists and occupational therapists. Several variations exist based on method and length
 17 of restraint, and type and duration of therapy (e.g., environment and provider). The therapy
 18 involves constraining the unaffected arm or hand with a sling, glove or mitt. CIMT
 19 typically involves intensive individualized therapy with up to six–eight hours of therapy
 20 provided per day. However, other forms of modified CIMT have been developed with less
 21 therapy provided, but longer periods of restraint (Wolf, 2007). Veterans Affairs/Dept of
 22 Defense (VA/DoD) published guidelines that have also been endorsed by American Heart
 23 Association/American Stroke Association (AHA/ASA)—Clinical Practice Guideline for
 24 the Management of Adult Stroke Rehabilitation Care (Bates, et al., 2005). The guidelines
 25 note that, “Use of constraint-induced therapy should be considered for a select group of
 26 patients—that is, patients with 20 degrees of wrist extension and 10 degrees of finger
 27 extension, who have no sensory and cognitive deficits.” Indicating a recommendation that
 28 the intervention may be considered). The Royal College of Physicians/Intercollegiate
 29 Stroke Working Party (United Kingdom) and the Ottawa Panel (2006) agree with these
 30 recommendations.
 31

32
 33 CIMT has demonstrated inconsistent effectiveness for treatment of patients post-stroke
 34 (Abdullahi et al., 2020; Pulman et al., 2013; McIntyre et al., 2012; Corbetta et al., 2010;
 35 Sirtori et al., 2009; Abdullahi et al., 2021a; Abdullahi et al., 2021b; Alaca and Ocal, 2022).
 36 Future randomized controlled trials need to have accurate characteristics in terms of
 37 methodological quality, larger samples, longer follow up, reliable and relevant measure
 38 and report of adverse events. Some evidence demonstrates that modified CIMT could
 39 reduce the level of disability, improve the ability to use the paretic upper extremity, and
 40 enhance spontaneity during movement time, but evidence is still limited about the

1 effectiveness of modified CIMT in kinematic analysis (Pollack et al., 2014; Shi et al.,
2 2011). Research suggests that modified CIMT and intensive CIMT produce similar results
3 (Peurala et al., 2012).

4
5 CIMT has also been used for the treatment of children with cerebral palsy (CP). Research
6 is not conclusive with regards to the effectiveness of CIMT for this population; however
7 there appears to be modest evidence to support its use in a modified format (Novak et al.,
8 2020; Taub et al., 2004; Sakzewski et al., 2009; Eliasson et al., 2005; Hoare et al., 2007;
9 Chen et al., 2014; Chiu and Ada, 2016; Eliasson et al., 2014, Hoare et al., 2019; Martínez-
10 Costa Montero et al., 2020; Ramey et al., 2021; Walker et al., 2022; Dionisio and Terrill,
11 2022; Jackman et al., 2022; Baker et al., 2022). Further research using adequately powered
12 RCTs [randomized controlled trials], rigorous methodology and valid, reliable outcome
13 measures is essential to provide higher level support of the effectiveness of CIMT for
14 children with hemiplegic cerebral palsy.

15 16 **Intensive Model of Therapy (IMOT) Programs**

17 Refer to *Intensive Model of Therapy (CPG 286 – S)* clinical practice guideline for more
18 information.

19 20 **Dry Hydrotherapy**

21 Dry hydrotherapy, also referred to as aqua massage, water massage, or hydromassage, is a
22 treatment that incorporates water with the intent of providing therapeutic massage. The
23 treatment is generally provided in chiropractor or therapy offices. There are several dry
24 hydrotherapy devices available that provide this treatment, including the following:

- 25 • Aqua Massage® (AMI Inc., Mystic, CT)
- 26 • AquaMED® (JTL Enterprises, Inc., Clearwater, FL)
- 27 • H2Omassage System™ (H2Omassage Systems, Winnipeg, MB, Canada)
- 28 • Hydrotherapy Tables (Sidmar Manufacturing, Inc., Princeton, MN)

29
30 Proponents of dry hydrotherapy maintain that it can be used in lieu of certain conventional
31 physical medicine therapeutic modalities and procedures, such as heat packs, wet
32 hydrotherapy, massage, and soft tissue manipulation. The assertions that have been made
33 by manufacturers of this device at their websites have not yet been proven. No published
34 studies or information regarding dry hydrotherapy devices or dry hydrotherapy treatment
35 were identified in the peer-reviewed scientific literature. In the absence of peer- reviewed
36 literature demonstrating the effectiveness of dry hydrotherapy and in the absence of
37 comparison to currently accepted treatment modalities, no definitive conclusions can be
38 drawn regarding the clinical benefits of this treatment.

39

1 **Non-invasive Interactive Neurostimulation (e.g., InterX®)**

2 Refer to *Non-invasive Interactive Neurostimulation (InterX®)* (CPG 277 – S) clinical
3 practice guideline for more information.

4
5 **Microcurrent Electrical Nerve Stimulation (MENS)**

6 Refer to *Electric Stimulation for Pain, Swelling and Function in the Clinic Setting* (CPG
7 272 – S) clinical practice guideline for more information.

8
9 **H-WAVE®**

10 Refer to *H-WAVE® Electrical Stimulation* (CPG 269 – S) clinical practice guideline for
11 more information.

12 **Equestrian Therapy (e.g., Hippotherapy)**

13 Equestrian therapy, also known as hippotherapy, is proposed to offer a person with a
14 disability a means of physical activity that aids in improving balance, posture, coordination,
15 the development of a positive attitude and a sense of accomplishment. It is proposed for
16 treatment of several conditions including autism spectrum disorders and cerebral palsy.
17 There is insufficient published evidence regarding the effects of this therapy on individuals
18 with impaired physical function resulting from illness, injury, congenital defect or surgery
19 (Bronson et al., 2010; Lee et al., 2014; O’Haire et al., 2014; De Guindos-Sanchez et al.,
20 2020; Marquez et al., 2020; White et al., 2020; Santos de Assis et al., 2022; Pantera et al.,
21 2022; Pérez-Gómez et al., 2022; Heussen and Häusler, 2022; Prieto et al., 2022). It is noted
22 that most studies are limited by methodological weaknesses.

23
24 **MEDEK Therapy**

25 Refer to *MEDEK Therapy* (CPG 276 – S) clinical practice guideline for more information.

26
27 **The Interactive Metronome Program**

28 Interactive Metronome® (IM) is purported to be an assessment and training tool that
29 measures and improves Neurotiming, or the synchronization of neural impulses within key
30 brain networks for cognitive, communicative, sensory and motor performance. It is
31 designed to improve processing speed, focus, and coordination. Patients wear headphones
32 and match a beat using a hand or foot sensor along with visual and auditory feedback. The
33 IM program has been promoted as a treatment for children with attention-deficit
34 hyperactivity disorder (ADHD) and for other special needs children to increase
35 concentration, focus, and coordination. It has also been promoted to improve athletic
36 performance, to assess and improve academic performance of normal children, and to
37 improve children’s performance in the arts (e.g., dance, music, theater, creative arts).
38 Additionally, it has been implemented as part of a therapy program for patients with
39 balance disorders, cerebrovascular accident, limb amputation, multiple sclerosis,

1 Parkinson's disease, and traumatic brain injury. However, based on peer-reviewed
2 literature, evidence is insufficient to support effectiveness of the IM program. Well-
3 designed clinical studies are needed to determine the effectiveness of the IM program and
4 whether a clinically significant improvement is achieved.

5

6 **Taping/Elastic Therapeutic Tape (e.g., Kinesio™ Tape, Spidertech™ Tape)**

7 Refer to *Strapping and Taping (CPG 143 – S)* clinical practice guideline for more
8 information.

9

10 **Dry Needling**

11 Refer to *Dry Needling (CPG 178 – S)* clinical practice guideline for more information.

Laser Therapy (LT)

Refer to *Laser Therapy (LT) (CPG 30 – S)* clinical practice guideline for more information.

10. CODING/BILLING INFORMATION

Note: 1) This list of codes may not be all-inclusive.
 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Covered When Medically Necessary

CPT® Code	CPT® Code Description
97010	Application of a modality to 1 or more areas; hot or cold packs
97012	Application of a modality to 1 or more areas; traction, mechanical
97014	Application of a modality to 1 or more areas; electrical stimulation (unattended)
97016	Application of a modality to 1 or more areas; vasopneumatic devices
97018	Application of a modality to 1 or more areas; paraffin bath
97022	Application of a modality to 1 or more areas; whirlpool
97024	Application of a modality to 1 or more areas; diathermy (e.g., microwave)
97026	Application of a modality to 1 or more areas; infrared
97028	Application of a modality to 1 or more areas; ultraviolet
97032	Application of a modality to 1 or more areas; electrical stimulation (manual), each 15 minutes
97033	Application of a modality to 1 or more areas; iontophoresis, each 15 minutes
97034	Application of a modality to 1 or more areas; contrast baths, each 15 minutes
97035	Application of a modality to 1 or more areas; ultrasound, each 15 minutes
97036	Application of a modality to 1 or more areas; Hubbard tank, each 15 minutes
97110	Therapeutic procedure, 1 or more areas, each 15 minutes; therapeutic exercises to develop strength and endurance, range of motion and flexibility
97112	Therapeutic procedure, 1 or more areas, each 15 minutes; neuromuscular reeducation of movement, balance, coordination,

CPT® Code	CPT® Code Description
	kinesthetic sense, posture, and/or proprioception for sitting and/or standing activities
97113	Therapeutic procedure, 1 or more areas, each 15 minutes; aquatic therapy with therapeutic exercises
97116	Therapeutic procedure, 1 or more areas, each 15 minutes; gait training (includes stair climbing)
97124	Therapeutic procedure, 1 or more areas, each 15 minutes; massage, including effleurage, petrissage and/or tapotement (stroking, compression, percussion)
97140	Manual therapy techniques (e.g., mobilization/manipulation, manual lymphatic drainage, manual traction), 1 or more regions, each 15 minutes
97165	Occupational therapy evaluation, low complexity, requiring these components: An occupational profile and medical and therapy history, which includes a brief history including review of medical and/or therapy records relating to the presenting problem; An assessment(s) that identifies 1-3 performance deficits (i.e., relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of low complexity, which includes an analysis of the occupational profile, analysis of data from problem-focused assessment(s), and consideration of a limited number of treatment options. Patient presents with no comorbidities that affect occupational performance. Modification of tasks or assistance (e.g., physical or verbal) with assessment(s) is not necessary to enable completion of evaluation component. Typically, 30 minutes are spent face-to-face with the patient and/or family.
97166	Occupational therapy evaluation, moderate complexity, requiring these components: An occupational profile and medical and therapy history, which includes an expanded review of medical and/or therapy records and additional review of physical, cognitive, or psychosocial history related to current functional performance; An assessment(s) that identifies 3-5 performance deficits (i.e., relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of moderate analytic complexity, which includes an analysis of the occupational profile, analysis of data from detailed assessment(s), and consideration of several treatment options. Patient may present with comorbidities that affect occupational performance. Minimal to moderate

CPG 155 Revision 18 – S

Occupational Therapy Medical Policy/Guideline

Revised – December 21, 2023

To CQT for review 11/13/2023

CQT reviewed 11/13/2023

To QIC for review and approval 12/05/2023

QIC reviewed and approved 12/05/2023

To QOC for review and approval 12/21/2023

QOC reviewed and approved 12/21/2023

To MA-UMC for review and approval 06/28/2024

MA-UMC reviewed and approved 06/28/24

CPT® Code	CPT® Code Description
	modification of tasks or assistance (e.g., physical or verbal) with assessment(s) is necessary to enable patient to complete evaluation component. Typically, 45 minutes are spent face-to-face with the patient and/or family.
97167	Occupational therapy evaluation, high complexity, requiring these components: An occupational profile and medical and therapy history, which includes review of medical and/or therapy records and extensive additional review of physical, cognitive, or psychosocial history related to current functional performance; An assessment(s) that identify 5 or more performance deficits (i.e., relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and a clinical decision-making is of high analytic complexity, which includes an analysis of the patient profile, analysis of data from comprehensive assessment(s), and consideration of multiple treatment options. Patient presents with comorbidities that affect occupational performance. Significant modification of tasks or assistance (e.g., physical or verbal) with assessment(s) is necessary to enable patient to complete evaluation component. Typically, 60 minutes are spent face-to-face with the patient and/or family.
97168	Re-evaluation of occupational therapy established plan of care, requiring these components: An assessment of changes in patient functional or medical status with revised plan of care; An update to the initial occupational profile to reflect changes in condition or environment that affect future interventions and/or goals; and a revised plan of care. A formal reevaluation is performed when there is a documented change in functional status or a significant change to the plan of care is required. Typically, 30 minutes are spent face-to-face with the patient and/or family
97530	Therapeutic activities, direct (one-on-one) patient contact (use of dynamic activities to improve functional performance), each 15 minutes
97535	Self-care/home management training (e.g., activities of daily living (ADL) and compensatory training, meal preparation, safety procedures, and instructions in use of assistive technology devices/adaptive equipment) direct one-on-one contact, each 15 minutes
97542	Wheelchair management (e.g., assessment, fitting, training), each 15 minutes

CPG 155 Revision 18 – S

Occupational Therapy Medical Policy/Guideline

Revised – December 21, 2023

To CQT for review 11/13/2023

CQT reviewed 11/13/2023

To QIC for review and approval 12/05/2023

QIC reviewed and approved 12/05/2023

To QOC for review and approval 12/21/2023

QOC reviewed and approved 12/21/2023

To MA-UMC for review and approval 06/28/2024

MA-UMC reviewed and approved 06/28/24

CPT® Code	CPT® Code Description
97760	Orthotic(s) management and training (including assessment and fitting when not otherwise reported), upper extremity(s), lower extremity(s) and/or trunk, initial orthotic(s) encounter, each 15 minutes
97761	Prosthetic training, upper and/or lower extremity(s), initial prosthetic(s) encounter, each 15 minutes
97763	Orthotic(s)/prosthetic(s) management and/or training, upper extremity(ies), lower extremity(ies), and/or trunk, subsequent orthotic(s)/prosthetic(s) encounter, each 15 minutes

HCPCS Code	HCPCS Code Description
G0129	Occupational therapy services requiring the skills of a qualified occupational therapist, furnished as a component of a partial hospitalization treatment program, per session (45 minutes or more)
G0152	Services performed by a qualified occupational therapist in the home health or hospice setting, each 15 minutes
G0158	Services performed by a qualified occupational therapist assistant in the home health or hospice setting, each 15 minutes
G0160	Services performed by a qualified occupational therapist, in the home health setting, in the establishment or delivery of a safe and effective therapy maintenance program, each 15 minutes
G0237	Therapeutic procedures to increase strength or endurance of respiratory muscles, face-to-face, one-on-one, each 15 minutes (includes monitoring)
G0238	Therapeutic procedures to improve respiratory function, other than described by G0237, one-on-one, face-to-face, per 15 minutes (includes monitoring)
G0239	Therapeutic procedures to improve respiratory function or increase strength or endurance of respiratory muscles, two or more individuals (includes monitoring)
S9129	Occupational therapy, in the home, per diem

1

2 **Training in Nature/Not Medically Necessary/Not Covered**

CPT® Code	CPT® Code Description
97169	Athletic training evaluation, low complexity, requiring these components: A history and physical activity profile with no comorbidities that affect physical activity; An examination of affected body area and other symptomatic or related systems addressing 1-2 elements from any of the following: body structures, physical activity, and/or participation deficiencies; and Clinical decision making of low complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 15 minutes are spent face-to-face with the patient and/or family
97170	Athletic training evaluation, moderate complexity, requiring these components: A medical history and physical activity profile with 1-2 comorbidities that affect physical activity. An examination of affected body area and other symptomatic or related systems addressing a total of 3 or more elements from any of the following: body structures, physical activity, and/or participation deficiencies; and Clinical decision making of moderate complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 30 minutes are spent face-to-face with the patient and/or family.
97171	Athletic training evaluation, high complexity, requiring these components: A medical history and physical activity profile, with 3 or more comorbidities that affect physical activity; A comprehensive examination of body systems using standardized tests and measures addressing a total of 4 or more elements from any of the following: body structures, physical activity, and/or participation deficiencies; Clinical presentation with unstable and unpredictable characteristics; and Clinical decision making of high complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 45 minutes are spent face-to-face with the patient and/or family.
97172	Re-evaluation of athletic training established plan of care requiring these components: An assessment of patient’s current functional status when there is a documented change, and A revised plan of care using a standardized patient assessment instrument and/or measurable assessment of functional outcome with an update in management options, goals, and interventions. Typically, 20 minutes are spent face-to-face with the patient and/or family.

CPT® Code	CPT® Code Description
97537	Community/work reintegration training (e.g., shopping, transportation, money management, avocational activities and/or work environment/modification analysis, work task analysis, use of assistive technology device/adaptive equipment), direct one-on-one contact, each 15 minutes
97545	Work hardening/conditioning; initial 2 hours
97546	Work hardening/conditioning; each additional hour (List separately in addition to code for primary procedure)

1

HCPCS Code	HCPCS Code Description
S8990	Physical or manipulative therapy performed for maintenance rather than restoration
S9117	Back school, per visit

1
2 Unproven and not covered when used to report constraint-induced movement therapy or
3 dry hydrotherapy/aqua massage/hydromassage, equestrian therapy (e.g., hippotherapy),
4 elastic therapeutic tape/taping, low-level laser:
5

HCPCS Code	HCPCS Code Description
S8940	Equestrian/hippotherapy, per session
S8948	Application of a modality (requiring constant provider attendance) to one or more areas, low-level laser; each 15 minutes

6
7 **References**

8 Abdullahi A, Van Criekeing T, Umar NA, Zakari UU, Truijen S, Saeys W. Effect of
9 constraint-induced movement therapy on persons-reported outcomes of health status
10 after stroke: a systematic review and meta-analysis. *Int J Rehabil Res.* 2020 Nov 23.

11
12 Abdullahi A, Aliyu NU, Useh U, et al. Comparing two different modes of task practice
13 during lower limb constraint-induced movement therapy in people with stroke: A
14 randomized clinical trial. *Neural Plast.* 2021a;2021:6664058.

15
16 Abdullahi A, Truijen S, Umar NA, et al. Effects of lower limb constraint induced
17 movement therapy in people with stroke: A systematic review and meta-analysis.
18 *Front Neurol.* 2021b;12:638904
19

20 Agency for Healthcare Research and Quality. Multidisciplinary Postacute Rehabilitation
21 for Moderate to Severe Traumatic Brain Injury in Adults. Effective Health Care
22 Program. Comparative Effectiveness Review, 2012;72. Retrieved on April 18, 2023
23 from
24 https://www.ncbi.nlm.nih.gov/books/NBK98993/pdf/Bookshelf_NBK98993.pdf
25

26 Alaca N, Öcal NM. Proprioceptive based training or modified constraint-induced
27 movement therapy on upper extremity motor functions in chronic stroke patients: A
28 randomized controlled study. *NeuroRehabilitation.* 2022;51(2):271-282.
29 doi:10.3233/NRE-220009
30

- 1 American Medical Association. (current year). Current Procedural Terminology (CPT)
2 Current year (rev. ed.). Chicago: AMA.
3
- 4 American Occupational Therapy Association. Retrieved on April 18, 2023 from
5 <http://www.aota.org>
6
- 7 American Occupational Therapy Association. Official Documents. Retrieved on April 18,
8 2023 from <http://www.aota.org/en/Practice/Manage/Official.aspx>
9
- 10 American Occupational Therapy Association, Inc. (AOTA). Consumer Information.
11 Retrieved on April 18, 2023 from [http://www.aota.org/en/About-Occupational-](http://www.aota.org/en/About-Occupational-Therapy/Patients-Clients.aspx)
12 [Therapy/Patients-Clients.aspx](http://www.aota.org/en/About-Occupational-Therapy/Patients-Clients.aspx)
13
- 14 American Occupational Therapy Association, Inc. (AOTA). OT Services in Work
15 Rehabilitation. Retrieved on April 18, 2023 from [http://www.aota.org/About-](http://www.aota.org/About-Occupational-Therapy/Professionals/WI.aspx)
16 [Occupational-Therapy/Professionals/WI.aspx](http://www.aota.org/About-Occupational-Therapy/Professionals/WI.aspx)
17
- 18 American Occupational Therapy Association. Occupational Therapy Scope of Practice.
19 *Am J Occup Ther.* 2021;75(Supplement_3):7513410020.
20 doi:10.5014/ajot.2021.75S3005
21
- 22 American Occupational Therapy Association; Standards of Practice for Occupational
23 Therapy. *Am J Occup Ther* November/December 2021, Vol. 75(Supplement_3),
24 7513410030. doi: <https://doi.org/10.5014/ajot.2021.75S3004>
25
- 26 Aqua Massage [product description]. AMI Inc. Retrieved on April 18, 2023 from
27 http://amiaqua.com/PR_overview.htm
28
- 29 AquaMED Dry Hydrotherapy. JTL Enterprises, Inc. Retrieved on April 18, 2023 from
30 <http://www.hydrmassage.com/>
31
- 32 Baker A, Niles N, Kysh L, Sargent B. Effect of Motor Intervention for Infants and Toddlers
33 With Cerebral Palsy: A Systematic Review and Meta-analysis. *Pediatr Phys Ther.*
34 2022;34(3):297-307. doi:10.1097/PEP.0000000000000914
35
- 36 Boyd R, Sakzewski L, Ziviani J, Abbott DF, Badawy R, Gilmore R, et al. INCITE: A
37 randomized trial comparing constraint induced movement therapy and bimanual
38 training in children with congenital hemiplegia. *BMC Neurol.* 2010 Jan 12;10:4.
39

- 1 Boyles R, Fowler R, Ramsey D, Burrows E. Effectiveness of trigger point dry needling for
- 2 multiple body regions: a systematic review. J Man Manip Ther. 2015 Dec;23(5):276-
- 3 93.

- 1 Brogårdh C, Flansbjer UB, Lexell J. What is the long-term benefit of constraint-induced
2 movement therapy? A four-year follow-up. Clin Rehabil. 2009 May;23(5):418-23.
3 Epub 2009 Apr 6.
4
- 5 Bronson C, Brewerton K, Ong J, Palanca C, Sullivan SJ. Does hippotherapy improve
6 balance in persons with multiple sclerosis: a systematic review. Eur J Phys Rehabil
7 Med. 2010 Sep;46(3):347-53.
8
- 9 Cameron M. Physical Agents in Rehabilitation: An Evidence-Based Approach to Practice.
10 6th Edition. Elsevier; 2022.
11
- 12 Centers for Medicare and Medicaid Services (CMS). Pub. 100-02, Chapter 15, Sections
13 220 and 230 Therapy Services. Coverage of Outpatient Rehabilitation Therapy
14 Services (Physical Therapy, Occupational Therapy, and Speech-Language Pathology
15 Services) Under Medical Insurance (Rev. 11905, 03-16-23). Retrieved on April 18,
16 2023 from <http://www.cms.hhs.gov/manuals/Downloads/bp102c15.pdf>
17
- 18 Centers for Medicare & Medicaid Services (CMS). Local Coverage Article: Medical
19 Necessity of Therapy Services (A52775). Retrieved on October 18, 2022 from
20 [https://www.cms.gov/medicare-coverage-](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=52775&ver=15&keyword=medical+necessity+of+therapy+services&keywordType=starts&areaId=all&docType=NCA%2cCAL%2cNCD%2cMEDCAC%2cTA%2cMCD%2c6%2c3%2c5%2c1%2cF%2cP&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
21 [database/view/article.aspx?articleid=52775&ver=15&keyword=medical+necessity+o](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=52775&ver=15&keyword=medical+necessity+of+therapy+services&keywordType=starts&areaId=all&docType=NCA%2cCAL%2cNCD%2cMEDCAC%2cTA%2cMCD%2c6%2c3%2c5%2c1%2cF%2cP&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
22 [f+therapy+services&keywordType=starts&areaId=all&docType=NCA%2cCAL%2c](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=52775&ver=15&keyword=medical+necessity+of+therapy+services&keywordType=starts&areaId=all&docType=NCA%2cCAL%2cNCD%2cMEDCAC%2cTA%2cMCD%2c6%2c3%2c5%2c1%2cF%2cP&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
23 [NCD%2cMEDCAC%2cTA%2cMCD%2c6%2c3%2c5%2c1%2cF%2cP&contractOp](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=52775&ver=15&keyword=medical+necessity+of+therapy+services&keywordType=starts&areaId=all&docType=NCA%2cCAL%2cNCD%2cMEDCAC%2cTA%2cMCD%2c6%2c3%2c5%2c1%2cF%2cP&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
24 [tion=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&Ke](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=52775&ver=15&keyword=medical+necessity+of+therapy+services&keywordType=starts&areaId=all&docType=NCA%2cCAL%2cNCD%2cMEDCAC%2cTA%2cMCD%2c6%2c3%2c5%2c1%2cF%2cP&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
25 [yWordSearchType=Exact](https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=52775&ver=15&keyword=medical+necessity+of+therapy+services&keywordType=starts&areaId=all&docType=NCA%2cCAL%2cNCD%2cMEDCAC%2cTA%2cMCD%2c6%2c3%2c5%2c1%2cF%2cP&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
26
- 27 Centers for Medicare and Medicaid. Local Coverage Determination (LCD): Outpatient
28 Physical and Occupational Therapy Services (L33631). Retrieved on October 18, 2022
29 from [https://www.cms.gov/medicare-coverage-database/details/lcd-](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?lcdid=33631&ver=51&keyword=outpatient%20physical%20and%20occupational%20therapy&keywordType=starts&areaId=all&docType=NCA,CAL,NCD,MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
30 [details.aspx?lcdid=33631&ver=51&keyword=outpatient%20physical%20and%20occ](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?lcdid=33631&ver=51&keyword=outpatient%20physical%20and%20occupational%20therapy&keywordType=starts&areaId=all&docType=NCA,CAL,NCD,MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
31 [upational%20therapy&keywordType=starts&areaId=all&docType=NCA,CAL,NCD,](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?lcdid=33631&ver=51&keyword=outpatient%20physical%20and%20occupational%20therapy&keywordType=starts&areaId=all&docType=NCA,CAL,NCD,MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
32 [MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=AAAA](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?lcdid=33631&ver=51&keyword=outpatient%20physical%20and%20occupational%20therapy&keywordType=starts&areaId=all&docType=NCA,CAL,NCD,MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
33 [AAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?lcdid=33631&ver=51&keyword=outpatient%20physical%20and%20occupational%20therapy&keywordType=starts&areaId=all&docType=NCA,CAL,NCD,MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact)
34
- 35 Chen YP, Pope S, Tyler D, Warren GL. Effectiveness of constraint-induced movement
36 therapy on upper-extremity function in children with cerebral palsy: A systematic
37 review and meta-analysis of randomized controlled trials. Clin Rehabil.
38 2014;28(10):939-953.

- 1 Chiu HC, Ada L. Constraint-induced movement therapy improves upper limb activity and
2 participation in hemiplegic cerebral palsy: A systematic review. *J Physiother.*
3 2016;62(3):130-137.
4
- 5 Christiansen AS, Lange C. Intermittent versus continuous physiotherapy in children with
6 cerebral palsy. *Dev Med Child Neurol.* 2008 Apr;50(4):290-3.
7
- 8 Comparative Effectiveness Review No. 169. (Prepared by the Pacific Northwest Evidence-
9 based Practice Center under Contract No. 290-2012-00014-I.) AHRQ Publication No.
10 16-EHC004-EF. Rockville, MD: Agency for Healthcare Research and Quality;
11 February 2016.
12
- 13 Corbetta D, Sirtori V, Moja L, Gatti R. Constraint-induced movement therapy in stroke
14 patients: systematic review and meta-analysis. *Eur J Phys Rehabil Med.* 2010
15 Dec;46(4):537-44.
16
- 17 Csapo R, Alegre LM. Effects of Kinesio(®) taping on skeletal muscle strength-A meta-
18 analysis of current evidence. *J Sci Med Sport.* 2014 Jun 27.
19
- 20 Cunningham R, Uyeshiro Simon A, Preissner K. Occupational Therapy Practice
21 Guidelines for Adults With Multiple Sclerosis. *Am J Occup Ther.*
22 2022;76(5):7605397010. doi:10.5014/ajot.2022.050088
23
- 24 de Brito Brandão M, Mancini MC, Vaz DV, Pereira de Melo AP, Fonseca ST. Adapted
25 version of constraint-induced movement therapy promotes functioning in children with
26 cerebral palsy: a randomized controlled trial. *Clin Rehabil.* 2010 Jul;24(7):639-47.
27
- 28 De Guindos-Sanchez L, Lucena-Anton D, Moral-Munoz JA, et al. The effectiveness of
29 hippotherapy to recover gross motor function in children with cerebral palsy: A
30 systematic review and meta-analysis. *Children (Basel).* 2020;7(9):106
31
- 32 De Miguel A, De Miguel MD, Lucena-Anton D, Rubio MD. Effects of hippotherapy on the
33 motor function of persons with Down's syndrome: A systematic review. *Rev Neurol.*
34 2018;67(7):233-241.
35
- 36 Dionisio MC, Terrill AL. Constraint-Induced Movement Therapy for Infants With or at
37 Risk for Cerebral Palsy: A Scoping Review. *Am J Occup Ther.*
38 2022;76(2):7602205120. doi:10.5014/ajot.2022.047894

- 1 Dixon L, Duncan D, Johnson P, Kirkby L, O’Connell H, Taylor H, Deane KH.
 2 Occupational therapy for patients with Parkinson’s disease. *Cochrane Database Syst*
 3 *Rev.* 2007 Jul 18;(3):CD002813.
 4
- 5 Dromerick AW, Edwards DF, Hahn M. Does the application of constraint-induced
 6 movement therapy during acute rehabilitation reduce arm impairment after ischemic
 7 stroke? *Stroke.* 2000 Dec;31(12):2984-8.
 8
- 9 Dromerick AW, Lang CE, Birkenmeier RL, Wagner JM, Miller JP, Videen TO, et al. Very
 10 Early Constraint-Induced Movement during Stroke Rehabilitation (VECTORS): A
 11 single-center RCT. *Neurology.* 2009 Jul 21;73(3):195-201. Epub 2009 May 20.
 12
- 13 Dry Hydromassage. Princeton, MN: Sidmar Manufacturing, Inc.; 2001-2005. Accessed
 14 April 18, 2023. Available at URL address: <http://www.sidmar.com/>
 15
- 16 Eliasson AC, Krumlinde-sundholm L, Shaw K, Wang C. Effects of constraint-induced
 17 movement therapy in young children with hemiplegic cerebral palsy: an adapted model.
 18 *Dev Med Child Neurol.* 2005 Apr;47(4):266-75.
 19
- 20 Eliasson AC, Krumlinde-Sundholm L, Gordon AM, et al; European network for Health
 21 Technology Assessment (EUnetHTA). Guidelines for future research in constraint-
 22 induced movement therapy for children with unilateral cerebral palsy: An expert
 23 consensus. *Dev Med Child Neurol.* 2014
 24
- 25 Espí-López GV, Serra-Añó P, Vicent-Ferrando J, Sánchez-Moreno-Giner M, Arias-Burúa
 26 JL, Cleland J, Fernández-de-Las-Peñas C. Effectiveness of Inclusion of Dry Needling
 27 in a Multimodal Therapy Program for Patellofemoral Pain: A Randomized Parallel-
 28 Group Trial. *J Orthop Sports Phys Ther.* 2017 Jun;47(6):392-401.
 29
- 30 Fields B, Smallfield S. Occupational Therapy Practice Guidelines for Adults With Chronic
 31 Conditions. *Am J Occup Ther.* 2022;76(2):7602397010. doi:10.5014/ajot.2022/762001
 32
- 33 Forwell S. Occupational therapy practice guidelines for adults with neurodegenerative
 34 diseases. Bethesda (MD): American Occupational Therapy Association (AOTA);
 35 2014.
 36
- 37 Frontera W, Silver J, Rizzo TD editors. *Essentials of physical medicine and rehabilitation.*
 38 3rd ed. Philadelphia, PA: Saunders, an imprint of Elsevier Inc.; 2014.

- 1 García-Pérez P, Rodríguez-Martínez MDC, Lara JP, Cruz-Cosme C. Early Occupational
 2 Therapy Intervention in the Hospital Discharge after Stroke. *Int J Environ Res Public*
 3 *Health*. 2021;18(24):12877. Published 2021 Dec 7. doi:10.3390/ijerph182412877
 4
- 5 Golisz K. Occupational therapy practice guidelines for adults with traumatic brain injury.
 6 Bethesda (MD): American Occupational Therapy Association (AOTA); 2016.
 7
- 8 Grampurohit N, Pradhan S, Kartin D. Efficacy of adhesive taping as an adjunct to physical
 9 rehabilitation to influence outcomes post-stroke: a systematic review. *Top Stroke*
 10 *Rehabil*. 2015 Feb; 22(1):72-82.
 11
- 12 Grotta JC, Noser EA, Ro T, Boake C, Levin H, Aronowski J, Schallert T. Constraint-
 13 induced movement therapy. *Stroke*. 2004 Nov;35(11 Suppl 1):2699-701.
 14
- 15 Guindos-Sanchez L, Lucena-Anton D, Moral-Munoz JA, Salazar A, Carmona-Barrientos
 16 I. The Effectiveness of Hippotherapy to Recover Gross Motor Function in Children
 17 with Cerebral Palsy: A Systematic Review and Meta-Analysis. *Children (Basel)*. 2020
 18 Aug 19;7(9):106.
 19
- 20 H2Omassage System. Winnipeg, MB, Canada. Retrieved on April 18, 2023 from
 21 <http://www.h2omassage.com/>
 22
- 23 Hoare BJ, Imms C, Rawicki HB, Carey L. Modified constraint-induced movement therapy
 24 or bimanual occupational therapy following injection of Botulinum toxin-A to improve
 25 bimanual performance in young children with hemiplegic cerebral palsy: a randomized
 26 controlled trial methods paper. *BMC Neurol*. 2010 Jul 5;10(1):58.
 27
- 28 Hoare BJ, Wasiak J, Imms C, Carey L. Constraint-induced movement therapy in the
 29 treatment of the upper limb in children with hemiplegic cerebral palsy. *Cochrane*
 30 *Database Syst Rev*. 2007 Apr 18;(2):CD004149.
 31
- 32 Hoare BJ, Wallen MA, Thorley MN, Jackman ML, Carey LM, Imms C. Constraint-
 33 induced movement therapy in children with unilateral cerebral palsy. *Cochrane*
 34 *Database of Systematic Reviews* 2019, Issue 4. Art. No.: CD004149. DOI:
 35 10.1002/14651858.CD004149.pub3.
 36
- 37 Hochberg MC, Altman RD, April KT, Benkhalti M, Guyatt G, McGowan J, et al. American
 38 College of Rheumatology. American College of Rheumatology 2012 recommendations

- 1 for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the
- 2 hand, hip, and knee. Arthritis Care Res (Hoboken). 2012 Apr;64(4):465-74.

- 1 Hoffmann T, Bennett S, Koh C, McKenna K. The Cochrane review of occupational therapy
2 for cognitive impairment in stroke patients. *Eur J Phys Rehabil Med.* 2011;47(3):513-
3 519.
- 4
- 5 Huang HH, Fethers L, Hale J, McBride A. Bound for success: a systematic review of
6 constraint-induced movement therapy in children with cerebral palsy supports
7 improved arm and hand use. *Phys Ther.* 2009 Nov;89(11):1126-41.
- 8
- 9 Heussen N, Häusler M. Equine-Assisted Therapies for Children With Cerebral Palsy: A
10 Meta-analysis. *Pediatrics.* 2022;150(1):e2021055229. doi:10.1542/peds.2021-
11 055229
- 12
- 13 Jackman M, Sakzewski L, Morgan C, et al. Interventions to improve physical function for
14 children and young people with cerebral palsy: international clinical practice
15 guideline. *Dev Med Child Neurol.* 2022;64(5):536-549. doi:10.1111/dmcn.15055
- 16
- 17 Jimmo v. Sebelius, No. 5:11-CV-17-CR (D. Vt. filed Jan. 18, 2011).
- 18
- 19 Koomar, J., Burpee, J. D., DeJean, V., Frick, S., Kawar, M. J., & Fischer, D. M. (2001).
20 Theoretical and clinical perspectives on the interactive metronome®: A view from
21 occupational therapy practice. *American Journal of Occupational Therapy*, 55(2),
22 163–166. <https://doi.org/10.5014/ajot.55.2.163>
- 23
- 24 Kraft KA, Weisberg J, Finch MD, et al. Hippotherapy in rehabilitation care for children
25 with neurological impairments and developmental delays: A Case Series. *Pediatr Phys*
26 *Ther.* 2019;31(1):E14-E21.
- 27
- 28 Kundakci B, Kaur J, Goh SL, et al. Efficacy of nonpharmacological interventions for
29 individual features of fibromyalgia: a systematic review and meta-analysis of
30 randomised controlled trials. *Pain.* 2022;163(8):1432-1445.
31 doi:10.1097/j.pain.0000000000002500
- 32
- 33 Lee CW, Kim SG, Yong MS. Effects of hippotherapy on recovery of gait and balance
34 ability in patients with stroke. *J Phys Ther Sci.* 2014 Feb;26(2):309-11.
- 35
- 36 Legg L, Drummond A, Leonardi-Bee J, Gladman JR, Corr S, Donkervoort M, et al.
37 Occupational therapy for patients with problems in personal activities of daily living
38 after stroke: systematic review of randomized trials. *BMJ.* 2007 Nov 3;335(7626):922.

- 1 Legg LA, Drummond AE, Langhorne P. Occupational therapy for patients with problems
2 in activities of daily living after stroke. *Cochrane Database Syst Rev.* 2006 Oct
3 18;(4):CD003585.
- 4
- 5 Legg LA, Lewis SR, Schofield-Robinson OJ, Drummond A, Langhorne P. Occupational
6 therapy for adults with problems in activities of daily living after stroke. *Cochrane*
7 *Database of Systematic Reviews* 2017, Issue 7. Art. No.: CD003585. DOI:
8 10.1002/14651858.CD003585.pub3.
- 9
- 10 Lindsay P, Bayley M, Hellings C, Hill M, Woodbury E, Phillips S. Stroke rehabilitation
11 and community reintegration. Outpatient and community-based rehabilitation. In:
12 *Canadian best practice recommendations for stroke care. CMAJ* 2008 Dec 2;179(12
13 *Suppl*):E58-61.
- 14
- 15 Marquez J, Weerasekara I, Chambers L. Hippotherapy in adults with acquired brain injury:
16 A systematic review. *Physiother Theory Pract.* 2020 Jul;36(7):779-790.
- 17
- 18 Martínez-Costa Montero MC, Cabeza AS. Effectiveness of constraint-induced movement
19 therapy in upper extremity rehabilitation in patients with cerebral palsy: A systematic
20 review. *Rehabilitacion (Madr).* 2020 Nov 30 [Online ahead of print].
- 21
- 22 McIntyre A, Viana R, Janzen S, Mehta S, Pereira S, Teasell R. Systematic review and
23 meta-analysis of constraint-induced movement therapy in the hemiparetic upper
24 extremity more than six months post stroke. *Top Stroke Rehabil.* 2012 Nov-
25 *Dec*;19(6):499-513.
- 26
- 27 Morris ME, Perry A, Bilney B, Curran A, Dodd K, Wittwer JE, Dalton GW. Outcomes of
28 physical therapy, speech pathology, and occupational therapy for people with motor
29 neuron disease: a systematic review. *Neurorehabil Neural Repair.* 2006 Sep;20(3):424-
30 34.
- 31
- 32 Nair HKR. Microcurrent as an adjunct therapy to accelerate chronic wound healing and
33 reduce patient pain. *J Wound Care.* 2018 May 2;27(5):296-306.
- 34
- 35 Nijland R, Kwakkel G, Bakers J, van Wegen E. Constraint-induced movement therapy for
36 the upper paretic limb in acute or sub-acute stroke: a systematic review. *Int J Stroke.*
37 2011 Oct;6(5):425-33.

- 1 Novak I, Morgan C, Fahey M, Finch-Edmondson M, Galea C, Hines A, Langdon K,
2 Namara MM, Paton MC, Popat H, Shore B, Khamis A, Stanton E, Finemore OP, Tricks
3 A, Te Velde A, Dark L, Morton N, Badawi N. State of the Evidence Traffic Lights
4 2019: Systematic Review of Interventions for Preventing and Treating Children with
5 Cerebral Palsy. *Curr Neurol Neurosci Rep.* 2020 Feb 21;20(2):3.
6
- 7 O'Haire ME. Animal-assisted intervention for autism spectrum disorder: a systematic
8 literature review. *J Autism Dev Disord.* 2013 Jul;43(7):1606-22.
9
- 10 Ottawa Panel. Ottawa panel evidence-based clinical practice guidelines for therapeutic
11 exercises and manual therapy in the management of osteoarthritis. *Phys Ther* 2005
12 Sep;85(9):907-71.
13
- 14 Ottawa Panel. Ottawa Panel evidence-based clinical practice guidelines for therapeutic
15 exercises in the management of rheumatoid arthritis in adults. *Phys Ther.* 2004
16 Oct;84(10):934-72.
17
- 18 Ottawa Panel, Khadilkar A, Phillips K, Jean N, Lamothe C, Milne S, Sarnecka J. Ottawa
19 panel evidence-based clinical practice guidelines for post-stroke rehabilitation. *Top*
20 *Stroke Rehabil.* 2006 Spring;13(2):1-269.
21
- 22 Pantera E, Froment P, Vernay D. Does Hippotherapy Improve the Functions in Children
23 with Cerebral Palsy? Systematic Review Based on the International Classification of
24 Functioning. *J Integr Complement Med.* 2022;28(9):705-720.
25 doi:10.1089/jicm.2021.0417
26
- 27 Peurala SH, Kantanen MP, Sjögren T, Paltamaa J, Karhula M, Heinonen A. Effectiveness
28 of constraint-induced movement therapy on activity and participation after stroke: a
29 systematic review and meta-analysis of randomized controlled trials. *Clin Rehabil.*
30 2012 Mar;26(3):209-23.
31
- 32 Pollock A, Baer G, Campbell P, Choo PL, Forster A, Morris J, et al. Physical rehabilitation
33 approaches for the recovery of function and mobility following stroke. *Cochrane*
34 *Database Syst Rev.* 2014 Apr 22;4:CD001920.
35
- 36 Prieto A, Martins Almeida Ayupe K, Nemetala Gomes L, Saúde AC, Gutierrez Filho P.
37 Effects of equine-assisted therapy on the functionality of individuals with disabilities:
38 systematic review and meta-analysis. *Physiother Theory Pract.* 2022;38(9):1091-1106.
39 doi:10.1080/09593985.2020.1836694

- 1 Pulman J, Buckley E, Clark-Carter D. A meta-analysis evaluating the effectiveness of two
 2 different upper limb hemiparesis interventions on improving health-related quality of
 3 life following stroke. *Top Stroke Rehabil.* 2013 Mar-Apr;20(2):189-96.
 4
- 5 Quinn É, Hynes SM. Occupational therapy interventions for multiple sclerosis: A scoping
 6 review. *Scand J Occup Ther.* 2021;28(5):399-414.
 7
- 8 Ramey SL, DeLuca SC, Stevenson RD, et al. Constraint-Induced Movement Therapy for
 9 Cerebral Palsy: A Randomized Trial. *Pediatrics.* 2021;148(5):e2020033878.
 10 doi:10.1542/peds.2020-033878
 11
- 12 Revised by the Commission on Practice, 2020:, Casto SC, Davis C, et al. Standards of
 13 Practice for Occupational Therapy. *Am J Occup Ther.*
 14 2022;75(Supplement_3):7513410030. doi:10.5014/ajot.2021.75S3004
 15
- 16 Sabari J, Lieberman D. Occupational therapy practice guidelines for adults with stroke.
 17 Bethesda (MD): American Occupational Therapy Association (AOTA); 2015.
 18
- 19 Sakzewski L, Gordon A, Eliasson AC. The state of the evidence for intensive upper limb
 20 therapy approaches for children with unilateral cerebral palsy. *J Child Neurol.* 2014
 21 Aug;29(8):1077-90.
 22
- 23 Sakzewski L, Ziviani J, Boyd RN. Efficacy of upper limb therapies for unilateral cerebral
 24 palsy: a meta-analysis. *Pediatrics.* 2014 Jan;133(1):e175-204
 25
- 26 Sall, J., Eapen, B. C., Tran, J. E., Bowles, A. O., Bursaw, A., & Rodgers, M. E. (2019).
 27 The Management of Stroke Rehabilitation: A Synopsis of the 2019 U.S. Department
 28 of Veterans Affairs and U.S. Department of Defense Clinical Practice Guideline.
 29 *Annals of internal medicine*, 171(12), 916–924. <https://doi.org/10.7326/M19-1695>
 30
- 31 Santos de Assis G, Schlichting T, Rodrigues Mateus B, Gomes Lemos A, Dos Santos AN.
 32 Physical therapy with hippotherapy compared to physical therapy alone in children
 33 with cerebral palsy: systematic review and meta-analysis. *Dev Med Child Neurol.*
 34 2022;64(2):156-161. doi:10.1111/dmcn.15042
 35
- 36 Scottish Intercollegiate Guidelines Network (SIGN). Management of patients with stroke:
 37 rehabilitation, prevention and management of complications, and discharge planning.
 38 A national clinical guideline. Edinburgh (Scotland): Scottish Intercollegiate Guidelines
 39 Network (SIGN); 2010 Jun (Updated 15 October 2014).

- 1 Shi YX, Tian JH, Yang KH, Zhao Y. Modified constraint-induced movement therapy
2 versus traditional rehabilitation in patients with upper-extremity dysfunction after
3 stroke: a systematic review and meta-analysis. *Arch Phys Med Rehabil.* 2011
4 Jun;92(6):972-82.
- 5
- 6 Silberstein N. Dry hydrotherapy: don't add water. *Rehab Manag.* 2006 Jun;19(5):22-3.
- 7
- 8 Silkwood-Sherer DJ, Killian CB, Long TM, Martin KS. Hippotherapy—an intervention to
9 habilitate balance deficits in children with movement disorders: a clinical trial. *Phys
10 Ther.* 2012 May;92(5):707-17.
- 11
- 12 Singh P, Pradhan B. Study to assess the effectiveness of modified constraint-induced
13 movement therapy in stroke subjects: A randomized controlled trial. *Ann Indian Acad
14 Neurol.* 2013 Apr;16(2):180-4.
- 15
- 16 Sirtori V, Corbetta D, Moja L, Gatti R. Constraint-induced movement therapy for upper
17 extremities in stroke patients. *Cochrane Database Syst Rev.* 2009 Oct 7;(4):CD004433.
- 18
- 19 Steultjens EM, Dekker J, Bouter LM, Cardol M, Van de Nes JC, Van den Ende CH.
20 Occupational therapy for multiple sclerosis. *Cochrane Database Syst Rev.*
21 2003;(3):CD003608.
- 22
- 23 Steultjens EM, Dekker J, Bouter LM, Leemrijse CJ, van den Ende CH. Evidence of the
24 efficacy of occupational therapy in different conditions: an overview of systematic
25 reviews. *Clin Rehabil.* 2005 May;19(3):247-54.
- 26
- 27 Steultjens EM, Dekker J, Bouter LM, van de Nes JC, van de Ende CH. Occupational
28 therapy for stroke patients. A systematic review. *Stroke.* 2003 Mar;34(3):676-87.
- 29
- 30 Steultjens EM, Dekker J, Bouter LM, van Schaardenburg D, van Kuyk MA, van den Ende
31 CH. Occupational therapy for rheumatoid arthritis. *Cochrane Database Syst Rev.*
32 2004;(1):CD003114.
- 33
- 34 Stevenson T, Thalman L, Christie H, Poluha W. Constraint-Induced Movement Therapy
35 Compared to Dose-Matched Interventions for Upper-Limb Dysfunction in Adult
36 Survivors of Stroke: A Systematic Review with Meta-analysis. *Physiother Can.* 2012
37 Fall;64(4):397-413.
- 38

- 1 Sung IY, Ryu JS, Pyun SB, Yoo SD, Song WH, Park MJ. Efficacy of forced-use therapy
- 2 in hemiplegic cerebral palsy. Arch Phys Med Rehabil. 2005 Nov;86(11):2195-8.

- 1 Taub E, Ramey SL, DeLuca S, Echols K. Efficacy of constraint-induced movement therapy
2 for children with cerebral palsy with asymmetric motor impairment. *Pediatrics*. 2004
3 Feb;113(2):305-12.
4
- 5 Trombly CA, Ma HI. A synthesis of the effects of occupational therapy for persons with
6 stroke, Part I: Restoration of roles, tasks, and activities. *Am J Occup Ther*. 2002 May-
7 Jun;56(3):250-9.
8
- 9 Walker MF, Leonardi-Bee J, Bath P, Langhorne P, Dewey M, Corr S, et al. Individual
10 patient data meta-analysis of randomized controlled trials of community occupational
11 therapy for stroke patients. *Stroke*. 2004 Sep;35(9):2226-32.
12
- 13 Walker C, Shierk A, Roberts H. Constraint Induced Movement Therapy in Infants and
14 Toddlers with Hemiplegic Cerebral Palsy: A Scoping Review. *Occup Ther Health
15 Care*. 2022;36(1):29-45. doi:10.1080/07380577.2021.1953206
16
- 17 White E, Zippel J, Kumar S. The effect of equine-assisted therapies on behavioural,
18 psychological and physical symptoms for children with attention deficit/hyperactivity
19 disorder: A systematic review. *Complement Ther Clin Pract*. 2020 May;39:101101.
20
- 21 Wolf SL, Winstein CJ, Miller JP, Taub E, Uswatte G, Morris D, et al; EXCITE
22 Investigators. Effect of constraint-induced movement therapy on upper extremity
23 function 3 to 9 months after stroke: the EXCITE randomized clinical trial. *JAMA*. 2006
24 Nov 1;296(17):2095-104.
25
- 26 Wood J, Henderson W, Foster ER. Occupational Therapy Practice Guidelines for People
27 With Parkinson's Disease. *Am J Occup Ther*. 2022;76(3):7603397010.
28 doi:10.5014/ajot.2022.763001
29
- 30 Yu H, Côté P, Shearer HM, Wong JJ, Sutton DA, Randhawa KA, Varatharajan S, Southerst
31 D, Mior SA, Ameis A, Stupar M, Nordin M, van der Velde GM, Carroll L, Jacobs CL,
32 Taylor-Vaisey AL, Abdulla S, Shergill Y. Effectiveness of passive physical modalities
33 for shoulder pain: systematic review by the Ontario protocol for traffic injury
34 management collaboration. *Phys Ther*. 2015 Mar;95(3):306-18.