

1 **Clinical Practice Guideline: Unhealthy Alcohol / Substance Use Screening and**
 2 **Intervention**

3
 4 **Date of Implementation: April 19, 2012**

5
 6 **Product: Specialty**
 7

8
 9 **GUIDELINES**

10 Unhealthy alcohol / substance use screenings and interventions by an appropriately trained
 11 healthcare professional are established as clinically effective, are professionally
 12 recognized, and have a favorable benefit: risk profile.

13
 14 **INTRODUCTION**

15 Substance Use Disorders are characterized by specific behavioral and physical symptoms,
 16 often including impaired control over substance use, social impairment, risky use,
 17 withdrawal, tolerance, and craving (DSM-V). The causes of substance-related disorders
 18 are multifaceted involving genetic, neurobiological, and psychosocial factors. The early
 19 signs of substance-related disorders may not always be evident to health care practitioners,
 20 but unhealthy substance use screening can help identify impaired control, social
 21 impairment, risky use, and pharmacological criteria. Treatment frequently involves detox,
 22 individual or group counseling, mutual support groups, and/or medications, however,
 23 prevention of the development of substance-related disorders is preferred to reduce
 24 healthcare costs and human suffering (National Institute on Alcohol Abuse and
 25 Alcoholism, 2021).

26
 27 **Unhealthy Alcohol Use**

28 Unhealthy alcohol use is associated with adverse effects on physical and mental health and
 29 is involved in socioeconomic issues as well. It is one of the leading causes of preventable
 30 death in the United States. (AHRQ 2024). Unhealthy alcohol use encompasses a wide
 31 range of behaviors, from drinking above the recommended limits (i.e., risky drinking) to
 32 alcohol use disorder (AUD). Types of unhealthy alcohol use and related terms are
 33 numerous and are not mutually exclusive. Examples include risky/at-risk use, binge
 34 drinking, excessive use, harmful use, and alcohol use disorder.

35
 36 **Definitions**

37 Unhealthy Alcohol Use: Unhealthy alcohol use includes any alcohol use that puts a
 38 person’s health or safety at risk or causes other alcohol-related problems. Includes binge
 39 drinking.

1 Binge Drinking: Binge drinking is when a male has five or more drinks in two hours, or a
 2 female has four or more drinks within two hours.

3
 4 Risky Use: The National Institute on Alcohol Abuse and Alcoholism (NIAAA) defines
 5 risky use as exceeding the recommended limits of 4 drinks per day (56 g/d based on the
 6 US standard of 14 g/drink) or 14 drinks per week (196 g/d) for healthy adult men aged 21
 7 to 64 years or 3 drinks per day or 7 drinks per week (42 g/d or 98 g/week) for all adult
 8 women of any age and men 65 years or older.

9
 10 Excessive Alcohol Use: Excessive alcohol use is defined by the CDC as binge drinking
 11 ($\geq 5/4$ drinks per occasion for men/women), heavy drinking ($\geq 15/8$ drinks per week for
 12 men/women), and any alcohol use by people younger than age 21 years and by pregnant
 13 women.

14
 15 Harmful Use: A pattern of alcohol consumption causing health problems directly related
 16 to alcohol.

17
 18 Alcohol Use Disorder: The National Institute on Alcohol Abuse and Alcoholism (NIAAA)
 19 defines Alcohol use disorder (AUD) as a medical condition (a brain disorder) characterized
 20 by an impaired ability to stop or control alcohol use despite adverse social, occupational or
 21 health consequences. DSM V defines alcohol use disorder as, “A problematic pattern of
 22 alcohol use leading to clinically significant impairment or distress, as manifested by at least
 23 two of the following, occurring within a 12-month period: Craving, or a strong desire or
 24 urge to use alcohol.”

25
 26 Standard Drink: Per the NIAAA, examples of a standard drink include a 12-oz beer (5%
 27 alcohol), 5 oz of wine (12% alcohol), or 1.5 oz of distilled spirits (40% alcohol).

28
 29 **USPSTF Recommendation Levels:**

Grade	Definition	Suggestions for Practice
A	The USPSTF <i>recommends</i> the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
B	The USPSTF <i>recommends</i> the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
C	The USPSTF recommends <i>selectively</i> offering or providing this service based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small.	Offer or provide this service for selected patients depending on circumstances.

Grade	Definition	Suggestions for Practice
D	The USPSTF recommends <i>against</i> the service. There is moderate or high certainty of either no net benefit or that the harms outweigh the benefits.	Discourage the use of this service.
I	The USPSTF concludes that the current evidence is <i>insufficient</i> to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.	If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

A comprehensive review of the USPSTF rating process can be found in the ASH policy *Preventive Care Guidelines* (CPG 140 – S) or at the USPSTF website: <http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions>.

USPSTF Recommendations:

Unhealthy Alcohol Use in Adolescents and Adults: Screening and Behavioral Counseling Interventions
Unhealthy Alcohol Use (2018)

Grade B Recommendation: The U.S. Preventive Services Task Force (USPSTF) recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women, and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use.

Grade I Recommendation: The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening and brief behavioral counseling interventions for alcohol use in primary care settings in adolescents aged 12 to 17 years.

UNHEALTHY ALCOHOL USE

Benefits of Early Detection and Behavioral Counseling:
 The USPSTF found no studies that directly evaluated whether screening for unhealthy alcohol use in primary care settings in adolescents and adults, including pregnant women, leads to reduced unhealthy alcohol use; improved risky behaviors; or improved health, social, or legal outcomes.

The USPSTF found adequate evidence that brief behavioral counseling interventions in adults who screen positive are associated with reduced unhealthy alcohol use. There were reductions in both the odds of exceeding recommended drinking limits and heavy use episodes at 6- to 12-month follow-up. In pregnant women, brief counseling interventions increased the likelihood that women remained abstinent from alcohol use during pregnancy. The magnitude of these benefits is moderate. Epidemiologic literature links reductions in alcohol use with reductions in risk for morbidity and mortality and provides

1 indirect support that reduced alcohol consumption may help improve some health
2 outcomes.

3
4 The USPSTF found inadequate evidence that brief behavioral counseling interventions in
5 adolescents were associated with reduced alcohol use. However, no harms were noted for
6 brief behavioral counseling in the evidence.

7
8 Screening Tools—Unhealthy Alcohol Use:

9 Of the available screening tools, the USPSTF determined that 1- to 3-item screening
10 instruments have the best accuracy for assessing unhealthy alcohol use in adults 18 years
11 or older. These instruments include the abbreviated Alcohol Use Disorders Identification
12 Test–Consumption (AUDIT-C) and the NIAAA-recommended Single Alcohol Screening
13 Question (SASQ). When patients screen positive on a brief screening instrument (e.g.,
14 SASQ or AUDIT-C), clinicians should ensure follow-up with a more in-depth risk
15 assessment to confirm unhealthy alcohol use and determine the next steps of care. Evidence
16 supports the use of brief instruments with higher sensitivity and lower specificity as initial
17 screening, followed by a longer instrument with greater specificity (e.g., AUDIT). The
18 AUDIT has 10 questions: 3 questions covering frequency of alcohol use, typical amount
19 of alcohol use, and occasions of heavy use, and 7 questions on the signs of alcohol
20 dependence and common problems associated with alcohol use (e.g., being unable to stop
21 once you start drinking). It requires approximately 2 to 5 minutes to administer. If AUDIT
22 is used as an initial screening test, clinicians may use a lower cutoff (such as 3, 4, or 5) to
23 balance sensitivity and specificity in screening for the full spectrum of unhealthy alcohol
24 use.

25
26 Behavioral Counseling Interventions:

27 Behavioral counseling interventions for unhealthy alcohol use vary in their specific
28 components, administration, length, and number of interactions. They may include
29 cognitive behavioral strategies, such as action plans, drinking diaries, stress management,
30 or problem solving. Interventions may be delivered by face-to-face sessions, written self-
31 help materials, computer- or Web-based programs, or telephone counseling. For the
32 purposes of this recommendation statement, the USPSTF uses the following definitions of
33 intervention intensity: very brief single contact (≤ 5 minutes), brief single contact (6 to 15
34 minutes), brief multicontact (each contact is 6 to 15 minutes), and extended multicontact
35 (≥ 1 contact, each > 15 minutes). Thirty percent of the interventions reviewed by the
36 USPSTF were web-based. Nearly all of the interventions consisted of 4 or fewer sessions;
37 the median number of sessions was 1 (range, 0-21). The median length of time of contact
38 was 30 minutes (range, 1-600 minutes). Most of the interventions had a total contact time
39 of 2 hours or less. Primary care settings often used the Screening, Brief Intervention, and
40 Referral to Treatment (SBIRT) approach. Most interventions involved giving general
41 feedback to participants (e.g., how their drinking fits with recommended limits, or how to
42 reduce alcohol use). The most commonly reported intervention component was use of

1 personalized normative feedback sessions, in which participants were shown how their
 2 alcohol use compares with that of others; more than half of the included trials and almost
 3 all trials in young adults used this technique. Personalized normative feedback was often
 4 combined with motivational interviewing or more extensive cognitive behavioral
 5 counseling. Other cognitive behavioral strategies, such as drinking diaries, action plans,
 6 alcohol use “prescriptions,” stress management, or problem solving were also frequently
 7 used. The USPSTF was unable to identify specific intervention characteristics or
 8 components that were clearly associated with improved outcomes.

9
 10 The USPSTF found no studies that directly evaluated whether screening for unhealthy
 11 alcohol use in primary care settings in adolescents and adults, including pregnant women,
 12 leads to reduced unhealthy alcohol use; improved risky behaviors; or improved health,
 13 social, or legal outcomes.

14 **UNHEALTHY DRUG USE: Screening**

15 **Unhealthy Drug Use (2020)**

16
 17 ***Grade B Recommendation:*** The USPSTF recommends screening by asking questions
 18 about unhealthy drug use in adults age 18 years or older. Screening should be
 19 implemented when services for accurate diagnosis, effective treatment, and
 20 appropriate care can be offered or referred. (Screening refers to asking questions
 21 about unhealthy drug use, not testing biological specimens.)
 22

23
 24 ***Grade I Recommendation:*** The USPSTF concludes that the current evidence is
 25 insufficient to assess the balance of benefits and harms of screening for unhealthy
 26 drug use in adolescents.
 27

28 **Screening Tools for Unhealthy Drug Use**

29 Several screening tools that ask questions about drug use are available for identifying 1 or
 30 more classes of unhealthy drug use, the frequency or severity of use, or drug-related health,
 31 social, or legal consequences that characterize unhealthy use or drug use disorders.
 32 Interviewer-administered tools and self-administered tools appear to have similar accuracy.
 33 Screening tools are not meant to diagnose drug dependence, abuse, addiction, or drug use
 34 disorders. Patients with positive screening results may, therefore, need to be offered or
 35 referred for diagnostic assessment. Brief tools (e.g., NIDA [National Institute on Drug
 36 Abuse] Quick Screen, which asks 4 questions about use of alcohol, tobacco, nonmedical
 37 use of prescription drugs, and illegal drugs in the past year) may be more feasible in busy
 38 primary care settings, but longer tools (e.g., the 8-item ASSIST [Alcohol, Smoking and
 39 Substance Involvement Screening Test]) that assess risks associated with unhealthy drug
 40 use or comorbid conditions may reveal information signaling the need for prompt
 41 diagnostic assessment. Tools with questions about nonmedical use of prescription drugs
 42 (e.g., TAPS [Tobacco, Alcohol, Prescription Medication, and Other Substance Use]) may

1 be useful when clinicians are concerned about prescription misuse. One study reported that
 2 drug use questions in the PRO (Prenatal Risk Overview) risk assessment tool were
 3 reasonably accurate for identifying drug abuse or dependence in pregnant women.

4
 5 Other screening tools for drug use include DAST (Drug Abuse Screening Test), a 10 item
 6 tool that takes five minutes to administer. And SQDS (Single-question drug screening,
 7 ASSIST (Alcohol, Smoking and Substance Involvement Screening Test) and TAPS
 8 (Tobacco, alcohol, prescription medication, and other substance use) may be helpful as
 9 they cover use of multiple substances in one tool.

10 **CPT Code and Documentation Requirements to Substantiate Medical Necessity**

11 When performing an Alcohol/Substance Screening and Intervention (CPT codes 99408 and
 12 99409) the documentation in the medical records should identify the use of a validated
 13 and/or standardized screening questionnaire and the nature and character of the
 14 intervention. Please see the Practitioner Resource section below for examples of screening
 15 and intervention guidelines.

16
 17
 18 Medical record documentation should reflect what screening tools were utilized and that
 19 the results were reviewed with the patient. The practitioner should provide feedback on the
 20 screening results, advice on safe consumption limits and recommendations about behavior
 21 change.

22
 23 If unhealthy substance use is identified through the screening process, the practitioner
 24 should provide, at a minimum, feedback about screening results and advice on safe
 25 consumption limits and recommended behavior change. A referral for chemical
 26 dependency treatment should also be offered or patient's response and plan documented.

CPT Code	Description
99408	Alcohol and/or substance abuse structured screening and brief intervention services; 15 to 30 minutes
99409	Alcohol and/or substance abuse structured screening and brief intervention services; greater than 30 minutes

28 **Practitioner Resources**

- 29 • CDC's Alcohol Screening and Brief Intervention Efforts
 30 (<https://www.cdc.gov/ncbddd/fasd/alcohol-screening.html>)
- 31 • National Institutes on Alcohol Abuse and Alcoholism (<http://www.niaaa.nih.gov>)
- 32 • Substance Abuse and Mental Health Services (<https://www.samhsa.gov/>)
- 33 • National Institute of Drug Abuse (<https://nida.nih.gov/>)
- 34 • Institute for Research, Education, and Training in Addictions (<http://ireta.org/>)
- 35

- Healthy People (<https://health.gov/healthypeople/objectives-and-data/browse-objectives/drug-and-alcohol-use>)

Member Resources

Educating patients about substance abuse disorders, treatment options and available resources can assist the recovery process. Publicly available resources can be found at:

- Substance Abuse & Mental Health Services (<https://www.samhsa.gov/>) 1-800-662-HELP (4357)
- National Institute on Alcohol Abuse and Alcoholism (<https://www.niaaa.nih.gov>)
- Alcoholics Anonymous Number – 1-212-870-3400
- National Council on Alcoholism and Drug Dependence, Inc. (NCADD) 1-800-NCA-CALL (1-800-622-2255)
- Suicide and Crisis Lifeline (mental health, alcohol or substance use concerns) call or text 988

PRACTITIONER SCOPE AND TRAINING

Practitioners should practice only in the areas in which they are competent based on their education, training and experience. Levels of education, experience, and proficiency may vary among individual practitioners. It is ethically and legally incumbent on a practitioner to determine where they have the knowledge and skills necessary to perform such services and whether the services are within their scope of practice.

It is *best practice* for the practitioner to appropriately render services to a patient only if they are trained to competency, equally skilled, and adequately competent to deliver a service compared to others trained to perform the same procedure. If the service would be most competently delivered by another health care practitioner who has more skill and training, it would be best practice to refer the patient to the more expert practitioner.

Best practice can be defined as a clinical, scientific, or professional technique, method, or process that is typically evidence-based and consensus driven and is recognized by a majority of professionals in a particular field as more effective at delivering a particular outcome than any other practice (Joint Commission International Accreditation Standards for Hospitals, 2020).

Depending on the practitioner’s scope of practice, training, and experience, a patient’s condition and/or symptoms during examination or the course of treatment may indicate the need for referral to another practitioner or even emergency care. In such cases it is essential for the practitioner to refer the patient for appropriate co-management (e.g., to their primary care physician) or if immediate emergency care is warranted, to contact 911 as appropriate. See the *Managing Medical Emergencies in a Health Care Facility (CPG 159 – S)* clinical practice guideline for information.

1 **References**

2 American Academy of Family Physicians. (2023). *Addressing Alcohol and Other*
 3 *Substance Use Practice Manual*. Retrieved December 27, 2024 from
 4 [https://www.aafp.org/content/dam/AAFP/documents/patient_care/alcohol/alcohol%20](https://www.aafp.org/content/dam/AAFP/documents/patient_care/alcohol/alcohol%20practice%20manual%202023.pdf)
 5 [practice%20manual%202023.pdf](https://www.aafp.org/content/dam/AAFP/documents/patient_care/alcohol/alcohol%20practice%20manual%202023.pdf)

6
 7 American Medical Association. (current year). *Current Procedural Terminology (CPT)*
 8 *Current year* (rev. ed.). Chicago: AMA

9
 10 American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental*
 11 *disorders* (5th ed.)

12
 13 EvidenceNOW: Managing Unhealthy Alcohol Use. Content last reviewed October 2024.
 14 Agency for Healthcare Research and Quality, Rockville, MD.
 15 <https://www.ahrq.gov/evidencenow/projects/alcohol/index.html>

16
 17 Joint Commission International. (2020). *Joint Commission International Accreditation*
 18 *Standards for Hospitals* (7th ed.): Joint Commission Resources

19
 20 Kowalchuk A, Gonzalez SJ, Zoorob RJ. Substance Misuse in Adults: A Primary Care
 21 Approach. *Am Fam Physician*. 2024 May;109(5):430-440. PMID: 38804757

22
 23 National Institute on Alcohol Abuse and Alcoholism. (2024). *Alcohol Facts and Statistics*.
 24 Retrieved on December 27, 2024 from
 25 [https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/alcohol-facts-and-](https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/alcohol-facts-and-statistics)
 26 [statistics](https://www.niaaa.nih.gov/publications/brochures-and-fact-sheets/alcohol-facts-and-statistics)

27
 28 O'Connor, E.A., Perdue, L.A., Senger, C.A., Rushkin, M., Patnode, C.D., Bean, S.I., &
 29 Jonas, D.E. (2018). Screening and Behavioral Counseling Interventions to Reduce
 30 Unhealthy Alcohol Use in Adolescents and Adults: An Updated Systematic Review for
 31 the U.S. Preventive Services Task Force. *JAMA*. 2018;320(18):1910–1928

32
 33 Patnode, C. D., Perdue, L. A., Rushkin, M., Dana, T., Blazina, I., Bougatsos, C., Grusing,
 34 S., O'Connor, E. A., Fu, R., & Chou, R. (2020). Screening for Unhealthy Drug Use:
 35 Updated Evidence Report and Systematic Review for the US Preventive Services Task
 36 Force. *JAMA*, 323(22), 2310–2328. <https://doi.org/10.1001/jama.2019.21381>

37
 38 U.S. Preventive Health Services Task Force (USPSTF). (2018). *Unhealthy Alcohol Use in*
 39 *Adolescents and Adults: Screening and Behavioral Counseling Interventions*.
 40 Retrieved December 27, 2024, from
 41 <https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStat>

1 ementFinal/unhealthy-alcohol-use-in-adolescents-and-adults-screening-and-
2 behavioral-counseling-interventions

3

4 U.S. Preventive Health Services Task Force (USPSTF). (2020). *Unhealthy Drug Use:*
5 *Screening*. Retrieved December 27, 2024, from
6 [http://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/](http://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/drug-use-illicit-screening)
7 drug-use-illicit-screening