

1 **Clinical Practice Guideline: Medical Nutrition Therapy for Weight**
 2 **Management**

4 **Date of Implementation: April 19, 2012**

6 **Product: Specialty**

9 **GUIDELINES**

10 American Specialty Health – Specialty (ASH) considers Medical Nutrition Therapy
 11 (MNT) and patient specific nutritional evaluation and counseling for weight management
 12 in which appropriate diet and eating habits are essential to the overall treatment program
 13 provided by a qualified healthcare professional as medically necessary.

15 **INTRODUCTION**

16 Practitioners and the American public are grappling with the obesity epidemic. Medical
 17 nutrition therapy offers a science-based, expert guided framework and specific tools to help
 18 obese individuals develop skills and behaviors to promote weight loss and maintain stable
 19 healthy weight in the long term.

21 The USPSTF concluded with moderate certainty that offering or referring adults with
 22 obesity to intensive behavioral interventions or behavior-based weight loss maintenance
 23 interventions has a moderate net benefit. Effective behavioral interventions targeted
 24 achieving and maintaining weight loss greater than or equal to 5%. Most interventions
 25 studied combined dietary changes and increased physical activity and lasted for 1-2 years,
 26 with the majority having ≥ 12 sessions in the first year. The Guidelines note the US Food
 27 and Drug Administration (FDA) considers a weight loss of 5% as clinically important.

29 **SCREENING RECOMMENDATIONS**

30 Applicable recommendations for the preventive health screenings covered in this policy
 31 are based on the United States Preventive Services Task Force (USPSTF).
 32

Grade	Definition	Suggestions for Practice
A	The USPSTF <i>recommends</i> the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
B	The USPSTF <i>recommends</i> the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.

Grade	Definition	Suggestions for Practice
C	The USPSTF recommends <i>selectively</i> offering or providing this service based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small.	Offer or provide this service for selected patients depending on circumstances.
D	The USPSTF recommends <i>against</i> the service. There is moderate or high certainty of either no net benefit or that the harms outweigh the benefits.	Discourage the use of this service.
I	The USPSTF concludes that the current evidence is <i>insufficient</i> to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.	If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

1
2 A comprehensive review of the USPSTF rating process can be found in the ASH policy
3 *Preventive Care Guidelines* (CPG 140 – S) or at the USPSTF website:
4 <http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions>.
5

6 **USPSTF Recommendations (2018) – Weight Loss to Prevent Obesity-Related**
7 **Morbidity and Mortality in Adults: Behavioral Interventions**

8
9 **Grade B Recommendation:**

10 The USPSTF recommends that clinicians offer or refer adults with a body mass index
11 (BMI) of 30 or higher (calculated as weight in kilograms divided by height in meters
12 squared) to intensive, multicomponent behavioral interventions.
13

14 **USPSTF Recommendations (2024) – High Body Mass Index in Children and**
15 **Adolescents: Interventions**

16
17 **For children and adolescents 6 years or older, the USPSTF recommends that**
18 **clinicians provide or refer children and adolescents 6 years or older with a high**
19 **body mass index (BMI) (\geq 95th percentile for age and sex) to comprehensive,**
20 **intensive behavioral interventions. SCREENING TOOLS**

21
22 **Adults**

23 The USPSTF has determined measurement of BMI by the practitioner as the appropriate
24 screening method for obesity/overweight. There are several online BMI calculators
25 available upon searching.

1 BMI of 25-29.9 kg/m² indicates overweight and BMI ≥ 30 kg/m² indicates obesity.
 2 Obesity is further differentiated into 3 classes: I = BMI 30-34.9 kg/m² (obese); II =BMI
 3 35-39.9 kg/m² (severely obese); III =BMI 40+ kg/m² (morbidly obese).

4
 5 Waist circumference maybe an acceptable alternative to BMI measurement in some patient
 6 sub-populations.

7 8 **Children And Adolescents**

9 The USPSTF is using the following terms to define categories of increased BMI:
 10 overweight is defined as an age- and sex-specific BMI between the 85th and 95th
 11 percentiles, and obesity is defined as an age- and sex-specific BMI at ≥95th percentile.

12 13 **DOCUMENTATION REQUIREMENTS TO SUBSTANTIATE MEDICAL** 14 **NECESSITY**

15 Short term evaluation and counseling should include performing an initial dietary
 16 evaluation, counseling the patient about sample menu planning, and teaching the patient
 17 the impact of diet on their health condition(s). The goals of MNT are to promote health,
 18 reduce the incidence of preventable disease and improve quality of life.

19
 20 The 5A's is a framework frequently used in clinical practice and should be documented
 21 within the clinical record to guide behavioral interventions.

- 22 1. Assess the health risk
- 23 2. Advise the patient on behavior change
- 24 3. Agree collaboratively with patient on an action plan
- 25 4. Assist the patient in making changes and adhering to the plan
- 26 5. Arrange follow-up

27
 28 The practitioner's medical record should also reflect:

- 29 • Performance of a nutrition assessment determining the nutrition diagnosis
- 30 • BMI measurement
- 31 • Identification of treatment goals
- 32 • Planning and implementing a nutrition intervention that is culturally appropriate
 33 and uses evidence-based nutrition practice guidelines
- 34 • Development of a nutritional recommendation/plan
- 35 • Monitoring and evaluating an individual's progress over subsequent visits with the
 36 clinician
- 37 • Establishment of a patient's self-management training and goal setting
- 38 • Nutrition intervention most appropriate for the management or treatment of
 39 patients' condition are chosen after review of all available data

1 PRACTITIONER RESOURCES

- 2 • Tool to identify MNT professionals:
 - 3 ○ Academy of Nutrition and Dietetics: <https://www.eatright.org/find-an-expert>
- 4 • Tools to offer for assessing health risk (waist circumference and BMI):
 - 5 ○ BMI: https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmicalc.htm
 - 6 ○ Child and Teen BMI Calculator: <https://www.cdc.gov/bmi/child-teen-calculator/index.html>

9 MEMBER RESOURCES

- 10 • myplate.gov: <https://www.myplate.gov/resources/tools>
- 11 • <https://www.eatright.org/food>
- 12 • <https://healthy10challenge.org> (interactive 10 week program that focuses on
- 13 building in healthy food and activity habits)
- 14 • <https://www.fda.gov/food/new-nutrition-facts-label/whats-new-nutrition-facts-label>

17 PRACTITIONER SCOPE AND TRAINING

18 Practitioners should practice only in the areas in which they are competent based on their
 19 education training and experience. Levels of education, experience, and proficiency may
 20 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
 21 to determine where they have the knowledge and skills necessary to perform such services
 22 and whether the services are within their scope of practice.

24 It is best practice for the practitioner to appropriately render services to a patient only if
 25 they are trained to competency, equally skilled, and adequately competent to deliver a
 26 service compared to others trained to perform the same procedure. If the service would be
 27 most competently delivered by another health care practitioner who has more skill and
 28 training, it would be best practice to refer the patient to the more expert practitioner.

30 Best practice can be defined as a clinical, scientific, or professional technique, method, or
 31 process that is typically evidence-based and consensus driven and is recognized by a
 32 majority of professionals in a particular field as more effective at delivering a particular
 33 outcome than any other practice (Joint Commission International Accreditation Standards
 34 for Hospitals, 2020).

36 Depending on the practitioner’s scope of practice, training, and experience, a patient’s
 37 condition and/or symptoms during examination or the course of treatment may indicate the
 38 need for referral to another practitioner or even emergency care. In such cases it is essential
 39 for the practitioner to refer the patient for appropriate co-management (e.g., to their primary
 40 care physician) or if immediate emergency care is warranted, to contact 911 as appropriate.
 41 See the *Managing Medical Emergencies in a Health Care Facility (CPG 159 – S)* clinical
 42 practice guideline for information.

1 **REFERENCES**

2 American Medical Association. (current year). *Current Procedural Terminology (CPT)*
 3 *Current year* (rev. ed.). Chicago: AMA

4
 5 Centers for Disease Prevention and Control. Overweight and Obesity. Retrieved on January
 6 2, 2025 from <https://www.cdc.gov/obesity/index.html>

7
 8 Centers for Disease Prevention and Control. Childhood Overweight and Obesity. Retrieved
 9 on January 2, 2025 from <https://www.cdc.gov/obesity/childhood/index.html>

10
 11 Hawk C, Schneider MJ, Vallone S, Hewitt EG. Best Practices for Chiropractic Care of
 12 Children: A Consensus Update. *J Manipulative Physiol Ther.* 2016 Mar-
 13 Apr;39(3):158-68. doi: 10.1016/j.jmpt.2016.02.015. Epub 2016 Mar 31. PMID:
 14 27040034

15
 16 Hawk, C., Schneider, M. J., Haas, M., Katz, P., Dougherty, P., Gleberzon, B., Killinger, L.
 17 Z., & Weeks, J. (2017). Best Practices for Chiropractic Care for Older Adults: A
 18 Systematic Review and Consensus Update. *Journal of manipulative and physiological*
 19 *therapeutics*, 40(4), 217–229. <https://doi.org/10.1016/j.jmpt.2017.02.001>

20
 21 Hill, J.O.; Wyatt, H.R.; Peters, J.C. (2013) The Importance of Energy Balance. *European*
 22 *Endocrinology*, 9(2): 111-115. Doi: 10.17925/EE.2013.09.02.111

23
 24 Joint Commission International. (2020). Joint Commission International Accreditation
 25 Standards for Hospitals (7th ed.): Joint Commission Resources

26
 27 Matarese, L, and Pories, W. Diets: Adult Weight Loss Diets: Metabolic effects and
 28 outcomes. *Nutr Clin Pract* 2014; 29: 759-767

29
 30 Trust for America’s Health (TFAH.org). (2021). The State of Obesity: Better Policies for
 31 a Healthier America. Retrieved January 2, 2025 from [https://www.tfah.org/report-](https://www.tfah.org/report-details/state-of-obesity-2021/)
 32 [details/state-of-obesity-2021/](https://www.tfah.org/report-details/state-of-obesity-2021/)

33
 34 US Preventive Services Task Force. Interventions for High Body Mass Index in Children
 35 and Adolescents: US Preventive Services Task Force Recommendation
 36 Statement. *JAMA.* 2024;332(3):226–232. doi:10.1001/jama.2024.11146 U.S

37
 38 U.S. Preventive Services Task Force (USPSTF). (2018). Weight Loss to Prevent Obesity-
 39 Related Morbidity and Mortality in Adults: Behavioral Interventions. Retrieved
 40 January 2, 2025, from
 41 [https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/obesity-in-](https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/obesity-in-adults-interventions)
 42 [adults-interventions](https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/obesity-in-adults-interventions)