

1 **Clinical Practice Guideline: Vestibular Rehabilitation**

2

3 **Date of Implementation: April 17, 2014**

4

5 **Product: Specialty**

6

7

8 **Table of Contents**

9 GUIDELINES **Error! Bookmark not defined.**

10 DESCRIPTION/BACKGROUND 1

11 EVIDENCE REVIEW 3

12 Vestibular Rehabilitation (VR) 3

13 Benign Paroxysmal Positional Vertigo (BPPV) 9

14 Concussion 14

15 Persistent Postural-Perceptual Dizziness (PPPD) 17

16 PRACTITIONER SCOPE AND TRAINING 18

17 *References* **Error! Bookmark not defined.**

18

19 **GUIDELINES**

20 American Specialty Health – Specialty (ASH) considers the use of vestibular rehabilitation,
 21 consisting of vestibular rehabilitation exercises, for the treatment of non-specific unilateral
 22 and bilateral peripheral vestibular dysfunction as medically necessary.

23

24 ASH considers the use of the Dix-Hallpike test for the diagnosis of benign paroxysmal
 25 positional vertigo (BPPV) as medically necessary. Additionally, the use of the Epley
 26 maneuver and the Semont (liberatory) maneuver for the treatment of BPPV are medically
 27 necessary for the treatment of BPPV.

28

29 ASH considers manual therapy mobilization or manipulation as not medically necessary
 30 for the treatment of isolated cervicogenic dizziness. The literature is insufficient to
 31 conclude that it is either clinically effective or ineffective in the treatment of this condition.
 32 Additional clinical trials are required to determine the effectiveness of manual therapy
 33 mobilization or manipulation for the treatment of cervicogenic dizziness for individual
 34 patients in order to determine its benefit: risk profile.

35

36 **DESCRIPTION/BACKGROUND**

37 Dizziness is a common patient complaint resulting in an estimated 7 million doctor visits
 38 per year (Hillier and McDonnell, 2011). Vertigo is a related symptom that occurs when
 39 subjects perceive movement despite being still. In a 2009 review, Neuhauser and Lempert

1 summarized the epidemiology of vertigo (Neuhauser and Lempert, 2009). Per the review
2 findings, community-based surveys indicated that as many as 20-30% of the population
3 reports complaints of dizziness or vertigo. A more detailed neurologic screening indicated
4 that the lifetime prevalence of vertigo is 7.4%, the one-year prevalence is 4.9%, and the
5 annual incidence is 1.4% in adults ages 18-79. Additional epidemiology findings showed
6 that the incidence of vertigo is 2.7 times more common in females than males and
7 prevalence increases steadily with age.

8
9 According to Hillier and McDonnell (2011), the most common source of dizziness and
10 vertigo is the vestibular system which accounts for 25% of cases. Various conditions can
11 cause vestibular pathology including surgical procedures in this region, head or neck
12 trauma, Meniere’s disease, vestibular neuritis or labyrinthitis, perilymphatic fistula,
13 acoustic neuroma, and benign paroxysmal positional vertigo (BPPV). Differential
14 diagnosis can be difficult, so many studies group patients under the category unilateral
15 peripheral vestibular dysfunction (UPVD) or hypofunction. Central nervous system
16 pathologies may also cause vestibular dysfunction, but these are less common and are often
17 excluded from studies of vestibular rehabilitation.

18
19 Patients with UPVD will report dizziness with associated visual or gaze disturbance,
20 disequilibrium, and balance abnormalities. Oscillopsia may be reported which is a visual
21 disturbance characterized by blurring or movement of the surroundings during gaze. Gaze
22 disturbances may be mediated through interruption of the vestibular-ocular reflex (VOR),
23 which functions to coordinate eye and head movements to allow for steady gaze as the
24 body moves through space. Various tests and measures have been used to measure baseline
25 status and change over time. There was considerable variation in the applied outcomes
26 measures within the studies under investigation, with authors reporting results on various
27 scales ranging from one item dichotomous (symptom resolution/not), ordinal, or visual
28 analog measures to the Vertigo Symptom Scale (14 items; 0-60 scoring). Gait disturbances
29 may be measured with gait speed or the Dynamic Gait Index (8 tasks, 0-24 scale), while
30 the Dizziness Handicap Inventory (DHI) measures participation restrictions. More
31 objective physiological measures such as electronystagmography tests for VOR were not
32 considered because they have not been correlated with function (Hillier and McDonnell,
33 2011).

34
35 Vestibular rehabilitation (VR) is frequently recommended to manage the signs and
36 symptoms of UPVD. VR typically consists of various components, each targeted to a
37 specific aspect of the pathology, including:

- 38 • Habituation exercises, which utilize repeated symptom producing motions to
39 decrease the sensitivity to stimuli via neural plasticity. These may also be termed
40 compensatory or neuroplastic strategies (Hillier and McDonnell, 2011).

- 1 • Adaptation exercises, where patients fix their gaze on a distant point while turning
- 2 their head in various directions. These are designed to train the VOR and reduce
- 3 retinal “slip” (Herdman, 2013).
- 4 • Substitution exercises, designed to sharpen other sensory organs to assist the
- 5 vestibular system in balance.
- 6 • Education on strategies to avoid provocative motions and promote safe activity
- 7 despite vestibular hypofunction.

8
 9 Medications for UPVD such as anti-nausea drugs or vestibular suppressants may be used
 10 to reduce symptoms but are seldom long-term solutions. Surgery may be used for extreme
 11 cases, including procedures such as labyrinthectomy or vestibular nerve resection. They
 12 may also be useful for specific pathologies such as an acoustic neuroma or peri-lymphatic
 13 fistula (Hillier and McDonnell, 2011). Other conservative interventions for vertigo and
 14 dizziness include canalith repositioning maneuvers, specifically for BPPV, and manual
 15 therapy, advocated for cervicogenic dizziness.

16
 17 Benign paroxysmal positional vertigo (BPPV) is characterized by short bouts of vertigo or
 18 dizziness, often with nausea, brought on by changes in position (e.g., bending down) or
 19 rapid head movements, particularly neck extension. Symptoms may resolve spontaneously
 20 and may also recur after a period of time without symptoms. BPPV may be associated with
 21 a variety of causes such as head trauma (including concussion), vestibular neuritis, and ear
 22 infection. Most cases are idiopathic. The female to male ratio is 2:1 for idiopathic; other
 23 causes are more evenly distributed. It is more common among ages 50-70. A positive Dix-
 24 Hallpike test is diagnostic for BPPV. This maneuver involves taking a patient through rapid
 25 changes in position that produce nystagmus, dizziness, and nausea.

26
 27 Persistent postural-perceptual dizziness (PPPD) is a disorder of functional dizziness that in
 28 the International Classification of Diseases in its 11th revision (ICD-11) supersedes phobic
 29 postural vertigo and chronic subjective dizziness. PPPD manifests with one or more
 30 symptoms of dizziness, unsteadiness, or non-spinning vertigo that are present on most days
 31 for three months or more and are exacerbated by upright posture, active or passive
 32 movement, and exposure to moving or complex visual stimuli. PPPD may be precipitated
 33 by conditions that disrupt balance or cause vertigo, unsteadiness, or dizziness, including
 34 peripheral or central vestibular disorders, other medical illnesses, or psychological distress.
 35 PPPD may be present alone or co-exist with other conditions.

36 **EVIDENCE REVIEW**

37 **Vestibular Rehabilitation (VR)**

38
 39
 40 Hillier and McDonnell (2011) provided a comprehensive systematic review of vestibular
 41 rehabilitation. Their comprehensive literature review included community dwelling
 42 subjects with a physician’s diagnosis of UPVD and symptoms of vertigo, dizziness, a

1 balance disorder, and/or visual or gaze disturbances. Subjects with Meniere’s disease could
2 be included if they were in later, non-fluctuating stages. There was no age limitation
3 although the majority of patients in the studies were age 65 and over. Studies which utilized
4 exercise and movement-based therapies were included, while studies that focused on
5 specific repositioning maneuvers were excluded. Comparison groups received placebo,
6 sham, usual care, no treatment, specific alternative treatment such as medication or surgery,
7 or another type of vestibular rehabilitation. Relevant outcomes included symptoms,
8 functional measures including balance, or an alternative vestibular rehabilitation approach.

9
10 The authors’ exhaustive search included articles published through July 2010. A total of
11 27 studies were included, with 10 additional articles added since the previous update was
12 published in 2007. Sample size for these articles ranged from 14-360 subjects with an
13 average of 64. Four of the studies took place in the acute hospital setting, while the rest of
14 the studies were performed in outpatient clinics. Most studies utilized a combination of
15 therapy approaches (habituation, adaptation, substitution, balance training, and education);
16 only a few studies isolated a particular therapeutic approach. Controls were most often
17 usual care or a sham exercise approach. There was a great deal of heterogeneity in outcome
18 measures, as there appears to be no generally accepted standardized measure of vestibular
19 symptoms. In their assessment of risk of bias, the authors noted generally poor reporting
20 of randomization and allocation procedures with generally low risk of bias in 4 other
21 categories.

22
23 There were 13 studies that compared VR to a sham or usual care control. Most studies
24 favored VR, but the variety of outcome measures made it difficult to formulate an overall
25 summary. When subjective improvement of dizziness was dichotomized (4 studies), VR
26 was favored with an odds ratio of 2.67 (95% CI: 1.85-3.86). Hillier and McDonald reported
27 combined standardized mean difference (SMD) of -0.67 on the Vertigo Symptom Scale (3
28 studies), -0.80 on the DHI (4 studies), and -0.92 on the Dynamic Gait Index (DGI) (3
29 studies). Various other secondary outcomes also generally supported the use of VR versus
30 a control intervention. VR was compared to alternative treatment in 6 trials. There were 2
31 studies involving subjects with a diagnosis of benign paroxysmal positional vertigo
32 (BPPV). VR was much less effective in “curing” BPPV induced dizziness than physical
33 maneuvers (odds ratio=0.13; 95% CI: 0.03-0.51); however, VR plus physical maneuvers
34 was more effective than physical maneuvers alone for the DGI (SMD=-0.87), while there
35 were non-significant findings for dizziness symptoms. One weak study, which was not
36 included in the meta-analysis due to inadequate data, compared home VR exercise to
37 betahistine medication (a vestibular suppressant) and found VR superior for relief of
38 dizziness symptoms and quality of life. Other studies comparing VR to electrical
39 stimulation or physical maneuvers had either non-significant or mixed findings.

40
41 There were 5 studies that compared one type of VR to another. In general, there were no
42 significant differences between VR approaches. There were significant differences on the

1 Vertigo Symptom Scale for a home VR program plus simulator activities versus home VR
2 alone, and for a formal program for balance and fall prevention versus a home program.
3 Hillier and McDonald noted there were generally low drop-out rates in the studies reviewed
4 and some studies showed gains lasting to 12 months (moderate evidence). The optimal
5 dosage is unclear from the literature, but they noted that “even a minimalist approach” can
6 be effective. No adverse effects were reported in any of the studies included in their review.

7
8 A systematic review published by Ricci et al. (2010) focused on the effectiveness of VR in
9 studies published in the previous 10 years that included subjects > 40 years old. They
10 located 4 studies with subjects >40 years of age, and 5 with subjects > 60 years of age.
11 Most studies included subjects with general diagnoses such as vestibular hypofunction with
12 subject complaints of dizziness, vertigo, or imbalance. They utilized the PEDro criteria for
13 scoring study quality. Nine studies were included, with 4 of 9 rated as “good” quality
14 ($\geq 6/11$ on PEDro scale). Most interventions were based on a Cawthorne and Cooksey
15 approach originally developed in the 1940s. Control subjects generally received no
16 exercise or placebo exercise; in one study, control subjects received Tai Chi. These authors
17 reported results that generally favored VR on various outcomes when compared to no
18 treatment or placebo (6 studies) but generally no significant differences when compared to
19 an alternative treatment. There were no reports of adverse reactions to VR.

20
21 In an informal review of the literature, Herdman located two additional small randomized
22 controlled trials and one crossover study that supported the effectiveness of VR for patients
23 with dizziness complaints (Herdman, 2013). McDonnell and Hillier (2015) completed an
24 update of a Cochrane review first published in 2007 and previously updated in 2011 to
25 assess the effectiveness of vestibular rehabilitation in the adult, community-dwelling
26 population of people with symptomatic unilateral peripheral vestibular dysfunction. Thirty-
27 nine studies involving 2,441 participants with unilateral peripheral vestibular disorders
28 were included in the review. Authors concluded that there was moderate to strong evidence
29 that vestibular rehabilitation is a safe, effective management for unilateral peripheral
30 vestibular dysfunction, based on several high-quality randomized controlled trials. There
31 was moderate evidence that vestibular rehabilitation resolves symptoms and improves
32 functioning in the medium term. However, there is evidence that for the specific diagnostic
33 group of BPPV, physical (repositioning) maneuvers are more effective in the short term
34 than exercise-based vestibular rehabilitation; although a combination of the two is effective
35 for longer-term functional recovery. There was insufficient evidence to discriminate
36 between differing forms of vestibular rehabilitation. Hall et al. (2016) authored an
37 evidence-based clinical practice guideline on vestibular rehabilitation for peripheral
38 vestibular hypofunction. A systematic review of the literature was performed in 5 databases
39 published after 1985 and 5 additional sources for relevant publications were searched.
40 Article types included meta-analyses, systematic reviews, randomized controlled trials,
41 cohort studies, case control series, and case series for human subjects, published in English.
42 A total of 135 articles were identified as relevant to this clinical practice guideline. Based

1 on strong evidence and a preponderance of benefit over harm, clinicians should offer
2 vestibular rehabilitation to persons with unilateral and bilateral vestibular hypofunction
3 with impairments and functional limitations related to the vestibular deficit. Based on
4 strong evidence and a preponderance of harm over benefit, clinicians should not include
5 voluntary saccadic or smooth-pursuit eye movements in isolation (i.e., without head
6 movement) as specific exercises for gaze stability. Based on moderate evidence, clinicians
7 may offer specific exercise techniques to target identified impairments or functional
8 limitations. Based on moderate evidence and in consideration of patient preference,
9 clinicians may provide supervised vestibular rehabilitation. As a general guide, persons
10 without significant comorbidities that affect mobility and with acute or subacute unilateral
11 vestibular hypofunction may need once a week supervised sessions for 2 to 3 weeks;
12 persons with chronic unilateral vestibular hypofunction may need once-a-week sessions
13 for 4 to 6 weeks; and persons with bilateral vestibular hypofunction may need once-a-week
14 sessions for 8 to 12 weeks. In addition to supervised sessions, patients are to be provided a
15 daily home exercise program.

16
17 Arnold et al. (2017) compared the effectiveness of vestibular rehabilitation interventions
18 (adaptation, substitution, and habituation) in people with unilateral peripheral vestibular
19 hypofunction, exclusionary of benign paroxysmal positional vertigo and Meniere's disease.
20 Seven papers were selected for inclusion. Results suggest that vestibular therapy for
21 unilateral peripheral vestibular hypofunction is effective. When considering all 7 studies,
22 it was difficult to determine the superiority of one intervention over another in treating
23 unilateral peripheral vestibular hypofunction except when patient outcomes are captured
24 by the dynamic gait index or dizziness handicap inventory. Maslovara et al. (2019)
25 compared the impact of VR in patients with chronic unilateral vestibular hypofunction
26 (UVH) and bilateral vestibular hypofunction (BVH). Authors concluded that well-planned
27 and individually adjusted system of vestibular exercises leads to a significant decrease in
28 clinical symptoms and improvement of functioning and confidence in activities in both the
29 chronic UVH and the BVH patient.

30
31 Tramontano et al. (2021) critically assessed the effectiveness of VR administered either
32 alone or in combination with other neurorehabilitation strategies in patients with neurologic
33 disorders. All clinical studies carried out on adult patients with a diagnosis of neurologic
34 disorders who performed VR provided alone or in combination with other therapies were
35 included. Twelve studies were included in the review. All the included studies, with 1
36 exception, report that improvements provided by customized VR in subject affected by a
37 central nervous system diseases are greater than traditional rehabilitation programs alone.
38 Authors concluded that because of the lack of high-quality studies and heterogeneity of
39 treatments protocols, clinical practice recommendations on the efficacy of VR cannot be
40 made. Results show that VR programs are safe and could easily be implemented with
41 standard neurorehabilitation protocols in patients affected by neurologic disorders. Hence,

1 more high-quality randomized controlled trials of VR in patients with neurologic disorders
2 are needed.

3
4 Hall et al. (2022) authored a revision of the 2016 guidelines published by the American
5 Physical Therapy Association and the Academy of Neurologic Physical Therapy and
6 involved a systematic review of the literature published since 2015 through June 2020
7 across 6 databases. Article types included meta-analyses, systematic reviews, randomized
8 controlled trials, cohort studies, case-control series, and case series for human subjects,
9 published in English. Sixty-seven articles were identified as relevant to this clinical practice
10 guideline and critically appraised for level of evidence. The purpose of this revised clinical
11 practice guideline is to improve quality of care and outcomes for individuals with acute,
12 subacute, and chronic unilateral and bilateral vestibular hypofunction by providing
13 evidence-based recommendations regarding appropriate exercises. The following are
14 reported:

- 15 • Based on strong evidence, clinicians should offer vestibular rehabilitation to adults
16 with unilateral and bilateral vestibular hypofunction who present with impairments,
17 activity limitations, and participation restrictions related to the vestibular deficit.
- 18 • Based on strong evidence and a preponderance of harm over benefit, clinicians
19 should not include voluntary saccadic or smooth-pursuit eye movements in
20 isolation (i.e., without head movement) to promote gaze stability.
- 21 • Based on moderate to strong evidence, clinicians may offer specific exercise
22 techniques to target identified activity limitations and participation restrictions,
23 including virtual reality or augmented sensory feedback.
- 24 • Based on strong evidence and in consideration of patient preference, clinicians
25 should offer supervised vestibular rehabilitation.
- 26 • Based on moderate to weak evidence, clinicians may prescribe weekly clinic visits
27 plus a home exercise program of gaze stabilization exercises consisting of a
28 minimum of: (a) 3 times per day for a total of at least 12 minutes daily for
29 individuals with acute/subacute unilateral vestibular hypofunction; (b) 3 to 5 times
30 per day for a total of at least 20 minutes daily for 4 to 6 weeks for individuals with
31 chronic unilateral vestibular hypofunction; (c) 3 to 5 times per day for a total of 20
32 to 40 minutes daily for approximately 5 to 7 weeks for individuals with bilateral
33 vestibular hypofunction.
- 34 • Based on moderate evidence, clinicians may prescribe static and dynamic balance
35 exercises for a minimum of 20 minutes daily for at least 4 to 6 weeks for individuals
36 with chronic unilateral vestibular hypofunction and based on expert opinion, for a
37 minimum of 6 to 9 weeks for individuals with bilateral vestibular hypofunction.
- 38 • Based on moderate evidence, clinicians may use achievement of primary goals,
39 resolution of symptoms, normalized balance and vestibular function, or plateau in
40 progress as reasons for stopping therapy.

- Based on moderate to strong evidence, clinicians may evaluate factors, including time from onset of symptoms, comorbidities, cognitive function, and use of medication that could modify rehabilitation outcomes.

In summary, recent evidence supports the original recommendations from the 2016 guidelines. There is strong evidence that vestibular physical therapy provides a clear and substantial benefit to individuals with unilateral and bilateral vestibular hypofunction. Limitations of this guideline includes that the focus of the guideline was on peripheral vestibular hypofunction; thus, the recommendations of the guideline may not apply to individuals with central vestibular disorders. One criterion for study inclusion was that vestibular hypofunction was determined based on objective vestibular function tests. This guideline may not apply to individuals who report symptoms of dizziness, imbalance, and/or oscillopsia without a diagnosis of vestibular hypofunction. These recommendations are intended as a guide to optimize rehabilitation outcomes for individuals undergoing vestibular physical therapy.

Rezaeian et al. (2023) aimed to investigate the effect of vestibular rehabilitation (VR) versus control/other interventions on the quality of life in patients with Meniere's disease (MD) in a systematic review and meta-analysis. Overall, 3 studies with a total of 465 patients were included in the meta-analysis. Authors concluded that VR can improve the quality of life in patients with MD immediately after treatment. Since all the included studies had a high risk of bias and none had long-term follow-ups, further high-quality research is required to determine the short-, intermediate-, and long-term effects of VR compared to control/other interventions.

Meng et al. (2023) aimed to evaluate the effects of vestibular rehabilitation therapy (VRT) in addition to usual rehabilitation compared with usual rehabilitation on improving balance and gait for patients after stroke in a systematic review. Fifteen randomized controlled trials with 769 participants were included. VRT was effective in improving balance for patients after stroke, particularly for patients after stroke that occurred within 6 months with moderate certainty of evidence. Subgroup analysis showed that VRT provided as gaze stability exercises combined with swivel chair training and head movements could significantly improve balance. Four-week VRT had better effect on balance improvement than the less than 4-week VRT. The pooled mean difference of values of Timed Up-and-Go test showed that VRT could significantly improve gait function for patients after stroke, particularly for patients after stroke that occurred within 6 months with moderate certainty of evidence. Authors concluded that there is moderate certainty of evidence supporting the positive effect of VRT in improving balance and gait of patients after stroke.

Kamo et al. (2023) investigated the effect of early vestibular rehabilitation on physical function and dizziness in patients with acute vestibular disorders. The inclusion criteria in terms of the study participants were patients 20 years and older with an acute unilateral

1 peripheral vestibular disorder. Early vestibular rehabilitation was defined as rehabilitation
2 within 14 days of vestibular disorder onset or surgery. Main outcome measures were gait,
3 balance (eyes open, eyes closed), activities of daily living, dizziness, and vestibular
4 function. Twelve trials involving 542 participants were included. Early vestibular
5 rehabilitation improved the Dizziness Handicap Inventory by -7.18, and dizziness by -1.47
6 compared with no intervention or placebo. Authors concluded that this study demonstrated
7 that early vestibular rehabilitation improved the Dizziness Handicap Inventory, balance
8 (eyes closed), and subjective dizziness in a patient with acute vestibular disorders. This
9 result indicates that early vestibular rehabilitation can promote vestibular compensation.

10
11 Huang et al. (2024) evaluated the efficacy of vestibular rehabilitation in vestibular neuritis
12 in a systematic review and meta-analysis. This study included 12 randomized controlled
13 trials involving 536 patients with vestibular neuritis. Vestibular rehabilitation was
14 comparable with steroids in dizziness handicap inventory score at the first, sixth, and 12th
15 months; caloric lateralization at the third, sixth, and 12th months; and abnormal numbers
16 of vestibular-evoked myogenic potentials at the first, sixth, and 12th months. Patients
17 receiving a combination of rehabilitation and steroid exhibited significant improvement in
18 dizziness handicap inventory score at the first, third, and 12th months; caloric lateralization
19 at the first and third months; and numbers of vestibular-evoked myogenic potentials at the
20 first and third months than did those receiving steroids alone. Authors concluded that
21 vestibular rehabilitation is recommended for patients with vestibular neuritis. A
22 combination of vestibular rehabilitation and steroids is more effective than steroids alone
23 in the treatment of patients with vestibular neuritis.

24 **Benign Paroxysmal Positional Vertigo (BPPV)**

25
26
27 Helminski et al. (2010) explains the two mechanisms that have been proposed to explain
28 BPPV. In normal vestibular function, calcite particles (otoconia) are attached to the sensory
29 membrane in the semicircular canals. They serve as weights which make hair-like sensors
30 in the canals sensitive to acceleration movements in their fluid-filled environment. In the
31 mechanism known as canalithiasis, BPPV is hypothesized to result when otoconia break
32 loose and float free in the endolymph, where their movement continues even after the head
33 has stopped moving, thereby causing vestibular symptoms. The other mechanism is termed
34 cupulolithiasis, where the calcite particles become embedded in the cupula, the gelatinous
35 membrane of the canal, causing abnormal weighting in the sensory organ. BPPV may be
36 divided into three types based on canal involvement: posterior, horizontal, and anterior
37 semicircular canal BPPV. The posterior semicircular canal is most often involved in this
38 mechanism. BPPV cases involving the horizontal semicircular canal are reportedly less
39 common and can be more difficult to treat. Anterior Canal BPPV is considered rare and
40 deemed more likely to be self-treated, or resolved, due to gravity. (Gupta et al., 2019).

1 The first treatments for BPPV were habituation exercise, first reported in the 1950s. Later,
2 a physical maneuver was advocated by Epley that uses gravity and 4 position changes
3 designed to move any loose particles through the posterior semicircular canal into the
4 vestibule, where they will not produce symptoms. The success of the Epley maneuver as a
5 treatment for BPPV has led to favoring of the canalithiasis mechanism for BPPV (Hilton
6 and Pinder, 2004). Following, a second physical maneuver known as the Semont or
7 liberatory maneuver was developed to address cases of cupulolithiasis and canalithiasis,
8 involving a rapid 180-degree movement from side-lying on the involved side to side-lying
9 on the uninvolved side to loosen any particles lodged in the cupula. Collectively, these are
10 known as particle repositioning maneuvers. There are home versions of each maneuver and
11 postural/neck range of motion restrictions may be advised for 24-48 hours following
12 treatment.

13
14 In an update of a 2004 Cochrane review, Hilton and Pinder (2014) included studies
15 published through May 2010 that included patients with a positive Dix-Hallpike test, limited
16 to randomized controlled trials studying the Epley versus no treatment, placebo, or an
17 alternative mode of treatment. Key outcomes for inclusion were incidence and severity of
18 vertigo, patient ratings of improvement, and/or a negative Dix-Hallpike test. Their search
19 yielded 22 randomized controlled trials, however 17 were excluded due to high risk of bias
20 (mostly randomization procedure and lack of blinded allocation). For the 5 studies with
21 low risk of bias, the sample sizes were generally small (36-81 total) and included patients
22 with symptoms less than 2 weeks. Four of the studies used a sham control while one study
23 used a no treatment control group. Four weeks was the longest follow-up. Meta-analysis
24 revealed a pooled odds ratio of 4.42 (2.62, 7.44) in favor of the Epley maneuver for
25 complete resolution of symptoms, and a pooled odds ratio of 6.4 (3.6, 11.3) for a negative
26 Dix-Hallpike outcome. They found widely varying estimates of natural recovery, from 15-
27 84%. Only one study reported adverse effects – inability to tolerate the Epley maneuver
28 due to vomiting or pre-existing neck pain – but the adverse event rate was not reported.

29
30 A companion systematic review by Hunt et al. (2012) focused on adjuncts to the Epley
31 maneuver including limiting cervical movements and maintaining upright posture for 24-
32 48 hours following maneuver, perhaps with a soft collar, and mastoid vibration, using a
33 mechanical device attached to a headband. They included randomized controlled trials
34 involving patients with confirmed BPPV. They located 11 randomized controlled trials that
35 met their inclusion criteria; nine investigated postural restrictions and 2 studies involved
36 oscillation to mastoid during the Epley maneuver. Sample sizes varied from 38-106, and
37 follow-up was typically 1 week, although a few studies had a longer follow-up period.
38 They found that the addition of postural restrictions yielded significantly better results for
39 conversion of the Dix-Hallpike test with a risk ratio = 1.13 (1.05, 1.22). Adverse events
40 were tracked in 5 studies; neck stiffness was more common in the intervention group (27%
41 versus none in one study); development of horizontal BPPV, transient nausea and
42 disequilibrium also occurred rarely but not more common in the experimental versus the

1 control group. The Epley maneuver plus mastoid oscillation was compared to the Epley
 2 maneuver alone in 2 studies; there were no significant differences in conversion of Dix-
 3 Hallpike or in the intensity of symptoms.

4
 5 Helminski et al. (2010) performed a systematic review to determine the effectiveness of
 6 particle positioning maneuvers, including the Epley or the Semont (liberatory) method, to
 7 treat BPPV. Their search included randomized controlled trials (RCTs) or quasi
 8 randomized controlled trials published through 2009. Randomized controlled trials
 9 provided strong evidence that the canalith repositioning procedure (CRP) resolves
 10 posterior canal (PC) benign paroxysmal positional nystagmus (BPPN); whereas quasi-
 11 RCTs suggested that the CRP or the liberatory maneuver performed by a health care
 12 practitioner, or with proper instruction at home by the patient, resolves PC BPPN. Their
 13 preferred measure of success was the conversion from a positive to a negative Dix-Hallpike
 14 test since vertigo symptoms are dependent on activity levels. Their search yielded 10
 15 studies total:

- 16 • There were 2 true RCTs and 2 quasi-randomized controlled trials that all found the
 17 Epley maneuver superior (67-95% success) to a sham intervention (10-38%
 18 success). In the 2 true RCTs the odds of resolution of the Dix-Hallpike test were
 19 22-37 times higher for the treatment group.
- 20 • There were 2 quasi-RCTs that compared the Semont (liberatory) maneuver to no
 21 treatment that favored the experimental group with 80-85% success versus 35-38%
 22 in controls, with an odds ratio of 7-10.
- 23 • There were 2 quasi-RCTs that compared the Semont (liberatory) maneuver to the
 24 Epley maneuver but found no difference overall.
- 25 • There were 3 quasi-RCTs that looked at the effectiveness of self-treatment using a
 26 particle repositioning maneuver with or without an in-clinic treatment. They found
 27 90-95% success overall, with 58% for liberatory and 24% for VR exercise only.
 28 The odds ratio was 3.5 for Epley + self-administered versus Epley alone. Self-
 29 treatment using the Epley maneuver was more effective than using the self-
 30 liberatory maneuver (odds ratio = 12.5).

31
 32 Clinical practice guidelines by the American Academy of Otolaryngology—Head and
 33 Neck Surgery Foundation (Bhattacharyya et al., 2008) strongly supported use of the Dix-
 34 Hallpike test for diagnosis and canalith repositioning maneuvers for treatment of posterior
 35 canal BPPV. Wegner et al. (2014) evaluated the effectiveness the Epley maneuver
 36 compared to vestibular rehabilitation for BPPV. Only 5 of 373 relevant articles satisfied
 37 the eligibility criteria. Results demonstrated that the Epley maneuver is more effective in
 38 treating BPPV than vestibular rehabilitation at 1-week follow-up. There is inconsistent
 39 evidence for the effectiveness of the Epley maneuver compared with vestibular
 40 rehabilitation at 1-month follow-up. An update of the Cochrane Review (Hilton and Pinder,
 41 2014) concluded that there is evidence that the Epley maneuver is a safe, effective
 42 treatment for posterior canal BPPV, based on the results of 11, mostly small, randomized

1 controlled trials with relatively short follow-up. There is a high recurrence rate of BPPV
2 after treatment (36%). Outcomes for Epley maneuver treatment are comparable to
3 treatment with Semont and Gans maneuvers, but superior to Brandt-Daroff exercises.
4 Adverse effects were infrequently reported. There were no serious adverse effects of
5 treatment. Rates of nausea during the repositioning maneuver varied from 16.7% to 32%.
6 Some patients were unable to tolerate the maneuver because of cervical spine problems.
7 Oh et al. (2017) compared the efficacy between repetition of Epley maneuver and switch
8 to alternate Semont maneuver in treating posterior canal benign paroxysmal positional
9 vertigo (PC-BPPV) that does not respond to the initial Epley maneuver. 144 (28.5%)
10 patients, who did not respond to the therapy, were randomized to the repetition of Epley
11 maneuver ($n = 70$) or switch to Semont maneuver ($n = 74$). The therapeutic efficacy was
12 determined within 1 hour by a blinded examiner after the trial of each second maneuver.
13 The efficacy did not differ between the repetition of Epley maneuver and switch to Semont
14 maneuver groups. However, the patients with a long duration ($p < 0.001$, linear regression)
15 and latency ($p = 0.01$) of the positional nystagmus during Dix-Hallpike maneuver showed
16 a higher rate of the initial and second treatment failures. Either Epley or Semont maneuver
17 may be applied as a second treatment to the patients with PC-BPPV refractory to the initial
18 Epley maneuver. This study provides Class I evidence that repeated Epley and switch to
19 Semont maneuver shows a similar efficacy in treating PC-BPPV that does not respond to
20 the initial Epley maneuver.

21
22 Bhattacharyya et al. (2017) updated the clinical practice guideline. Changes from the prior
23 guideline include a consumer advocate added to the update group; new evidence from 2
24 clinical practice guidelines, 20 systematic reviews, and 27 randomized controlled trials;
25 enhanced emphasis on patient education and shared decision making; a new algorithm to
26 clarify action statement relationships; and new and expanded recommendations for the
27 diagnosis and management of BPPV. The primary purposes of this guideline were to
28 improve the quality of care and outcomes for BPPV by improving the accurate and efficient
29 diagnosis of BPPV, reducing the inappropriate use of vestibular suppressant medications,
30 decreasing the inappropriate use of ancillary testing such as radiographic imaging, and
31 increasing the use of appropriate therapeutic repositioning maneuvers. The primary
32 outcome considered in this guideline was the resolution of the symptoms associated with
33 BPPV. Secondary outcomes considered included an increased rate of accurate diagnoses
34 of BPPV, a more efficient return to regular activities and work, decreased use of
35 inappropriate medications and unnecessary diagnostic tests, reduction in recurrence of
36 BPPV, and reduction in adverse events associated with undiagnosed or untreated BPPV.
37 The update group made strong recommendations that clinicians should (1) diagnose
38 posterior semicircular canal BPPV when vertigo associated with torsional, upbeat
39 nystagmus is provoked by the Dix-Hallpike maneuver, and (2) treat, or refer to a clinician
40 who can treat, patients with posterior canal BPPV with a canalith repositioning procedure.
41 The update group made a strong recommendation against postprocedural postural
42 restrictions after canalith repositioning procedure for posterior canal BPPV. The update

1 group made recommendations that the clinician should (1) perform, or refer to a clinician
 2 who can perform, a supine roll test to assess for lateral semicircular canal BPPV if the
 3 patient has a history compatible with BPPV and the Dix-Hallpike test exhibits horizontal
 4 or no nystagmus; (2) differentiate, or refer to a clinician who can differentiate, BPPV from
 5 other causes of imbalance, dizziness, and vertigo; (3) assess patients with BPPV for factors
 6 that modify management, including impaired mobility or balance, central nervous system
 7 disorders, a lack of home support, and/or increased risk for falling; (4) reassess patients
 8 within 1 month after an initial period of observation or treatment to document resolution
 9 or persistence of symptoms; (5) evaluate, or refer to a clinician who can evaluate, patients
 10 with persistent symptoms for unresolved BPPV and/or underlying peripheral vestibular or
 11 central nervous system disorders; and (6) educate patients regarding the impact of BPPV
 12 on their safety, the potential for disease recurrence, and the importance of follow-up. The
 13 update group made recommendations against (1) radiographic imaging for a patient who
 14 meets diagnostic criteria for BPPV in the absence of additional signs and/or symptoms
 15 inconsistent with BPPV that warrant imaging, (2) vestibular testing for a patient who meets
 16 diagnostic criteria for BPPV in the absence of additional vestibular signs and/or symptoms
 17 inconsistent with BPPV that warrant testing, and (3) routinely treating BPPV with
 18 vestibular suppressant medications such as antihistamines and/or benzodiazepines. The
 19 guideline update group provided the options that clinicians may offer (1) observation with
 20 follow-up as initial management for patients with BPPV and (2) vestibular rehabilitation,
 21 either self-administered or with a clinician, in the treatment of BPPV.

22
 23 Rodrigues et al. (2019) evaluated the additional effects of vestibular rehabilitation
 24 exercises as a therapeutic resource in the treatment of BPPV, to improve symptoms and
 25 reduce recurrence. Thirty-two individuals, both men and women, over 18 years of age with
 26 BPPV were randomly assigned to two groups: the control group ($n=15$) performing only
 27 the maneuver technique as treatment and the experimental group ($n=17$) performing the
 28 maneuvers and vestibular rehabilitation exercises. Results demonstrated that the
 29 experimental group had a lower level of dizziness in the posttreatment period ($p<0.05$) and
 30 a lower incidence of recurrences ($p=0.038$) than the control group. Authors concluded that
 31 vestibular exercises performed after repositioning treatments for BPPV increased the
 32 overall efficacy of treatment by improving symptoms with a lower rate of recurrence.
 33 Power et al. (2020) outlined the incidence of BPPV in specialized vestibular physiotherapy
 34 clinics and discusses the various nuances encountered during assessment and treatment of
 35 BPPV. Interventions included canalith repositioning maneuvers (CRP) for posterior canal
 36 (PC) or horizontal canal (HC) BPPV depending on the canal and variant of BPPV.
 37 Outcome measures included negative Dix-Hallpike or supine roll test examination. Results
 38 indicated that in 91% of cases, PC BPPV was effectively treated in 2 maneuvers or less.
 39 Similarly, 88% of HC BPPV presentations were effectively managed with 2 treatments.
 40 Bilateral PC, multiple canal or canal conversions required a greater number of treatments.
 41 There was no noticeable difference in treatment outcomes for patients who had nystagmus
 42 and symptoms during the Epley maneuver (EM) versus those who did not have nystagmus

1 and symptoms throughout the EM. Nineteen percent of patients experienced post treatment
 2 down-beating nystagmus and vertigo or “otolithic crisis” after the first or even the second
 3 consecutive EM. Authors concluded that based on the data collected, repeated testing and
 4 treatment of BPPV within the same session is promoted as a safe and effective approach to
 5 the management of BPPV with a low risk of canal conversion. Secondly, vertigo and
 6 nystagmus throughout the EM is not indicative of treatment success. Thirdly, clinicians
 7 must remain vigilant and mindful of the possibility of post treatment otolithic crisis
 8 following the treatment of BPPV. This is to ensure patient safety and to prevent possible
 9 injurious falls.

10
 11 Li et al. (2022) compared the efficacy of different treatments for posterior semicircular
 12 canal benign paroxysmal positional vertigo (PC-BPPV) by using direct and indirect
 13 evidence from existing randomized data. A total of 41 parallel, randomized controlled
 14 studies were included. The Epley with vestibular rehabilitation (EVR), Epley, Semont and
 15 Hybrid maneuvers were effective in eliminating nystagmus during a Dix-Hallpike test at 1
 16 week of follow-up, among which EVR showed the best efficacy. However, at 1 month of
 17 follow-up, only the Semont and Epley maneuvers were effective in eliminating nystagmus
 18 during a Dix-Hallpike test. In the pairwise subgroup meta-analysis, for patients younger
 19 than 55 years of age, the efficacy of the Epley maneuver was comparable to that of the
 20 Semont maneuver; for patients with a longer duration before treatment, the effect of the
 21 Epley maneuver was equivalent to that of a sham maneuver. Authors concluded that among
 22 the 12 types of PC-BPPV treatments, the Epley, Semont, EVR, and Hybrid maneuvers
 23 were effective in eliminating nystagmus during a Dix-Hallpike test for PC-BPPV at 1 week
 24 of follow-up, whereas only the Epley and Semont maneuvers were effective at 1 month of
 25 follow-up.

26
 27 Pauwels et al. (2023) assessed the influence of BPPV and treatment effects of particle
 28 repositioning maneuvers (PRM) on gait, falls, and fear of falling. Twenty of the 25 included
 29 studies were suitable for meta-analysis. BPPV increases the odds of falls and negatively
 30 impacts spatiotemporal parameters of gait. PRM improves falls, fear of falling, and gait
 31 during level walking. Additional rehabilitation might be necessary to improve gait while
 32 walking with head movements or tandem walking.

33 34 **Concussion**

35 Murray et al. (2017) systematically evaluated the evidence supporting the efficacy,
 36 prescription, and progression patterns of VRT in patients with concussion. Following a
 37 double review of abstract and full-text articles, 10 studies met the inclusion criteria:
 38 randomized controlled trial ($n=2$), uncontrolled studies ($n=3$) and case studies ($n=5$). 4
 39 studies evaluated VRT as a single intervention. 6 studies incorporated VRT in multimodal
 40 interventions (including manual therapy, strength training, occupational tasks, counselling,
 41 or medication). 9 studies reported improvement in outcomes but level I evidence from only
 42 1 study was found that demonstrated increased rates of medical clearance for return to sport

1 within 8 weeks, when VRT (combined with cervical therapy) was compared with usual
2 care. Heterogeneity in study type and outcomes precluded meta-analysis. Habituation and
3 adaptation exercises were employed in 8 studies and balance exercises in 9 studies. Authors
4 concluded that the current evidence for optimal prescription and efficacy of VRT in
5 patients with mTBI/concussion is limited. Available evidence, although weak, shows
6 promise in this population. Further high-level studies evaluating the effects of VRT in
7 patients with mTBI/concussion with vestibular and/or balance dysfunction are required.

8
9 Park et al. (2018) investigated whether VRT, rather than continued prescription of rest
10 (cognitive and physical), reduce recovery time and persistent symptoms of dizziness,
11 unsteadiness, and imbalance in adolescents (12-18 y) who suffer post-concussive
12 syndrome following a sports-related concussion. Authors noted that VRT was an effective
13 intervention for this population. Adolescents presenting with this cluster of symptoms may
14 also demonstrate verbal and visual memory loss linked to changes in the vestibular system
15 post-concussion. Authors concluded that moderate evidence supports that adolescents who
16 suffer from persistent symptoms of dizziness, unsteadiness, and imbalance following sport
17 concussion should be evaluated more specifically and earlier for vestibular dysfunction and
18 can benefit from participation in individualized VRT. Early evaluation and treatment may
19 result in a reduction of time lost from sport as well as a return to their pre-morbid condition.
20 For these adolescents, VRT may be more beneficial than continued physical and cognitive
21 rest when an adolescent's symptoms last longer than 30 days. Storey et al. (2018) sought
22 to determine whether active vestibular rehabilitation is associated with an improvement in
23 visuovestibular signs and symptoms in children with concussion. One hundred nine
24 children were included in the study with a mean age of 11.8 (3.4) years. Among this group,
25 59 (54%) were male and 48 (44%) had a sports-related concussion. Authors concluded that
26 vestibular rehabilitation in children with concussion is associated with improvement in
27 symptoms as well as visuovestibular performance. This active intervention may benefit
28 children with persistent symptoms after concussion. Future prospective studies are needed
29 to determine the efficacy and optimal postinjury timing of vestibular rehabilitation.
30 Schlemmer and Nicholson (2022) synthesized the best available evidence regarding the
31 effectiveness of VRT as a treatment option for adults with mTBIs. Five studies were
32 included in the systematic review: 1 randomized controlled trial, 2 retrospective chart
33 reviews, 1 pre-/post-intervention study, and 1 case series. Four of the 5 studies found VRT
34 to be effective at reducing postconcussion symptoms after head injury. Self-reported
35 measures were included in all studies; performance-based measures were included in four
36 out of five studies. None of the studies reported adverse effects of intervention. Authors
37 concluded that results suggest VRT is an effective treatment option for patients with
38 persistent/lingering symptoms after concussion/mTBI, as demonstrated by self-reported
39 and performance-based outcome measures.

40
41 Reid et al. (2022) investigated the effect of physical interventions (subthreshold aerobic
42 exercise, cervical, vestibular and/or oculomotor therapies) on days to recovery and

1 symptom scores in the management of concussion. Twelve trials met the inclusion criteria:
2 7 on subthreshold aerobic exercise, 1 on vestibular therapy, 1 on cervical therapy and 3 on
3 individually tailored multimodal interventions. The trials were of fair to excellent quality
4 on the PEDro scale. Eight trials were included in the quantitative analysis. Subthreshold
5 aerobic exercise had a significant small to moderate effect in improving symptom scores
6 but not in reducing days to symptom recovery in both acutely concussed individuals and
7 those with persistent symptoms. There was limited evidence for stand-alone cervical,
8 vestibular and oculomotor therapies. Concussed individuals with persistent symptoms (>2
9 weeks) were approximately 3 times more likely to have returned to sport by 8 weeks if they
10 received individually tailored, presentation-specific multimodal interventions (cervical,
11 vestibular, and oculo-motor therapy). In addition, the multimodal interventions had a
12 moderate effect in improving symptom scores when compared with control. Authors
13 concluded that subthreshold aerobic exercise appears to lower symptom scores but not time
14 to recovery in concussed individuals. Individually tailored multimodal interventions have
15 a worthwhile effect in providing faster return to sport and clinical improvement,
16 specifically in those with persistent symptoms.

17
18 LeMarshall et al. (2023) aimed to identify, synthesize, and assess the quality of studies
19 reporting on the effectiveness of virtual reality for the rehabilitation of vestibular and
20 balance impairments post-concussion in a scoping review. Additionally, this review aimed
21 to summarize the volume of scientific literature and identify the knowledge gaps in current
22 research pertaining to this topic. Data was charted from studies and outcomes were
23 categorized into one of three categories: balance, gait, or functional outcome measures.
24 Three randomized controlled trials, 3 quasi-experimental studies, 3 case studies, and 1
25 retrospective cohort study were ultimately included, using a thorough eligibility criteria.
26 All studies were inclusive of different virtual reality interventions. The ten studies had a
27 10-year range and identified 19 different outcome measures. Authors concluded that
28 findings from this review suggests that virtual reality is an effective tool for the
29 rehabilitation of vestibular and balance impairments post-concussion. Current literature
30 shows sufficient but low level of evidence, and more research is necessary to develop a
31 quantitative standard and to better understand appropriate dosage of virtual reality
32 intervention.

33
34 Schneider et al. (2023) evaluated interventions to facilitate recovery in children,
35 adolescents, and adults with a sport-related concussion (SRC). Thirteen studies met
36 inclusion (10 RCTs, 1 quasi-experimental and 2 cohort studies; 1 high-quality study, 7
37 acceptable and 5 at high risk of bias). Interventions, comparisons, timing, and outcomes
38 varied, precluding meta-analysis. For adolescents and adults with dizziness, neck pain
39 and/or headaches >10 days following concussion, individualized cervicovestibular
40 rehabilitation may decrease time to return to sport compared with rest followed by gradual
41 exertion and when compared with a subtherapeutic intervention. For adolescents with
42 vestibular symptoms/impairments, vestibular rehabilitation may decrease time to medical

1 clearance (vestibular rehab group 50.2 days compared with control 58.4). For adolescents
 2 with persisting symptoms >30 days, active rehabilitation and collaborative care may
 3 decrease symptoms. Authors concluded that cervicovestibular rehabilitation is
 4 recommended for adolescents and adults with dizziness, neck pain and/or headaches for
 5 >10 days. Vestibular rehabilitation (for adolescents with dizziness/vestibular impairments
 6 >5 days) and active rehabilitation and/or collaborative care (for adolescents with persisting
 7 symptoms >30 days) may be of benefit.

9 **Persistent Postural-Perceptual Dizziness (PPPD)**

10 Dieterich and Staab (2017) reviewed nomenclature, clinical features, possible
 11 pathomechanisms, and comorbidities of functional dizziness. The prevalence of functional
 12 dizziness as a primary cause of vestibular symptoms amounts to 10% in neuro-otology
 13 centers. Rates of psychiatric comorbidity in patients with structural vestibular syndromes
 14 are much higher with nearly 50% and with highest rates in patients with vestibular
 15 migraine, vestibular paroxysmia, and Ménière’s disease. Correct and early diagnosis of
 16 functional dizziness, as primary cause or secondary disorder after a structural vestibular
 17 syndrome, is very important to prevent further chronification and enable adequate
 18 treatment. Treatment plans that include patient education, vestibular rehabilitation,
 19 cognitive and behavioral therapies, and medications substantially reduce morbidity and
 20 offer the potential for sustained remission when applied systematically.

21
 22 Popkirov et al. (2018) reviewed different treatment strategies for this common functional
 23 neurological disorder. Authors noted that an emerging understanding of the underlying
 24 pathophysiology that considers vestibular, postural, cognitive, and emotional aspects can
 25 enable patients to profit from vestibular rehabilitation, as well as cognitive-behavioral
 26 therapy (CBT). Most importantly, approaches from CBT should inform and augment
 27 physiotherapeutic techniques, and vestibular exercises or relaxation techniques can be
 28 integrated into CBT programs. They conclude that, in PPPD and related disorders,
 29 vestibular rehabilitation combined with CBT can help patients escape a cycle of
 30 maladaptive balance control, recalibrate vestibular systems, and regain independence in
 31 everyday life. Staab (2020) reports in an article on PPPD that the diagnosis is made by
 32 identifying key symptoms in patients’ histories and conducting physical examinations and
 33 diagnostic testing of sufficient detail to establish PPPD as opposed to other illnesses.
 34 Ongoing research is providing insights into the pathophysiological mechanisms underlying
 35 PPPD and support for multimodality treatment plans incorporating specially adapted
 36 vestibular rehabilitation, serotonergic medications, and cognitive-behavior therapy. Cha
 37 (2021) authored an article that covered distinct causes of chronic dizziness including
 38 persistent postural perceptual dizziness, mal de débarquement syndrome, motion sickness
 39 and visually induced motion sickness, bilateral vestibulopathy, and persistent dizziness
 40 after mild concussion. Cha states that to date, none of these disorders has a cure but are
 41 considered chronic syndromes with fluctuations that are both innate and driven by
 42 environmental stressors. As such, the mainstay of therapy for chronic disorders of dizziness

1 involves managing factors that exacerbate symptoms and adding vestibular rehabilitation
2 or cognitive-behavioral therapy alone or in combination, as appropriate. These therapies
3 are supplemented by serotonergic antidepressants that modulate sensory gating and reduce
4 anxiety. Besides expectation management, ruling out concurrent disorders and recognizing
5 behavioral and lifestyle factors that affect symptom severity are critical issues in reducing
6 morbidity for each disorder.

7
8 Rogers et al. (2023) summarized dizziness and its evaluation and management in an article.
9 The physical examination may include orthostatic blood pressure measurement, a full
10 cardiac and neurologic examination, assessment for nystagmus, the Dix-Hallpike
11 maneuver (for patients with triggered dizziness), and the HINTS (head-impulse,
12 nystagmus, test of skew) examination when indicated. The treatment for dizziness is
13 dependent on the etiology of the symptoms. Canalith repositioning procedures (e.g., Epley
14 maneuver) are the most helpful in treating benign paroxysmal positional vertigo. Vestibular
15 rehabilitation is helpful in treating many peripheral and central etiologies. Other etiologies
16 of dizziness require specific treatment to address the cause. Pharmacologic intervention is
17 limited because it often affects the ability of the central nervous system to compensate for
18 dizziness.

19
20 Webster et al. (2023) assessed the benefits and harms of non-pharmacological interventions
21 for PPPD. Primary outcomes were: 1) improvement in vestibular symptoms 2) change in
22 vestibular and 3) serious adverse events. Secondary outcomes were: 4) disease-specific
23 health-related quality of life, 5) generic health-related quality of life and 6) other adverse
24 effects. Outcomes were reported at three time points: 3 to < 6 months, 6 to ≤ 12 months
25 and > 12 months. Of the few studies identified, only one followed up with participants for
26 at least three months, therefore most were not eligible for inclusion in this review. Authors
27 concluded that further work is necessary to determine whether any non-pharmacological
28 interventions may be effective for the treatment of PPPD and to assess whether they are
29 associated with any potential harms. As this is a chronic disease, future trials should follow
30 up participants for a sufficient period of time to assess whether there is a persisting impact
31 on the severity of the disease, rather than only observing short-term effects.

32 33 **PRACTITIONER SCOPE AND TRAINING**

34 Practitioners should practice only in the areas in which they are competent based on their
35 education, training, and experience. Levels of education, experience, and proficiency may
36 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
37 to determine where they have the knowledge and skills necessary to perform such services.

38
39 It is best practice for the practitioner to appropriately render services to a patient only if
40 they are trained, equally skilled, and adequately competent to deliver a service compared
41 to others trained to perform the same procedure. If the service would be most competently

1 delivered by another health care practitioner who has more skill and expert training, it
2 would be best practice to refer the patient to the more expert practitioner.

3
4 Best practice can be defined as a clinical, scientific, or professional technique, method, or
5 process that is typically evidence-based and consensus driven and is recognized by a
6 majority of professionals in a particular field as more effective at delivering a particular
7 outcome than any other practice (Joint Commission International Accreditation Standards
8 for Hospitals, 2020).

9
10 Depending on the practitioner’s scope of practice, training, and experience, a member’s
11 condition and/or symptoms during examination or the course of treatment may indicate the
12 need for referral to another practitioner or even emergency care. In such cases it is prudent
13 for the practitioner to refer the member for appropriate co-management (e.g., to their
14 primary care physician) or if immediate emergency care is warranted, to contact 911 as
15 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice
16 guideline for information.

17 **References**

18
19 Arnold SA, Stewart AM, Moor HM, Karl RC, Reneker JC. The Effectiveness of Vestibular
20 Rehabilitation Interventions in Treating Unilateral Peripheral Vestibular Disorders: A
21 Systematic Review. *Physiother Res Int.* 2017 Jul;22(3)

22
23 Bhattacharyya N, Baugh RF, Orvidas L, et al. Clinical practice guideline: Benign
24 paroxysmal positional vertigo. *Otolaryngol Head Neck Surg.* 2008;139(5 Suppl
25 4):S47-81

26
27 Bhattacharyya N, Gubbels SP, Schwartz SR, et al. Clinical Practice Guideline: Benign
28 Paroxysmal Positional Vertigo (Update). *Otolaryngol Head Neck Surg.*
29 2017;156(3_suppl):S1-S47

30
31 Brodovsky JR, Vnenchak MJ. Vestibular rehabilitation for unilateral peripheral vestibular
32 dysfunction. *Phys Ther.* 2013;93(3):293-298.

33
34 Cha YH. Chronic Dizziness. *Continuum (Minneap Minn).* 2021;27(2):420-446

35
36 Dieterich M, Staab JP. Functional dizziness: from phobic postural vertigo and chronic
37 subjective dizziness to persistent postural-perceptual dizziness. *Curr Opin Neurol.*
38 2017;30(1):107-113

39
40 Gupta AK, Sharma KG, Sharma P. Effect of Epley, Semont Maneuvers and Brandt-Daroff
41 Exercise on Quality of Life in Patients with Posterior Semicircular Canal Benign

- 1 Paroxysmal Positional Vertigo (PSCBPPV). *Indian J Otolaryngol Head Neck Surg.*
 2 2019;71(1):99-103
 3
- 4 Hall CD, Herdman SJ, Whitney SL, Cass SP, Clendaniel RA, Fife TD, Furman JM,
 5 Getchius TS, Goebel JA, Shepard NT, Woodhouse SN. Vestibular Rehabilitation for
 6 Peripheral Vestibular Hypofunction: An Evidence-Based Clinical Practice Guideline:
 7 From The American Physical Therapy Association Neurology Section. *J Neurol Phys*
 8 *Ther.* 2016 Apr;40(2):124-55
 9
- 10 Hall CD, Herdman SJ, Whitney SL, et al. Vestibular Rehabilitation for Peripheral
 11 Vestibular Hypofunction: An Updated Clinical Practice Guideline From the Academy
 12 of Neurologic Physical Therapy of the American Physical Therapy Association. *J*
 13 *Neurol Phys Ther.* 2022;46(2):118-177
 14
- 15 Helminski JO, Zee DS, Janssen I, Hain TC. Effectiveness of particle repositioning
 16 maneuvers in the treatment of benign paroxysmal positional vertigo: A systematic
 17 review. *Phys Ther.* 2010;90(5):663-678
 18
- 19 Herdman SJ. Vestibular rehabilitation. *Curr Opin Neurol.* 2013;26(1):96-101
 20
- 21 Hillier SL, McDonnell M. Vestibular rehabilitation for unilateral peripheral vestibular
 22 dysfunction. *Cochrane Database Syst Rev.* 2011;(2):CD005397. Doi(2):CD005397
 23
- 24 Hilton M, Pinder D. The Epley (canalith repositioning) maneuver for benign paroxysmal
 25 positional vertigo. *Cochrane Database Syst Rev.* 2004;(2)(2):CD003162
 26
- 27 Hilton MP, Pinder DK. The Epley (canalith repositioning) maneuver for benign
 28 paroxysmal positional vertigo. *Cochrane Database Syst Rev.* 2014 Dec
 29 8;(12):CD003162
 30
- 31 Huang HH, Chen CC, Lee HH, Chen HC, Lee TY, Tam KW, Kuan YC. Efficacy of
 32 Vestibular Rehabilitation in Vestibular Neuritis: A Systematic Review and Meta-
 33 analysis. *Am J Phys Med Rehabil.* 2024 Jan 1;103(1):38-46
 34
- 35 Hunt WT, Zimmermann EF, Hilton MP. Modifications of the Epley (canalith
 36 repositioning) maneuver for posterior canal benign paroxysmal positional vertigo
 37 (BPPV). *Cochrane Database Syst Rev.* 2012;4:CD008675
 38
- 39 Kamo T, Ogihara H, Azami M, Momosaki R, Fushiki H. Effects of Early Vestibular
 40 Rehabilitation in Patients With Acute Vestibular Disorder: A Systematic Review and
 41 Meta-Analysis. *Otol Neurotol.* 2023 Oct 1;44(9):e641-e647

- 1 LeMarshall SJ, Stevens LM, Ragg NP, Barnes L, Foster J, Canetti EFD. Virtual reality-
2 based interventions for the rehabilitation of vestibular and balance impairments post-
3 concussion: a scoping review. *J Neuroeng Rehabil.* 2023 Mar 3;20(1):31
4
- 5 Li D, Cheng D, Yang W, et al. Current Therapies in Patients With Posterior Semicircular
6 Canal BPPV, a Systematic Review and Network Meta-analysis. *Otol Neurotol.*
7 2022;43(4):421-428
8
- 9 Lystad RP, Bell G, Bonnevie-Svendsen M, Carter CV. Manual therapy with and without
10 vestibular rehabilitation for cervicogenic dizziness: A systematic review. *Chiropr Man
11 Therap.* 2011;19(1):21-709X-19-21
12
- 13 Maslovara S, Butkovic-Soldo S, Peric M, Pajic Matic I, Sestak A. Effect of vestibular
14 rehabilitation on recovery rate and functioning improvement in patients with chronic
15 unilateral vestibular hypofunction and bilateral vestibular hypofunction.
16 *NeuroRehabilitation.* 2019;44(1):95-102
17
- 18 McDonnell MN, Hillier SL. Vestibular rehabilitation for unilateral peripheral vestibular
19 dysfunction. *Cochrane Database Syst Rev.* 2015 Jan 13;1:CD005397
20
- 21 Meng L, Liang Q, Yuan J, Li S, Ge Y, Yang J, Tsang RCC, Wei Q. Vestibular rehabilitation
22 therapy on balance and gait in patients after stroke: a systematic review and meta-
23 analysis. *BMC Med.* 2023 Aug 25;21(1):322
24
- 25 Murray DA, Meldrum D, Lennon O. Can vestibular rehabilitation exercises help patients
26 with concussion? A systematic review of efficacy, prescription and progression
27 patterns. *Br J Sports Med.* 2017;51(5):442-451
28
- 29 Neuhauser HK, Lempert T. Vertigo: Epidemiologic aspects. *Semin Neurol.*
30 2009;29(5):473-481
31
- 32 Oh SY, Kim JS, Choi KD, Park JY, Jeong SH, Lee SH, Lee HS, Yang TH, Kim HJ. Switch
33 to Semont maneuver is no better than repetition of Epley maneuver in treating
34 refractory BPPV. *J Neurol.* 2017 Sep;264(9):1892-1898
35
- 36 Park K, Ksiazek T, Olson B. Effectiveness of Vestibular Rehabilitation Therapy for
37 Treatment of Concussed Adolescents With Persistent Symptoms of Dizziness and
38 Imbalance. *J Sport Rehabil.* 2018 Sep 1;27(5):485-490
39
- 40 Pauwels S, Casters L, Lemkens N, Lemmens W, Meijer K, Meyns P, van de Berg R,
41 Spildooren J. Gait and Falls in Benign Paroxysmal Positional Vertigo: A Systematic
42 Review and Meta-analysis. *J Neurol Phys Ther.* 2023 Jul 1;47(3):127-138

- 1 Popkirov S, Stone J, Holle-Lee D. Treatment of Persistent Postural-Perceptual Dizziness
2 (PPPD) and Related Disorders. *Curr Treat Options Neurol*. 2018 Oct 13;20(12):50
3
- 4 Power L, Murray K, Szmulewicz DJ. Characteristics of assessment and treatment in Benign
5 Paroxysmal Positional Vertigo (BPPV). *J Vestib Res*. 2020;30(1):55-62
6
- 7 Reid SA, Farbenblum J, McLeod S. Do physical interventions improve outcomes following
8 concussion: a systematic review and meta-analysis?. *Br J Sports Med*. 2022;56(5):292-
9 298. doi:10.1136/bjsports-2020-103470
10
- 11 Rezaeian A, Abtahi H, Moradi M, Farajzadegan Z. The effect of vestibular rehabilitation
12 in Meniere's disease: a systematic review and meta-analysis of clinical trials. *Eur Arch*
13 *Otorhinolaryngol*. 2023 Sep;280(9):3967-3975
14
- 15 Ricci NA, Aratani MC, Dona F, Macedo C, Caovilla HH, Gananca FF. A systematic review
16 about the effects of the vestibular rehabilitation in middle-age and older adults. *Rev*
17 *Bras Fisioter*. 2010;14(5):361-371
18
- 19 Rodrigues DL, Ledesma ALL, Pires de Oliveira CA, Bahmad F Jr. Effect of Vestibular
20 Exercises Associated With Repositioning Maneuvers in Patients With Benign
21 Paroxysmal Positional Vertigo: A Randomized Controlled Clinical Trial. *Otol*
22 *Neurotol*. 2019 Sep;40(8):e824-e829
23
- 24 Rogers TS, Noel MA, Garcia B. Dizziness: Evaluation and Management. *Am Fam*
25 *Physician*. 2023 May;107(5):514-523. PMID: 37192077
26
- 27 Schneider KJ, Critchley ML, Anderson V, Davis GA, Debert CT, Feddermann-Demont N,
28 Gagnon I, Guskiewicz KM, Hayden KA, Herring S, Johnstone C, Makdissi M, Master
29 CL, Moser RS, Patricios JS, Register-Mihalik JK, Ronksley PE, Silverberg ND, Yeates
30 KO. Targeted interventions and their effect on recovery in children, adolescents and
31 adults who have sustained a sport-related concussion: a systematic review. *Br J Sports*
32 *Med*. 2023 Jun;57(12):771-779
33
- 34 Schlemmer E, Nicholson N. Vestibular Rehabilitation Effectiveness for Adults With Mild
35 Traumatic Brain Injury/Concussion: A Mini-Systematic Review. *Am J Audiol*.
36 2022;31(1):228-242
37
- 38 Staab JP. Persistent Postural-Perceptual Dizziness. *Semin Neurol*. 2020;40(1):130-137
39
- 40 Storey EP, Wiebe DJ, D'Alonzo BA, Nixon-Cave K, Jackson-Coty J, Goodman AM,
41 Grady MF, Master CL. Vestibular Rehabilitation Is Associated With Visuovestibular
42 Improvement in Pediatric Concussion. *J Neurol Phys Ther*. 2018 Jul;42(3):134-141

- 1 Tramontano M, Russo V, Spitoni GF, et al. Efficacy of Vestibular Rehabilitation in Patients
2 With Neurologic Disorders: A Systematic Review. Arch Phys Med Rehabil.
3 2021;102(7):1379-1389
4
- 5 Webster KE, Kamo T, Smith L, Harrington-Benton NA, Judd O, Kaski D, Maarsingh OR,
6 MacKeith S, Ray J, Van Vugt VA, Burton MJ. Non-pharmacological interventions for
7 persistent postural-perceptual dizziness (PPPD). Cochrane Database Syst Rev. 2023
8 Mar 13;3(3):CD015333
9
- 10 Wegner I, Niesten ME, van Werkhoven CH, Grolman W. Rapid Systematic Review of the
11 Epley Maneuver versus Vestibular Rehabilitation for Benign Paroxysmal Positional
12 Vertigo. Otolaryngol Head Neck Surg. 2014 May 20;151(2):201-207