

1 **Clinical Practice Guideline: Temporomandibular Joint Disorder**

2  
3 **Date of Implementation: April 17, 2014**

4  
5 **Product: Specialty**

6  
7  
8 **GUIDELINES**

9 American Specialty Health – Specialty (ASH) considers conservative approaches (physical  
10 therapy and manual therapy such as active and passive exercises, postural training,  
11 mobilizations/manipulative therapy, and myofascial therapy) to be medically necessary  
12 when used in combination with one another.

13  
14 ASH considers electro-physiotherapy modalities (transcutaneous electrical nerve  
15 stimulation [TENS] and/or pulsed radio-frequency energy [PRFE]) and laser/light therapy  
16 (LLLT) for the treatment of temporomandibular joint disorder as not medically necessary.

17  
18 Clinical evidence does not support the use or the effectiveness of these modalities for  
19 treatment of Temporomandibular Disorder (TMD). Additionally, pulsed radio-frequency  
20 energy (PRFE) has a negative benefit-risk profile and presents a health and safety risk when  
21 used due to its physical properties. There is some evidence that LLLT may improve  
22 function, but further research is needed to confirm results. There is also some evidence that  
23 dry needling improves pain and function, but again, further research is needed to confirm  
24 results. For additional information, please see the *Electric Stimulation for Pain, Swelling*  
25 *and Function in a Clinic Setting (CPG 272-S)*, *Laser Therapy (LT) (CPG 30 – S)*, and  
26 *Passive Physiotherapy Modalities (CPG 121-S)* clinical practice guidelines.

27  
28 ASH considers the use of acupuncture for the symptomatic relief of temporomandibular  
29 joint pain as medically necessary. Please see the *Acupuncture Services Medical*  
30 *Policy/Guideline (CPG 264-S)* clinical practice guideline for additional information.

31  
32 **DESCRIPTION/BACKGROUND**

33 The temporomandibular joint (TMJ), a synovial hinge joint, is located where the mandible  
34 joins the temporal bone via an intra-articular disc. This complex synovial system is further  
35 comprised of articulating ligaments and masticatory muscles. The TMJ is functioning  
36 properly when the right-sided and left-sided joints are synchronized during movement. It  
37 is also one of the most frequently utilized joints within the body, used up to 2,000 times a  
38 day for such functions as mastication, swallowing, respiration and speech.

39  
40 TMD can be classified collectively as temporomandibular joint and muscle disorders that  
41 cause pain and dysfunction in the jaw joint and the muscles that control jaw movement or  
42 surrounding soft tissues. Normal mandible movement requires coordination between these

1 structures to maximize function and minimize the damage to surrounding structures. A  
2 rather unique feature of temporomandibular joint articulation is that it has two joints. The  
3 articular disc between the condyle and the temporal bone serves to separate the structures  
4 into two separate joint cavities. In the inferior joint between the head of the mandibular  
5 condyle and the articular disc, the movement is almost completely of a rotary or hinge type;  
6 whereas in the superior joint between the temporal bone and the articular disc the  
7 movement is gliding, or translational.

8  
9 The numerous epidemiologic studies on the occurrence of TMD in the general population  
10 indicate a number of consistent findings. Firstly, signs of TMD appear in about 60–70% of  
11 the general population, yet only about one in four people with signs are actually aware of  
12 or report any symptoms. The frequency of severe disorders that are accompanied by  
13 headache and facial pain, and that are characterized by urgent need of treatment is 1–2%  
14 in children, approximately 5% in adolescents and 5–12% in adults. Among those who seek  
15 treatment for TMD, by far the great majority are females, outnumbering males by at least  
16 four to one – although it is suspected that TMD affects both males and females in almost  
17 equal numbers in the general population.

18  
19 Similar to other musculoskeletal disorders, pain during function, or while at rest is the  
20 primary reason that therapy is sought. Less commonly, patients seek TMD therapy for  
21 temporomandibular joint catching and locking, masticatory stiffness, limited mandibular  
22 range of motion, temporomandibular joint dislocation, and occlusal changes.  
23 Temporomandibular joint noises (e.g., clicking, popping) are common among the general  
24 population, however, this is generally not a concern for patients and practitioners; hence  
25 are not commonly treated.

## 26 27 **DIAGNOSTIC CONSIDERATIONS IN THE TREATMENT OF TMD**

28 This disorder can be classified into three groups or types: disc displacement/internal  
29 derangement, muscle disorders, and arthroses. The most common disorder of the  
30 temporomandibular joint is disc displacement. In essence, this is when the articular disc,  
31 attached anteriorly to the superior head of the lateral pterygoid muscle and posteriorly to  
32 the retrodiscal tissue, becomes displaced from between the condyle and the fossa, so that  
33 the mandible and temporal bone contact is made on something other than the articular disc.  
34 This, as explained above, is usually very painful, because disc displacement can lead to the  
35 development of secondary inflammatory changes and progressive degradation of the  
36 articular cartilage (Maizlin et al., 2010). Muscle disorders include pain dysfunction  
37 syndrome, myofascial pain, and myofascial pain syndrome. This type presents with pain in  
38 the jaw, temple, face, preauricular area or inside the ear, at rest or during function. Lastly,  
39 arthroses TMD are comprised of arthritis (including osteo-, rheumatoid, traumatic, and  
40 psoriatic arthritis), arthrosis and ankyloses (such as ankylosing spondylitis affecting the  
41 temporomandibular joint). Arthroses present with joint sounds, limited mandibular  
42 movements and pain, and can be secondary to muscular or disc displacement TMD.

1 The quality of the pain is generally an ache, pressure, and/or dull pain and may include a  
 2 background burning sensation. There may also be episodes of sharp pain, and when the  
 3 pain worsens, the primary pain quality may become a throbbing sensation. Patients with  
 4 TMD tend to report that their pain is intensified by events such as stress, clenching, and  
 5 eating, while it is relieved by relaxing, applying heat to the painful area, and taking over-  
 6 the-counter analgesics. While the patient may be experiencing the aforementioned pain, it  
 7 is useful to note that TMD can also be associated with various comorbidities such as tension  
 8 headache, whiplash, fibromyalgia, tinnitus, vertigo, hearing loss, abnormal swallowing,  
 9 hyoid bone tenderness, and otalgia.

10  
 11 Current insight into TMD indicates its etiology is multifactorial; whereas historically,  
 12 occlusion of the jaw was considered the primary cause of TMD. Therefore, establishing a  
 13 concise mode of treatment for the condition presents a challenge to the health care  
 14 practitioner. A collaborative, interdisciplinary effort between practitioners in the diagnosis  
 15 and management of TMD is thus encouraged.

16  
 17 The first line of non-surgical treatment for TMD has traditionally been physiotherapy,  
 18 pharmacotherapy, and splint therapy. However, TMD treatment trends in recent decades  
 19 have leaned toward multi-modal as well as multi-disciplinary management, in line with  
 20 that of other chronic musculoskeletal conditions. Such strategies often suggest the use of  
 21 less invasive interventions such as biofeedback, cognitive and behavioral therapies,  
 22 chiropractic, and acupuncture.

## 23 **EVIDENCE REVIEW**

24  
 25 A systematic review by Brantingham et al. (2013) identifies 5 trials for the treatment of  
 26 TMD with what it calls “Manual and Manipulate Therapy” (MMT). The range of therapies  
 27 comprising MMT include exercise, mobilization, manual distraction, massage, muscle  
 28 relaxation and intra-oral myofascial therapy (IMT). Of these 5 clinical studies, 4 are  
 29 randomized clinical trials and 1 a non-randomized trial. The review concludes that there is  
 30 limited (level B) evidence supporting the use of MMT for TMD treatment. This is based  
 31 on the finding of “2 high-quality, 2 medium-quality and 1 low quality trials.” It further  
 32 concludes that the following interventions provide benefits for TMD: “intraoral myofascial  
 33 therapy (IMT), post isometric relaxation, manual distraction, and self-mobilization in  
 34 conjunction with a variety of exercises and gentle, high-velocity (very) low-amplitude  
 35 manipulation, soft tissue MMT, or extra-oral soft tissue mobilization alone or as  
 36 multimodal care.” Finally, the review notes that in addition to these 5 trials there is a large  
 37 body of mixed high, moderate, and low level evidence from a variety of studies including  
 38 case series, case reports, single cohort pre-post studies, etc.

39  
 40 Of the 5 studies reviewed, 3 have very small ( $n < 30$ ) sample sizes and would be more  
 41 properly viewed as pilot studies. Of the 2 larger studies (Kalamir et al., 2012; Minakuchi  
 42 et al., 2001), only the Kalamir study reported positive results. Additionally, the

1 heterogeneity of treatments, patient inclusion criteria and outcome measures represented  
2 by these studies are inconsistent and further studies with improved controls are necessary  
3 to demonstrate the effectiveness of manual manipulative therapy for the treatment of TMD.  
4 Two studies (Kalamir et al., 2010; Kalamir et al., 2012) did use a common treatment of  
5 intra-oral myofascial therapy (IMT). George et al. (2007) investigated the effects of manual  
6 therapy applied to the cervical-cranial junction to determine effects on mouth-opening  
7 capacity within an asymptomatic population. A total of 101 participants were randomly  
8 assigned to either an Active Release Technique (ART) group; high-velocity, low-  
9 amplitude manipulation (HVLA) group; or control group. A blinded investigator measured  
10 mouth opening using a TheraBite range of motion scale. Participants received ART to the  
11 suboccipital or HVLA to the cervical spine at C1 or sat with an investigator for 3 minutes  
12 with no treatment. After the treatment session, mouth opening was re-measured. ART and  
13 HVLA to the cervical spine did not significantly improve mouth opening in this  
14 asymptomatic population.

15  
16 Alves et al. (2013) conducted a systematic review to identify whether mandibular  
17 manipulation technique is an effective and safe technique for the treatment of the  
18 temporomandibular joint disk displacement without reduction. Only 2 studies of medium  
19 quality fulfilled all the inclusion criteria. There is no sufficient evidence to support the  
20 effectiveness of the mandibular manipulation therapy, and therefore its use remains  
21 questionable. The analysis of the results suggested that additional high-quality randomized  
22 clinical trials are necessary and should focus on methods for data randomization and  
23 allocation, on clearly defined outcomes, on a priori calculated sample size, and on an  
24 adequate follow-up strategy. There are 2 additional randomized controlled trials (RCTs)  
25 that are not identified by the Brantingham review which are relevant. Kalamir et al. (2013)  
26 carried out an RCT ( $n=46$ ) again comparing intra-oral myofascial therapies (IMT) to  
27 education, self-care, and exercise for TMD. This study evaluated short-term differences,  
28 over a course of 6 weeks (each patient receiving 2 therapy sessions per week), in pain and  
29 mouth opening range between IMT and an exercise program. While the study concluded  
30 that IMT presented a decrease in pain and increased mouth opening range, the results were  
31 not regarded as clinically significant.

32  
33 Calixtre et al. (2015) studied manual therapy for the management of pain and limited range  
34 of motion in subjects with signs and symptoms of temporomandibular disorder. Their aim  
35 of this systematic review is to synthesize evidence regarding the isolated effect of MT in  
36 improving maximum mouth opening (MMO) and pain in subjects with signs and symptoms  
37 of TMD. Myofascial release and massage techniques applied on the masticatory muscles  
38 were more effective than control (low to moderate evidence) but as effective as toxin  
39 botulinum injections (moderate evidence). Upper cervical spine thrust manipulation or  
40 mobilization techniques were more effective than control (low to high evidence), while  
41 thoracic manipulations were not. There was moderate-to-high evidence that MT techniques  
42 protocols were effective. In conclusion, there is widely varying evidence that MT improves

1 pain, MMO and pressure pain threshold (PPT) in subjects with TMD signs and symptoms,  
2 depending on the technique. Further studies should improve their design to strengthen  
3 clinical relevance.

4  
5 Martins et al. (2016) studied the efficacy of musculoskeletal manual approaches (e.g.,  
6 mobilization, manual traction, manipulation, myofascial release, trigger point therapy,  
7 manual translations) in the treatment of temporomandibular joint disorder within a  
8 systematic review with meta-analysis. From the 308 articles identified by the search  
9 strategy, only 8 articles met the inclusion criteria. The meta-analysis showed a significant  
10 difference ( $p < 0.0001$ ) and large effect on active mouth opening and on pain during active  
11 mouth in favor of musculoskeletal manual techniques when compared to other conservative  
12 treatments for TMD. Authors concluded that musculoskeletal manual approaches are  
13 effective for treating TMD. In the short term, there is a larger effect regarding the latter  
14 when compared to other conservative treatments for TMD.

15  
16 McNeely et al. (2006) reviewed the efficacy of exercise and postural therapy interventions  
17 for the treatment of TMD. This review is notable for its clear and explicit reporting of study  
18 quality on the 5-point Jadad scale. Four studies examined the effect of exercise  
19 interventions on TMD. However, the methodological quality of these 4 studies was  
20 considered weak. Two studies examined the effect of posture training (in combination with  
21 other therapies) on myogenous TMD and reported significant improvements in pain and  
22 oral opening in favor of the addition of postural exercise training. After 1 month,  
23 Komiyama et al. (1999) found a significant increase in mouth opening in patients who  
24 received postural training compared with patients receiving only cognitive intervention or  
25 compared with the control group. Wright et al. (2000) found a statistically significant  
26 improvement in maximum pain-free opening, pain threshold, and the modified symptom  
27 severity index in patients receiving postural treatment compared with patients receiving  
28 self-management instructions alone. Carmeli et al. (2001) compared the effect of manual  
29 therapy in combination with active exercise with the effect of treatment with occlusal splint  
30 therapy on anteriorly displaced temporomandibular disks on 36 patients with arthrogenous  
31 TMD. The authors reported significant improvement in pain and oral opening in favor of  
32 the manual therapy/exercise group. Grace et al. (2002) examined the benefit of an oral  
33 exercise device compared to traditional therapies, including when the oral exercise device  
34 was used as part of a home program, on oral opening, pain, and wellness in patients with  
35 mixed TMD. Results indicated that the study groups demonstrated significant clinical  
36 improvement. However, the groups did not differ significantly from each other in degree  
37 of patient improvement. McNeely et al. (2006) further reviewed the efficacy of various  
38 electro-physiotherapy modalities in the treatment of TMD pain and dysfunction and  
39 reported on 6 studies (2 strong studies and 4 weak studies). There was considerable  
40 heterogeneity among the studies in the type of TMD, the chosen modality and comparison  
41 group, and in the frequency and duration of the treatment.

1 In the double-blind, placebo-controlled study by Al-Badawi et al. (2004), forty patients  
2 received 6 treatments of pulsed radio-frequency energy (PRFE) therapy, however PRFE  
3 was not found to be significantly better than sham PRFE for arthrogenous TMD pain.  
4 Treacy et al. (1999) reported that 20 sessions of transcutaneous electrical nerve stimulation  
5 (TENS), were not significantly better than muscular awareness relaxation therapy (MART)  
6 or sham TENS ( $n=23$  patients). Significant improvements were found, however, in oral  
7 opening and electromyographic activity for the MART group when compared with  
8 treatment with TENS and sham TENS. The Treacy study is methodologically weak due to  
9 small sample size, lack of double blinding, and inadequate data collection methods.

10  
11 A review by List and Axelsson (2010) examined the set of systematic reviews for the entire  
12 range of treatments for TMD including surgery, occlusal appliances, medication, as well  
13 as physical and manual therapies. This review found that there was great variability in  
14 quality and methodology of the reviews as well as in the primary studies, making definitive  
15 conclusions impossible. This analysis concluded that occlusal appliances, acupuncture,  
16 behavioral therapy, jaw exercises, postural training, and some pharmacological treatments  
17 were effective for TMD. There was insufficient evidence for effectiveness for electro-  
18 physiotherapy modalities.

19  
20 Moraes et al. (2013) studied therapeutic exercises for the control of temporomandibular  
21 disorders. Their aim was to conduct a literature review concerning the types of exercises  
22 available and the efficacy for the treatment of muscular TMD. The results included 7  
23 articles which reported therapeutic exercises to be effective for the treatment of muscular  
24 TMD. However, these studies were deemed limited with regards to the conclusions because  
25 the exercises were combined with other conservative treatments. Other limitations  
26 included: small samples, lack of control group and no detailed exercise description, which  
27 should have included intensity, repetition, frequency, and duration. Authors conclude that  
28 although therapeutic exercises are considered effective in the management of muscular  
29 TMD, the development of randomized clinical trials is necessary, since many existing  
30 studies are still based on the clinical experience of professionals. Another study, Kraaijenga  
31 et al. (2014), compared in a randomized controlled clinical trial (RCT) the application of  
32 the TheraBite® (TB) Jaw Motion Rehabilitation System with a standard physical therapy  
33 (PT) exercise regimen for the treatment of myogenic temporomandibular disorder (TMD).  
34 Mandibular function was assessed with the mandibular function impairment questionnaire  
35 (MFIQ). Pain was evaluated using a visual analog scale, and maximum inter-incisor  
36 (mouth) opening (MIO) was measured using the disposable TB range of motion scale.  
37 After six-week follow-up, patients using the TB device reported a significantly greater  
38 functional improvement (MFIQ score) than the patients receiving regular PT exercises  
39 ( $P=0.0050$ ). At 6 weeks, no significant differences in pain, and active or passive MIO were  
40 found between the two groups. At 3 months, patients in both treatment groups did equally  
41 well, and showed a significant improvement in all parameters assessed. This RCT showed  
42 that both treatment modalities are equally effective in relieving myogenic TMD symptoms,

1 but that the use of the TB device has the benefit of achieving a significantly greater  
2 functional improvement within the first week of treatment.

3  
4 Rashid et al. (2013) investigated the perceived effectiveness of physiotherapy for patients  
5 with TMD among consultants in oral and maxillofacial surgery (OMFS). A total of 208  
6 responded (58%) and 72% considered physiotherapy to be effective. Amongst these  
7 respondents, jaw exercises (79%), ultrasound (52%), manual therapy (48%), acupuncture  
8 (41%) and laser therapy (15%) were considered to be effective. Twenty-eight percent of  
9 respondents did not consider physiotherapy to be effective. Reasons for this included lack  
10 of knowledge or expertise of the physiotherapist (41%) and lack of awareness of the  
11 benefits of physiotherapy (28%). Despite limited evidence to support its effectiveness,  
12 approximately three-quarters of OMFS consultants in the UK regard physiotherapy to be  
13 beneficial in the management of TMD. Chen et al. (2015) evaluated the efficacy of low-  
14 level laser therapy (LLLT) in the treatment of temporomandibular disorders (TMDs).  
15 Fourteen highly qualified RCTs reporting on a total of 454 patients, which evaluated the  
16 effectiveness of LLLT for patients suffering from TMDs were retrieved. The results  
17 indicated that LLLT was not better than placebo in reducing chronic TMD pain. However,  
18 the LLLT provided significant better functional outcomes in terms of maximum active  
19 vertical opening (MAVO), maximum passive vertical opening (MPVO), protrusion  
20 excursion (PE) and right lateral excursion (RLE). Authors conclude that this study indicates  
21 that using LLLT has limited efficacy in reducing pain in patients with TMDs. However,  
22 LLLT can significantly improve the functional outcomes of patients with TMDs.

23  
24 In an article by Shaffer et al. (2014), conservative management of TMJ disorders is  
25 discussed. Authors state that physical therapy is the preferred conservative management  
26 approach for TMD. They suggest that the potentially appropriate plan of care components  
27 may include joint and soft tissue mobilization, trigger point dry needling, friction massage,  
28 therapeutic exercise, patient education, modalities, and outside referral. Management  
29 options should address both symptom reduction and oral function. Satisfactory results can  
30 often be achieved when management focuses on patient-specific clinical variables.

31 Wieckiewicz et al. (2015) presented the concepts of TMD pain clinical management based  
32 on the most current treatment plans. Results reported that the most common conservative  
33 treatments are massage therapy and individually fabricated occlusal splints. In addition to  
34 massage, other popular methods include manual therapy and taping, warming/cooling of  
35 aching joints, and light and laser therapy. Drugs are also commonly used. In the most severe  
36 cases of the temporomandibular joint degeneration, surgical restoration of the joint is  
37 sometimes applied. Authors conclude that conservative treatment including counselling,  
38 exercises, occlusal splint therapy, massage, manual therapy, and others should be  
39 considered as a first-choice therapy for TMD pain because of their low risk of side effects.  
40 In the case of severe acute pain or chronic pain resulting from serious disorders,  
41 inflammation and/or degeneration pharmacotherapy, minimally invasive and invasive  
42 procedures should be considered.

1 Gauer and Semidey (2015) reported on standard treatment for patients with TMD. They  
2 report that most patients improve with a combination of noninvasive therapies, including  
3 patient education, self-care, cognitive behavior therapy, pharmacotherapy, physical  
4 therapy, and occlusal devices. Nonsteroidal anti-inflammatory drugs and muscle relaxants  
5 are recommended initially, and benzodiazepines or antidepressants may be added for  
6 chronic cases. Referral to an oral and maxillofacial surgeon is indicated for refractory  
7 cases.

8  
9 Armijo-Olivo et al. (2016) summarized evidence of randomized controlled trials that  
10 examined the effectiveness of MT and therapeutic exercise interventions compared with  
11 other active interventions or standard care for treatment of TMD. Randomized controlled  
12 trials involving adults with TMD that compared any type of MT intervention (e.g.,  
13 mobilization, manipulation) or exercise therapy with a placebo intervention, controlled  
14 comparison intervention, or standard care were included. The main outcomes were pain,  
15 range of motion, and oral function. Forty-eight studies met the inclusion criteria and were  
16 analyzed. The overall evidence for this systematic review was considered low, with an  
17 unclear or high risk of bias. Most of the effect sizes were low to moderate, with no clear  
18 indication of superiority of exercises versus other conservative treatments for TMD.  
19 However, MT alone or in combination with exercises at the jaw or cervical level showed  
20 promising effects. Overall, there was no high-quality evidence, indicating that there is  
21 uncertainty about the effectiveness of exercise and MT for treatment of TMD.

22  
23 According to Butts et al. (2017), a review of the literature revealed limited support of  
24 strengthening exercises targeting the muscles of mastication. There was also limited  
25 evidence for manual soft tissue work targeting muscles of mastication, which may be  
26 specifically related to the limited accessibility of the pterygoid muscles to palpation. For  
27 the reduction of pain, there was little to no evidence supporting splint therapy and  
28 electrophysical modalities, including laser therapy, ultrasound, TENs, and iontophoresis.  
29 However, for the reduction of pain and disability, non-thrust mobilization and high-  
30 velocity, low amplitude thrust manipulation techniques to the TMJ and/or upper cervical  
31 articulations that directly and indirectly target the TMJ joint capsule were generally  
32 supported in the literature. Studies that used dry needling or acupuncture of the lateral  
33 pterygoid and posterior, peri-articular connective tissue also led to significant  
34 improvements in pain and disability in patients with TMD. Thus, the most effective  
35 conservative management of TMD seems to be techniques best able to impact anatomic  
36 structures directly related to the etiology of TMD, to include the joint capsule, articular  
37 disc, and muscles of mastication, specifically the superior and inferior head of the lateral  
38 pterygoid.

39  
40 Garrigós-Pedró et al. (2018) investigated the effects of adding orofacial treatment to  
41 cervical physical therapy in patients with chronic migraine and temporomandibular  
42 disorders (TMD). A total of 45 participants with chronic migraine and TMD aged 18 to 65



1 years were randomized into two groups: a cervical group (CG) and a cervical and orofacial  
 2 group (COG). Both groups continued their medication regimens for migraine treatment  
 3 and received physical therapy. The CG received physical therapy only in the cervical  
 4 region, and the COG received physical therapy in both the cervical and orofacial regions.  
 5 Both groups received six sessions of treatment that consisted of manual therapy and  
 6 therapeutic exercise in the cervical region or the cervical and orofacial regions. Scores on  
 7 the Craniofacial Pain and Disability Inventory (CF-PDI) and the Headache Impact Test  
 8 (HIT-6) were primary outcome variables, and the secondary outcome variables were scores  
 9 on the Tampa Scale for Kinesiophobia (TSK-11), pain intensity measured on a visual  
 10 analog scale (VAS), pressure pain thresholds (PPTs) in the temporal, masseter and  
 11 extratrigeminal (wrist) regions, and maximal mouth opening (MMO). Data were recorded  
 12 at baseline, posttreatment, and after 12 weeks of follow-up. There were 22 CG participants  
 13 (13.6% men and 86.4% women) and 23 COG participants (13% men and 87% women).  
 14 The ANOVA analysis revealed statistically significant differences for group  $\times$  time  
 15 interaction in CF-PDI, HIT-6 in the last follow-up, pain intensity, PPTs in the trigeminal  
 16 region, and MMO, with a medium-large magnitude of effect. No statistically significant  
 17 differences were found in the PPTs of the extratrigeminal region or in the TSK-11. Authors  
 18 concluded that both groups reported a significant improvement in CF-PDI, HIT-6, and pain  
 19 intensity. Cervical and orofacial treatment was more effective than cervical treatment alone  
 20 for increasing PPTs in the trigeminal region and producing pain-free MMO. Physical  
 21 therapy alone was not effective for increasing the PPTs in the extratrigeminal region (wrist)  
 22 or decreasing the level of TSK-11.

23  
 24 Shimada et al. (2019) authored a review focused on the effects of exercise therapy for the  
 25 management of painful TMD. The aims of this review were to summarize the effects of  
 26 exercise therapy for major symptoms of painful TMD and to establish a guideline for the  
 27 management of painful TMD, resulting in higher quality and reliability of dental treatment.  
 28 In this review, exercise modalities are clearly defined as follows: mobilization exercise,  
 29 muscle strengthening exercise (resistance training), coordination exercise and postural  
 30 exercise. Furthermore, pain intensity and range of movements were focused as outcome  
 31 parameters in this review. Authors concluded that mobilization exercise including manual  
 32 therapy, passive jaw mobilization with oral appliances and voluntary jaw exercise appeared  
 33 to be a promising option for painful TMD conditions such as myalgia and arthralgia.  
 34 Calixtre et al. (2019) sought to determine whether mobilization of the upper cervical region  
 35 and craniocervical flexor training decreased orofacial pain, increased mandibular function  
 36 and pressure pain thresholds (PPTs) of the masticatory muscles and decreased headache  
 37 impact in women with TMD when compared to no intervention. Sixty-one women with  
 38 TMD were randomized into an intervention group (IG) and a control group (CG). The IG  
 39 received upper cervical mobilizations and neck motor control and stabilization exercises  
 40 for 5 weeks. The CG received no treatment. Pain intensity showed significant time-by-  
 41 group interaction, with significant between-group differences at four and five weeks, with  
 42 large effect sizes ( $d > 0.8$ ). The decrease in orofacial pain over time was clinically relevant

1 only in the IG. Change in headache impact was significantly different between groups, and  
2 the IG showed a clinically relevant decrease after the treatment. No effects were found for  
3 PPT or mandibular function. Authors concluded that women with TMD reported a  
4 significant decrease in orofacial pain and headache impact after 5 weeks of treatment aimed  
5 at the upper cervical spine compared to a CG.

6  
7 Vier et al. (2019) systematically reviewed the effects of dry needling on orofacial pain of  
8 myofascial origin in patients with temporomandibular joint dysfunction. Seven trials were  
9 considered eligible. There was discrepancy among dry needling treatment protocols. Meta-  
10 analysis showed that dry needling is better than other interventions for pain intensity as  
11 well as than sham therapy on pressure pain threshold, but there is very low-quality evidence  
12 and a small effect size. There were no statistically significant differences in other outcomes.  
13 Authors suggested that clinicians can use dry needling for the treatment of  
14 temporomandibular joint dysfunction. However, due to the low quality of evidence and  
15 high risk of bias of some included studies, larger and higher quality studies are needed to  
16 assess the effects of dry needling on orofacial pain associated with temporomandibular  
17 joint dysfunction. Madani et al. (2020) compared the efficacy of low-level laser therapy  
18 (LLLT) versus laser acupuncture therapy (LAT) in patients with temporomandibular  
19 disorders (TMDs). In this randomized, double-blind clinical trial, 45 TMD patients were  
20 randomly divided into three groups: group 1 (LLLT), group 2 (LAT), and group 3 (placebo)  
21 underwent treatment with sham laser. There was no significant difference in mouth opening  
22 between the groups, but the amount of lateral excursive and protrusive movements was  
23 significantly greater in LLLT and LAT groups than the placebo group at some intervals.  
24 The overall pain intensity and pain degree at masticatory muscles (except temporal muscle)  
25 and TMJs were significantly lower in both experimental groups than the placebo group at  
26 most intervals after therapy. Authors concluded that both LLLT and LAT were effective  
27 in reducing pain and increasing excursive and protrusive mandibular motion in TMD  
28 patients. LAT could be suggested as a suitable alternative to LLLT, as it provided effective  
29 results while taking less chair time.

30  
31 Reynolds et al. (2020) sought to determine the immediate and short-term effects of adding  
32 cervical spine high-velocity, low-amplitude thrust (HVLAT) to behavioral education, soft  
33 tissue mobilization, and a home exercise program on pain and dysfunction for people with  
34 a primary complaint of temporomandibular disorder (TMD) with myalgia. Fifty  
35 individuals with TMD were randomly assigned to receive cervical HVLAT or sham  
36 manipulation for 4 visits over 4 weeks. Participants in both groups received other  
37 treatments, including standardized behavioral education, soft tissue mobilization, and a  
38 home exercise program. Primary outcomes included maximal mouth opening, the numeric  
39 pain-rating scale, the Jaw Functional Limitation Scale (JFLS), the Tampa Scale of  
40 Kinesiophobia for TMD (TSK-TMD), and a global rating of change (GROC). Self-report  
41 and objective measurements were taken at baseline, immediately after initial treatment, and  
42 follow-ups of 1 week and 4 weeks. Results indicated that there was no significant

1 interaction for maximal mouth opening, the numeric pain-rating scale, or secondary  
2 measures. The HVLAT group had lower fear at 4 weeks and improved jaw function earlier  
3 (1 week). The GROC favored the HVLAT group, with significant differences in successful  
4 outcomes noted immediately after baseline treatment (thrust, 6/25; sham, 0/25) and at 4  
5 weeks (thrust, 17/25; sham, 10/25). Authors concluded that both groups improved over  
6 time; however, differences between groups were small. The additive clinical effect of  
7 cervical HVLAT to standard care remains unclear for treating TMD.

8  
9 Delgado de la Serna et al. (2020) investigated the effects of adding cervico-mandibular  
10 manual therapies into an exercise and educational program on clinical outcomes in  
11 individuals with tinnitus associated with temporomandibular disorders (TMDs). Sixty-one  
12 patients with tinnitus attributed to TMD were randomized into the physiotherapy and  
13 manual therapy group or physiotherapy alone group. All patients received 6 sessions of  
14 physiotherapy treatment including cranio-cervical and temporomandibular joint (TMJ)  
15 exercises, self-massage, and patient education for a period of 1 month. Patients allocated  
16 to the manual therapy group also received cervico-mandibular manual therapies targeting  
17 the TMJ and cervical and masticatory muscles. Primary outcomes included TMD pain  
18 intensity and tinnitus severity. Patients were assessed at baseline, 1 week, 3 months, and 6  
19 months after intervention by a blinded assessor. Authors reported that this clinical trial  
20 found that application of cervico-mandibular manual therapies in combination with  
21 exercise and education resulted in better outcomes than application of exercise/education  
22 alone in individuals with tinnitus attributed to TMD.

23  
24 Fisch et al. (2020) explored if physical therapy is an effective approach to treating patients  
25 with TMJ disorders. They sought to determine the effect of conservative physical therapy  
26 interventions on pain, maximal mouth opening, and TMJ disability index for patients with  
27 TMD. Medical records from 2013-2018 were retrospectively reviewed to identify patients  
28 and obtain demographic, baseline, and short-term outcomes of maximal mouth opening  
29 (MMO), pain, and temporomandibular disability index (TDI). A total of 100 patients were  
30 included. Significant changes were noted in MMO, pain rating, and TDI from initial  
31 evaluation to discharge from physical therapy. Sex, age, and weight did not affect the  
32 outcomes. There was also no correlation between the number of visits attended and change  
33 in MMO. Patients treated conservatively did show improvements in short term outcomes  
34 (MMO, pain rating, and TDI). These changes were statistically significant, indicating that  
35 conservative therapy may be a beneficial treatment option for patients with TMJ  
36 dysfunction. Future studies assessing the long-term outcomes of TMJ patients treated  
37 conservatively would determine if this treatment is beneficial in the long-term. In addition,  
38 researching the effectiveness of specific interventions for TMJ patients, and if certain TMJ  
39 disorders are more responsive to conservative care than others would be valuable in  
40 providing information on the effectiveness of conservative treatment in this patient  
41 population.

1 Fernández-de-Las-Peñas et al. (2020) aimed to discuss clinical reasoning based on  
2 nociceptive pain mechanisms for determining the most appropriate assessment and  
3 therapeutic strategy and to identify/map the most updated scientific evidence in relation to  
4 physical therapy interventions for patients with temporomandibular disorders (TMDs) in  
5 this narrative review. Authors conclude the following: the clinical examination of patients  
6 with TMDs should be based on nociceptive mechanisms and include the potential  
7 identification of the dominant, central, or peripheral sensitization driver. Additionally, the  
8 musculoskeletal drivers of these sensitization processes should be assessed with the aim of  
9 reproducing symptoms. Therapeutic strategies applied for managing TMDs can be grouped  
10 into tissue-based impairment treatments (bottom-up interventions) and strategies targeting  
11 the central nervous system (top-down interventions). Bottom-up strategies include joint-,  
12 soft tissue-, and nerve-targeting interventions, as well as needling therapies, whereas top-  
13 down strategies include exercises, grade motor imagery, and also pain neuroscience  
14 education. Evidence shows that the effectiveness of these interventions depends on the  
15 clinical reasoning applied, since not all strategies are equally effective for the different  
16 TMD subgroups. In fact, the presence or absence of a central sensitization driver could lead  
17 to different treatment outcomes. Authors report that it seems that multimodal approaches  
18 are more effective and should be applied in patients with TMDs. van der Meer et al. (2020)  
19 systematically evaluated the literature on the effectiveness of physical therapy on  
20 concomitant headache pain intensity in patients with TMD. Randomized or controlled  
21 clinical trials studying physical therapy interventions were included. Authors concluded  
22 physical therapy interventions presented small effect on reducing headache pain intensity  
23 on subjects with TMD, with low level of certainty. More studies of higher methodological  
24 quality are needed so better conclusions could be taken.

25  
26 Aisaiti et al. (2021) evaluated the effect of photobiomodulation therapy (PBMT) (i.e., low  
27 level laser therapy) on painful temporomandibular disorders (TMD) patients in a  
28 randomized, double-blinded, placebo-controlled manner. Participants were divided into a  
29 masseter myalgia group ( $n = 88$ ) and a temporomandibular joint (TMJ) arthralgia group ( $n = 87$ )  
30 according to the Diagnostic Criteria for Temporomandibular Disorders (DC/TMD).  
31 Both groups randomly received PBMT or placebo treatment once a day for 7 consecutive  
32 days, 1 session. The PBMT was applied with a gallium-aluminum-arsenide (GaAlAs) laser  
33 (wavelength = 810 nm) at pre-determined points in the masseter muscle (6 J/cm<sup>2</sup>, 3  
34 regions, 60 s) or TMJ region (6 J/cm<sup>2</sup>, 5 points, 30 s) according to their most painful site.  
35 Pain intensity was rated on a 0-10 numerical rating scale (NRS) and pressure pain  
36 thresholds (PPT), and mechanical sensitivity mapping were recorded before and after the  
37 treatment on day 1 and day 7. Jaw function was assessed by pain free jaw opening,  
38 maximum unassisted jaw opening, maximum assisted jaw opening, maximum protrusion  
39 and right and left excursion. Pain intensity in arthralgia patients decreased over time for  
40 both types of interventions, however, PBMT caused greater reduction in pain scores than  
41 placebo. For myalgia patients, pain intensity decreased over time but without difference  
42 between interventions. PPTs increased in both myalgia and TMJ arthralgia patients over

1 time but without difference between interventions. Overall, PBMT was associated with  
2 marginally better improvements in range of motion compared to placebo in both myalgia  
3 and arthralgia patients. Pain intensity, sensory function and jaw movements improve after  
4 both PBMT and placebo treatments in myalgia and arthralgia patients indicating a  
5 substantial non-specific effect of PBMT.

6  
7 Ahmad et al. (2021) evaluated the efficacy of LLLT in the treatment of temporomandibular  
8 joint disorder within a systematic review. Thirty-seven articles were considered eligible for  
9 this systematic review. Out of 37 studies, 33 (89.18%) were high methodological studies,  
10 which had an overall low risk of bias or with some concerns, while only 4 studies had a  
11 high risk of bias. Eighteen studies showed that LLLT was efficacious in diminishing TMD  
12 pain, whereas 12 studies showed that LLLT had similar efficacy as of  
13 placebo/controls/other intervention in TMD pain diminution. Four studies presented varied  
14 effects of LLLT on pain intensity, mandibular motion, EMG activity, and masticatory  
15 efficiency. Two studies revealed that LLLT improved the psychological and emotional  
16 aspects associated with TMDs, joint noises, masticatory efficiency, and EMG parameters,  
17 respectively. One study focused on subjective tinnitus, whereas another study suggested  
18 laser acupuncture (LAT) therapy as a suitable alternative to LLLT. The results demonstrate  
19 that LLLT appears to be efficient in diminishing TMD pain with variable effects on the  
20 outcome of secondary parameters. The results demonstrate that LLLT appears to be  
21 efficient in diminishing TMD pain with variable effects on the outcome of secondary  
22 parameters. Also, LLLT provides advantages as the therapeutic regimen is non-invasive,  
23 reversible, with fewer adverse effects, and may also improve the psychological and  
24 emotional aspects associated with TMDs. Therefore, this systematic review highlights the  
25 role of LLLT as a promising therapeutic regimen for TMDs.

26  
27 Zhang et al. (2021) compared the effects of exercise therapy and occlusal splint therapy on  
28 pain and mobility in individuals with painful temporomandibular disorders (TMD) in a  
29 systematic review. Six studies were included (498 patients: 251 occlusal splint therapy,  
30 247 therapeutic exercise). The results revealed that exercise therapy was not superior to  
31 occlusal splint therapy for pain reduction in patients with painful TMD. The effectiveness  
32 of occlusal splint therapy and exercise therapy was found to be equivalent in the maximum  
33 mouth-opening range, right laterotrusion, left laterotrusion, and protrusion for painful  
34 TMD patients. Authors concluded that given the limitations of the study, the small number  
35 of studies included in the sub-analysis for pain relief and the maximum mouth-opening  
36 range, and the small overall standardized mean difference for pain relief and mandibular  
37 movement observed, no high-quality evidence was found to distinguish the clinical  
38 effectiveness between occlusal splint therapy and exercise therapy for painful TMD  
39 patients. It appears that more randomized controlled trials comparing the effects of exercise  
40 therapy and occlusal splint therapy need to be implemented.

1 Urbański et al. (2021) compared the degree of relaxation of the anterior part of the temporal  
2 muscles and the masseter muscles, achieved through the use of post-isometric relaxation  
3 and myofascial release methods in patients requiring prosthetic treatment due to  
4 temporomandibular joint disorders with a dominant muscular component. Sixty patients  
5 who met the inclusion criteria were alternately assigned to one of the two study groups: (I)  
6 patients received post-isometric relaxation treatment (PIR), and (II) patients received  
7 myofascial release treatment (MR). The series of 10 treatments were performed in both  
8 groups. The comparative assessment was based on physiotherapeutic examination, a  
9 surface electromyography (sEMG) of the anterior temporal and masseter muscles and the  
10 intensity of spontaneous masticatory muscle pain, assessed using the Visual Analogue  
11 Scale (VAS). Authors observed a significant decrease in the electrical activity of examined  
12 muscles and a significant drop in the intensity of spontaneous pain in the masticatory  
13 muscles both in group I and II. There were no significant differences between groups. Both  
14 therapeutic methods may be used as successful forms of adjunctive therapy in the prosthetic  
15 treatment of TMD.

16  
17 Kulesa-Mrowiecka et al. (2021) aimed to present the occurrence of HJS among patients  
18 with myogenic TMD and disc displacement with reduction. The secondary goal was to  
19 assess the effectiveness of physiotherapy directed to TMD with coexisting HJS. The study  
20 involved 322 patients with symptoms of TMD. HJS was diagnosed using the Beighton  
21 Scale, which confirmed its occurrence in 26 cases. A total of 79 subjects (7 males and 72  
22 females; mean age,  $33.9 \pm 10.4$  years) were selected and divided into two groups: HJS +  
23 TMD ( $n = 26$ ; 2 males and 24 females; mean age,  $27.1 \pm 9.4$  years) and TMD ( $n = 53$ ; 5  
24 males and 48 females; mean age,  $37.4 \pm 9.2$  years). These patients completed 3 week  
25 physiotherapy management. Before and after physiotherapy, the myofascial pain severity  
26 on Numeric Pain Rating Scale, linear measurement of maximum mouth opening, and  
27 opening pattern, were assessed. A statistically significant improvement was obtained in  
28 decreasing myofascial pain in both groups. Coordination of mandibular movements was  
29 achieved in both groups. Generalized joint hypermobility occurred among patients with  
30 TMD. Physiotherapy directed to TMD was effective in reducing myofascial pain and  
31 restoring TMJ's coordination also in patients with HJS.

32  
33 Shousha et al. (2021) assessed the efficacy of low-level laser therapy (LLLT) as compared  
34 to occlusive splint therapy (OST) on the TMJ opening index (TOI) and sEMG of  
35 masticatory muscles. A total of 112 female subjects suffering from unilateral myogenous  
36 TMD, aged 21-30 years-old, were recruited and divided into 3 groups: LLLT, soft  
37 occlusive splint therapy OST, and a waitlist group as controls. Outcome measures included  
38 TMJ opening index (TOI), Visual analogue scale (VAS), and surface electromyography  
39 (sEMG). Results noted a significant reduction was reported in TOI, VAS and the sEMG  
40 within the LLLT and OST groups as well as significant decrease in all outcomes between  
41 groups in favor of the LLLT group. Authors concluded that findings supported an evident  
42 short term therapeutic effect of the LLLT on improving VAS, TOI and sEMG in females

1 suffering from myogenous TMD. Magri et al. (2021) sought to characterize short- and  
 2 long-term assessment of the low-intensity laser therapy (LLLT) effectiveness in women  
 3 with TMD of muscular origins and to evaluate whether the information about the treatment  
 4 received (active or placebo) modifies the pain intensity. Forty-one women with painful  
 5 TMD ( $31.7 \pm 5.2$  years) were divided into laser ( $n = 20$ ) and placebo ( $n = 21$ ) groups. The  
 6 pain intensity was measured at the baseline, after the LLLT (T8), 6 and 12 months. At the  
 7 6-month follow-up, the groups received information about the active or placebo treatment.  
 8 Results demonstrated that at T8 and 6-month, both active and placebo LLLT were effective  
 9 in reducing pain. After one year, the groups showed similar pain. Active LLLT was more  
 10 effective in reducing pain palpation and referred pain in the region of the TMJs. The  
 11 information about the treatment modified the perceived pain intensity. Authors concluded  
 12 that active and placebo LLLT are effective for painful TMD of muscular origins in the  
 13 short-term. Information about the treatment impairs the subjective perception of pain.

14  
 15 Dinsdale et al. (2022) evaluated the effectiveness of conservative interventions on self-  
 16 reported and physical measures of bite function in individuals with TMD in a systematic  
 17 review. Eleven studies were eligible for this review. Interventions included splinting,  
 18 photobiomodulation (PBM), needling, exercise, manual therapy, and patient education,  
 19 which were evaluated using mastication-related pain, self-reported chewing difficulty, and  
 20 bite force/endurance outcome measures. Findings suggested manual therapy, needling, oral  
 21 splinting, exercise, and PBM interventions may improve bite function in TMD, although  
 22 confidence in cumulative evidence ranged from moderate to very low. There was no  
 23 evidence that patient education improved bite function. Authors concluded that  
 24 conservative interventions may be helpful to address bite-related impairments associated  
 25 with TMD, although further research is needed to improve the quality of evidence and  
 26 direct clinical guidelines.

27  
 28 Asquini et al. (2021) aimed to evaluate the effectiveness of manual therapy applied  
 29 specifically to the craniomandibular structures (Cranio-Mandibular Manual Therapy  
 30 [CMMT]) on pain and maximum mouth opening in people with TMD. A total of 2,720  
 31 records were screened, of which only 6 (293 participants) satisfied the inclusion criteria.  
 32 All studies showed some concerns in risk of bias, except for one, which was high risk of  
 33 bias. The overall quality of evidence was very low for all outcomes because of high  
 34 heterogeneity and small sample sizes. All studies showed a significant improvement in pain  
 35 and maximum mouth opening for CMMT from baseline in the mid-term, but only 2 showed  
 36 superiorities compared to other interventions. Given the high heterogeneity and small  
 37 sample sizes of the included studies, a quantitative synthesis was not performed. Authors  
 38 concluded that there is the need for future high methodology research investigating  
 39 different manual therapy techniques applied to different regions and different populations  
 40 (e.g., chronic versus acute TMD) to determine what is most effective for pain and  
 41 maximum mouth opening in patients with TMDs. Tran et al. (2022) authored a knowledge-  
 42 to-action rapid review of systematic reviews published in the past 5 years and guidelines

1 published in the past 10 years concerning the management of TMD. In total, 62 systematic  
2 reviews and 9 guidelines considering a range of treatment modalities were included. In  
3 concordance with current guidelines, moderate evidence supports a multi-modal  
4 conservative approach towards initial management. Contrary to existing guidelines,  
5 occlusal splint therapy is not recommended due to a lack of supporting evidence. The  
6 evidence surrounding oral and topical pharmacotherapeutics for chronic TMD is low,  
7 whilst the evidence supporting injected pharmacotherapeutics is low to moderate. In  
8 concordance with current guidelines, moderate quality evidence supports the use of  
9 arthrocentesis or arthroscopy for arthrogenous TMD insufficiently managed by  
10 conservative measures, and open joint surgery for severe arthrogenous disease. Based on  
11 this, a management pathway showing escalation of treatment from conservative to invasive  
12 is proposed. La Touche et al. (2022) analyzed the effectiveness of exercise and manual  
13 therapy interventions in patients with disc displacement without reduction in a systematic  
14 review. Ten articles were included, according to the inclusion criteria. Most of the  
15 interventions showed statistically significant improvements in the primary outcomes.  
16 Results show that interventions based on therapeutic exercise or manual therapy may be  
17 beneficial and play a role in the treatment of disc displacement without reduction. Limited  
18 evidence suggests that exercise significantly improves mouth opening in comparison to  
19 splints. Due to the heterogeneity of the included studies, these results should be interpreted  
20 with caution.

21  
22 Al-Moraissi et al. (2021) aimed to identify the best treatment for adult patients with M-  
23 TMD in a network meta-analysis (NMA). Authors identified randomized clinical trials  
24 (RCTs) which are comparing 2 or more of the following treatment modalities in patients  
25 with M-TMD: counseling therapy; occlusal appliances; manual therapy; laser therapy; dry  
26 needling; intramuscular injection of local anesthesia (LA) or botulinum toxin-A (BTX-A);  
27 muscle relaxants; hypnosis/relaxation therapy; oxidative ozone therapy; and placebo or no  
28 treatment. Primary outcome variables were the reduction of pain and mechanical  
29 sensitivity. The secondary outcome was the maximal mouth opening (MMO). Included in  
30 this NMA were 52 RCTs. At the most follow up moments, manual therapy, counseling  
31 therapy, occlusal splints therapy, and needling using BTX-A or LA as well as dry needling  
32 significantly decreased post-treatment pain intensity in M-TMDs, when compared to  
33 placebo. At short term ( $\leq 5$  months), the 4 highest-ranked treatments for post-treatment pain  
34 reduction were manual therapy (83.5%, low quality evidence), ozone therapy (75.7%, very  
35 low quality evidence), counseling therapy (71.2%, moderate quality), and occlusal  
36 appliances (71.7%, moderate quality evidence). When intermediate term ( $\geq 6$  months) was  
37 considered, BTX-A (85.8%, very low quality evidence), counseling therapy (80%, low  
38 quality evidence), occlusal appliances (62.8%, low quality evidence) and hypnosis (50.6%,  
39 very low quality evidence) were the 4 highest-ranked treatments. This NMA reveals that  
40 manual therapy can be considered the most effective treatment for M-TMD, followed by  
41 counseling treatment, intramuscular injection of LA, and occlusal appliances. However,



1 considering the limitations of the studies included, and the scarcity of strong evidence, the  
2 present findings should be interpreted cautiously.

3  
4 Ekici et al. (2022) evaluated the effectiveness of high-intensity laser therapy (HILT) in the  
5 short and long term in the treatment of patients with the myogenic temporomandibular joint  
6 disorder (TMD). This prospective, double-blind, controlled clinical study was conducted  
7 on patients with myogenic TMD at a university's oral and maxillofacial surgery clinic.  
8 Seventy-six patients were randomized into 2 groups (HILT, and control group), including  
9 38 patients in one group. The patients were evaluated for pain, the range of motion of the  
10 jaw, disability, and quality of life. Assessments were performed before therapy (week 0)  
11 and after therapy (weeks 4 and 12). Data were evaluated using SPSS-20 and the level of  
12 significance was set at  $p < 0.05$ . There was no significant difference between the groups in  
13 terms of socio-demographic characteristics of the groups at the beginning of the study. In  
14 the 4th week, the VAS pain score was significantly decreased in the HILT group (47%)  
15 compared to the placebo HILT group (4%). The maximum mouth opening was  
16 significantly increased in the HILT group (27%) compared to the placebo HILT group  
17 (4%) at week 12. The HILT group showed a significant improvement in Jaw Functional  
18 Limitation Scale 20 (JFLS-20) and Oral Health Impact Profile (OHIP-14) compared to the  
19 placebo HILT group. Authors concluded that HILT is a highly effective, non-invasive  
20 therapeutic method for patients with myogenic TMD. Fertout et al. (2022) assessed the  
21 efficacy of transcutaneous electrical nerve stimulation (TENS) for the management of  
22 temporomandibular disorders (TMD) and to determine the indications and most  
23 appropriate application modalities. Fourteen articles were retained, corresponding to a total  
24 of 532 patients, among which, 285 had a TMD. Immediately after a TENS session,  
25 significant relief of pain (19.2% to 77%), significant functional improvement (mouth  
26 opening amplitude increased by between 8.7% and 19.46%), and reduced  
27 electromyographic activity of the anterior temporalis and masseter muscles were observed.  
28 However, studies comparing TENS to other physical medicine modalities (ultrasound and  
29 laser) reported equivalent results. Authors concluded that further randomized comparative  
30 clinical trials are necessary to optimize the use of TENS (program, duration of sessions,  
31 duration of treatment) for different types of TMD.

32  
33 Busse et al. (2023) completed a comparative effectiveness study of available therapies for  
34 chronic pain associated with temporomandibular disorders (TMD). Because current  
35 clinical practice guidelines are largely consensus-based and provide inconsistent  
36 recommendations, they wanted to summarize the current evidence. Based on findings,  
37 patients living with chronic pain ( $\geq 3$  months) associated with TMD, and compared with  
38 placebo or sham procedures, the guideline panel issued: (1) strong recommendations in  
39 favor of cognitive behavioral therapy (CBT) with or without biofeedback or relaxation  
40 therapy, therapist-assisted mobilization, manual trigger point therapy, supervised postural  
41 exercise, supervised jaw exercise and stretching with or without manual trigger point  
42 therapy, and usual care (such as home exercises, stretching, reassurance, and education);

1 (2) conditional recommendations in favor of manipulation, supervised jaw exercise with  
2 mobilization, CBT with non-steroidal anti-inflammatory drugs (NSAIDS), manipulation  
3 with postural exercise, and acupuncture; (3) conditional recommendations against  
4 reversible occlusal splints (alone or in combination with other interventions),  
5 arthrocentesis (alone or in combination with other interventions), cartilage supplement  
6 with or without hyaluronic acid injection, low level laser therapy (alone or in combination  
7 with other interventions), transcutaneous electrical nerve stimulation, gabapentin,  
8 botulinum toxin injection, hyaluronic acid injection, relaxation therapy, trigger point  
9 injection, acetaminophen (with or without muscle relaxants or NSAIDS), topical capsaicin,  
10 biofeedback, corticosteroid injection (with or without NSAIDS), benzodiazepines, and  $\beta$   
11 blockers; and (4) strong recommendations against irreversible oral splints, discectomy, and  
12 NSAIDS with opioids. These recommendations apply to patients living with chronic pain  
13 ( $\geq 3$  months duration) associated with TMD as a group of conditions, and do not apply to  
14 the management of acute TMD pain. When considering management options, clinicians  
15 and patients should first consider strongly recommended interventions, then those  
16 conditionally recommended in favor, then conditionally against. In doing so, shared  
17 decision making is essential to ensure patients make choices that reflect their values and  
18 preference, availability of interventions, and what they may have already tried. Further  
19 research is warranted and may alter recommendations in the future.

20  
21 Yao et al. (2023) explored the comparative effectiveness of available therapies for chronic  
22 pain associated with temporomandibular disorders (TMD). Two hundred thirty-three trials  
23 proved eligible for review, of which 153 (8,713 participants and 59 interventions or  
24 combinations of interventions) were included in network meta-analyses. All subsequent  
25 effects refer to comparisons with placebo or sham procedures. Effects on pain for 8  
26 interventions were supported by high to moderate certainty evidence. The 3 therapies  
27 probably most effective for pain relief were cognitive behavioral therapy (CBT) augmented  
28 with biofeedback or relaxation therapy for achieving the minimally important difference  
29 (MID) in pain relief of 1 cm on a 10 cm visual analogue scale: 36%, therapist-assisted jaw  
30 mobilization, and manual trigger point therapy. Five interventions were less effective, yet  
31 more effective than placebo: CBT, supervised postural exercise, supervised jaw exercise  
32 and stretching, supervised jaw exercise and stretching with manual trigger point therapy,  
33 and usual care (such as home exercises, self-stretching, reassurance). Moderate certainty  
34 evidence showed 4 interventions probably improved physical functioning: supervised jaw  
35 exercise and stretching, manipulation, acupuncture, and supervised jaw exercise and  
36 mobilization. The evidence for pain relief or physical functioning among other  
37 interventions, and all evidence for adverse events, was low or very low certainty. Authors  
38 concluded that when restricted to moderate or high certainty evidence, interventions that  
39 promote coping and encourage movement and activity were found to be most effective for  
40 reducing chronic TMD pain.

1 Gebka et al. (2023) evaluated the effectiveness of soft tissue therapy and therapeutic  
2 exercises in female patients with pain, increased masseter muscle tension, and limited  
3 mandibular mobility. The study was conducted on a group of 82 women (G1) with the Ib  
4 disorder diagnosed in DC/TMD (Ib-myofascial pain with restricted mobility). The control  
5 group (G2) consisted of 104 women without diagnosed TMDs (normal reference values  
6 for TMJ ROM and masseter muscle sEMG bioelectric activity). The G1 group was  
7 randomly divided into 3 therapeutic groups in which the therapy was carried out for 10  
8 days: therapeutic exercises (TE), manual therapy - massage and therapeutic exercises  
9 (MTM\_TE), manual therapy - post-isometric muscle relaxation (PIR) and therapeutic  
10 exercises (MTPIR\_TE). Each time after therapy, the intensity of pain and TMJ mobility  
11 were assessed. Massage, PIR, and self-therapy led to a decrease in sEMG at rest as well as  
12 in exercise. Each of the proposed forms of therapy showed a minimal clinically significant  
13 difference (MID) in the sEMG parameter at the endpoint, with the most considerable  
14 difference in the MTM\_TE group. The forms of MT used were effective in reducing the  
15 patients' pain intensity; however, a significant difference between therapies occurred after  
16 4 treatments. Analyzing the MID between methods, it was observed that self-therapy had  
17 an analgesic effect only after 8 treatments, while PIR after 3 and massage after 1 treatment.  
18 In terms of maximum mouth opening, a significant difference was obtained between  
19 monotherapy and each form of TM, i.e., massage and PIR. Analyzing mandibular lateral  
20 movements, the authors noted a significant difference in the proposed MT forms, of which  
21 massage treatments exceeded the effectiveness of PIR. Authors concluded that soft tissue  
22 manual therapy and therapeutic exercise are simple and safe interventions that can  
23 potentially benefit patients with myogenic TMDs, with massage showing better analgesic  
24 effects than PIR.

25  
26 Zhang et al. (2023) evaluated the efficacy of laser therapy in temporomandibular disorders  
27 (TMD). A total of 28 randomized controlled trials were included. Authors concluded that  
28 laser therapy can effectively reduce pain but have small effect on improving mandibular  
29 movement of TMD patients. More well-designed RCTs with large sample sizes are needed  
30 for further validation. These studies should report detailed laser parameters and provide  
31 complete outcome measure data.

32  
33 Serrano-Muñoz et al. (2023) aimed to determine the effectiveness of different electrical  
34 stimulation modalities in patients with temporomandibular disorders for reducing  
35 musculoskeletal pain, increasing the range of movement, and improving muscle activity.  
36 The main outcome measure was pain intensity. Seven studies were included in the  
37 qualitative analysis and in the quantitative analysis ( $n = 184$  subjects). The overall effect  
38 of electrical stimulation on pain reduction was statistically superior to sham/control. The  
39 overall effect on range of movement of the joint and muscle activity were not significant.  
40 Transcutaneous electrical nerve stimulation (TENS) and high-voltage current stimulation  
41 reduces pain intensity clinically in people with temporomandibular disorders with a  
42 moderate quality of evidence. On the other hand, there is no evidence of the effect of

1 different electrical stimulation modalities on range of movement and muscle activity in  
2 people with temporomandibular disorders with a moderate and low quality of evidence  
3 respectively.

4  
5 de Castro-Carletti et al. (2023) summarized the evidence from randomized controlled trials  
6 and controlled trials that examined the effectiveness of electrotherapy in the treatment of  
7 patients with orofacial pain. The overall quality of the evidence for pain intensity was very  
8 low. Although the results should be carefully used, transcutaneous electric nerve  
9 stimulation (TENS) therapy showed to be clinically superior to placebo for reducing pain  
10 after treatment and at follow-up and reduce tenderness after treatment and at follow-up in  
11 subjects with mixed temporomandibular disorders. Authors concluded that results of this  
12 systematic review support the use of TENS therapy for patients with mixed  
13 temporomandibular disorders to improve pain intensity, and tenderness demonstrating that  
14 transcutaneous electric nerve stimulation is superior to placebo. There is inconsistent  
15 evidence supporting the superiority of TENS against other therapies.

16  
17 Idáñez-Robles et al. (2023) analyzed the effectiveness of exercise therapy in improving  
18 pain and active or passive maximum mouth opening in patients with temporomandibular  
19 disorders. Randomized controlled trials evaluating the effect of exercise therapy on pain  
20 and on active and passive maximum mouth opening in patients with temporomandibular  
21 disorders were included (16 studies with 812 participants). Exercise therapy was effective  
22 in reducing pain and increasing the pain pressure threshold, active and passive maximum  
23 mouth opening. On pain pressure threshold, exercise therapy was better than physiotherapy  
24 approach (e.g., manual therapy and electrotherapy). Author concluded that therapeutic  
25 exercise is an effective therapy to reduce pain and increase pain pressure threshold and  
26 active and passive maximum mouth opening in patients with temporomandibular disorders.  
27 de Oliveira-Souza et al. (2023) determined the effectiveness of laser therapy for managing  
28 patients with orofacial pain (OFP). In addition, authors sought to determine which  
29 parameters provide the best treatment effects to reduce pain, improve function, and quality  
30 of life in adults with OFP. Eighty-nine studies were included. Most studies ( $n = 72$ , 80.9%)  
31 were considered to have a high risk of bias. The results showed that laser therapy was better  
32 than placebo in improving pain, maximal mouth open (MMO), protrusion, and tenderness  
33 at the final assessment, but with a low or moderate level of evidence. The best lasers and  
34 parameters to reduce pain were diode or gallium-aluminum-arsenide (GaAlAs) lasers, a  
35 wavelength of 400-800 or 800-1500 nm, and dosage of  $<25$  J/cm<sup>2</sup>. Authors concluded that  
36 laser therapy was better than placebo to improve pain, MMO, protrusion, and tenderness.  
37 Also, it was better than occlusal splint to improve pain, but not better than TENS and  
38 medication.

39  
40 Tanhan et al. (2023) investigated the efficacy of different types of physiotherapy  
41 approaches in individuals with cervical myofascial painful temporomandibular disorders  
42 (TMDs). Seventy-five participants with myofascial pain of jaw muscles and cervical

1 myofascial pain were randomized into 3 groups: exercise group (E), low-level laser therapy  
 2 group (LLLT), and manual pressure release group (MPR). All patients were assessed  
 3 before treatment and after 12 sessions of treatment. Significant improvement was seen in  
 4 all groups' pressure pain threshold (PPT) values. Some masticatory and neck muscles' PPT  
 5 changes in MRP and LLLT groups were significantly higher than the exercise group.  
 6 Authors concluded that exercise therapy is an effective approach for treatment of TMDs.  
 7 Additionally, LLLT combined with exercise and MPR combined with exercise have better  
 8 effects than only exercise therapy. Multimodal treatment approaches should include  
 9 exercise to achieve better results in clinical practice.

### 11 **Acupuncture**

12 Cho et al. (2010) assessed the effectiveness of acupuncture for the symptomatic treatment  
 13 of TMD. Nineteen studies were reviewed. There was moderate evidence that classical  
 14 acupuncture had a positive influence beyond those of placebo (3 trials; 65 participants);  
 15 had positive effects similar to those of occlusal splint therapy (3 trials; 160 participants);  
 16 and was more effective for TMD symptoms than physical therapy (4 trials; 397  
 17 participants), indomethacin plus vitamin B1 (2 trials; 85 participants), and a wait-list  
 18 control (3 trials; 138 participants). Only 2 RCTs addressed adverse events and reported no  
 19 serious adverse events. This review concluded that there is moderate evidence that  
 20 acupuncture is an effective intervention to reduce symptoms associated with TMD.

22 Jung et al. (2011) carried out a systematic review and meta-analysis of randomized,  
 23 placebo-controlled trials assessing the efficacy of acupuncture for treatment of TMD. A  
 24 total of 7 RCTs met the appropriate inclusion criteria for the purpose of this review. The  
 25 review and meta-analysis concluded that the evidence for acupuncture as a symptomatic  
 26 treatment of TMD is limited.

28 La Touche et al. (2010) carried out a systematic review and meta-analysis of randomized  
 29 controlled trials for the use of acupuncture treatment. A total of 4 RCTs were considered  
 30 acceptable. These 4 studies showed positive results such as reducing pain, improving  
 31 masticatory function, and increasing maximum interincisal opening. The results of this  
 32 meta-analysis suggest that acupuncture is a reasonable adjunctive treatment for producing  
 33 a short-term analgesic effect in patients with painful TMD symptoms. As a caveat, although  
 34 the results described are positive, the relevance of these results was limited by the fact that  
 35 the meta-analysis was carried out on a total of only 4 studies, representing a relatively small  
 36 global size ( $n=96$ ), which makes it more difficult to detect a sample bias. Two of the  
 37 systematic reviews (Jung; La Touche) identified essentially the same set of clinical trials.  
 38 All trials were very small, sample sizes ranging from only 10 to 20 subjects per treatment  
 39 group. The Cho review was less restrictive in its inclusion criteria and a few larger trials  
 40 were included. Notwithstanding, the evidence in this domain is limited to pilot-study-size  
 41 clinical trials.

1 Fernandes et al. (2017) sought to determine the effectiveness of acupuncture in treating  
2 myofascial pain in temporomandibular disorder (TMD) patients in a systematic review. A  
3 total of 4 randomized clinical trials using acupuncture (traditional, trigger point, and laser)  
4 for TMD treatment met the eligibility criteria and were included. Although the studies  
5 featured small sample sizes and short-term follow-up periods, acupuncture yielded results  
6 similar to those observed in groups treated with occlusal splints and were significantly  
7 superior to those obtained from placebo acupuncture-treated groups. Authors concluded  
8 that despite the weak scientific evidence supporting its efficacy, acupuncture treatment  
9 appears to relieve the signs and symptoms of pain in myofascial TMD. More controlled  
10 and randomized clinical trials with larger sample sizes are needed.

11  
12 A network meta-analysis (NMA) of RCTs was performed by Al-Moraissi et al. (2020)  
13 aiming to compare the treatment outcome of dry needling, acupuncture or wet needling  
14 using different substances in managing myofascial pain of the masticatory muscles (TMD-  
15 M). Twenty-one RCTs involving 959 patients were included. The quality of evidence of  
16 the included studies was low or very low. There was significant pain decrease after platelet-  
17 rich plasma (PRP) when compared to an active/passive placebo and acupuncture. There  
18 was a significant improvement of MMO after LA and dry needling therapy versus placebo.  
19 The 3 highest ranked treatments for short-term post-treatment pain reduction in TMD-M  
20 (1-20 days) were PRP (95.8%), followed by LA (62.5%) and dry needling (57.1%),  
21 whereas the 3 highest ranked treatments at intermediate-term follow-up (1-6 months) were  
22 LA (90.2%), dry needling (66.1%) and BTX-A (52.1%) (all very low-quality evidence).  
23 LA (96.4%) was the most effective treatment regarding the increase in MMO followed by  
24 dry needling (72.4%). Authors concluded that based on this NMA the effectiveness of  
25 needling therapy did not depend on needling type (dry or wet) or needling substance. The  
26 outcome of this NMA suggests that LA, BTX-A, granisetron and PRP hold some promise  
27 as injection therapies, but no definite conclusions can be drawn due to the low quality of  
28 evidence of the included studies. This NMA did not provide enough support for any of the  
29 needling therapies for TMD-M.

30  
31 Kalladka et al. (2021) provided an overview of the etiopathogenesis, clinical features and  
32 diagnosis of TMD, and summarized the current trends in the therapeutic management in  
33 review. Effective treatment requires a clear diagnosis based on an understanding of  
34 pathophysiologic mechanisms, a detailed history with assessment of predisposing local and  
35 systemic factors, perpetuating factors, a comprehensive clinical evaluation, and a  
36 diagnostic workup. Authors concluded that a thorough history and clinical examination are  
37 the gold standards for diagnosis of TMD. The treatment goals for TMD are to control pain,  
38 restore mandibular function and facilitate the return to normal daily activity and improve  
39 the overall quality of life of a patient. They report that based on the evidence, conservative  
40 modalities including home care regimens, pharmacotherapy, intraoral appliance therapy,  
41 local anesthetic trigger point injections, physiotherapy and complementary modalities may  
42 be beneficial in patients with TMDs.

1 Li et al. (2021) discussed the present thinking in the etiology and classification of TMD,  
2 followed by the diagnostic approach and the current trend and controversies in  
3 management. When focusing on the treatments, this review reports that physiotherapy has  
4 been suggested to be an important part in the management of TMD, which may be  
5 particularly useful for myalgia or myofascial pain. Understanding the loading of the  
6 stomatognathic system, and the existence of any tension and parafunctions, is important in  
7 delivering physiotherapy such as muscle training and changing of behavior. Evidence  
8 shows that physiotherapy is effective in treatment of TMD, in particular the headache  
9 symptoms associated with the condition; future research into this area will further ascertain  
10 these findings. For myogenous TMD, Botox injection and dry-needling techniques have  
11 been suggested. They note that Botox is not considered a standard treatment option for  
12 TMD, while dry-needling, or acupuncture, may be an effective method to reduce tension  
13 in some patients. Additionally, initial results regarding extracorporeal shock wave therapy  
14 for myogenous TMD appear to show positive results. Authors also note that there has been  
15 increasing evidence demonstrating that psychosocial assessment serves as a powerful tool  
16 in terms of predicting treatment outcome. For those patients with a significant psychosocial  
17 component, counselling seems to be a promising treatment adjunct, which might be most  
18 beneficial when included in a multimodal approach. Other conservative treatment options  
19 for TMD include stress reduction techniques and diet modification. In the past, a causative  
20 relationship between occlusion and TMD had been suggested, but it is now considered an  
21 outdated theory not supported by robust evidence, and occlusal adjustment is an  
22 irreversible treatment which is no longer supported by the recent literature.

23  
24 Liu et al. (2021) aimed to use a systematic review and meta-analysis method to understand  
25 the efficacy of warm needle acupuncture (WNA) for the treatment of TMD. The meta-  
26 analysis included 10 studies with a total of 670 patients, which included 340 patients in the  
27 experimental group and 330 patients in the control group. The data in this review showed  
28 that WNA is superior to treatments such as acupuncture alone, acupuncture therapy  
29 combined with TDP, drug therapy, and ultrasonic therapy in terms of effective rate and  
30 cure rate for the treatment of TMD. Authors concluded that this systematic review and  
31 meta-analysis provides new evidence for the effectiveness of WNA for the treatment of  
32 TMD. However, the above conclusions need to be further verified by multicenter  
33 prospective studies of larger samples and higher-quality RCTs.

34  
35 Park et al. (2023) aimed to assess the effectiveness and safety of acupuncture for TMD via  
36 a systematic review of randomized clinical trials. The qualitative analysis of randomized  
37 clinical trials with acupuncture as the intervention included 32 articles, 22 of which were  
38 included in the quantitative analysis (471 participants). Acupuncture significantly  
39 improved outcomes versus active controls. In the analysis of add-ons, acupuncture  
40 significantly improved the effect rate and pain intensity. However, the quality of evidence  
41 was determined to range from low to very low. Acupuncture in TMD significantly  
42 improved outcomes versus active controls and when add-on treatments were applied.

1 However, as the quality of evidence was determined to be low, well-designed clinical trials  
2 should be conducted in the future.

3  
4 Peixoto et al. (2023) evaluated current studies to establish and compare the efficacy of  
5 traditional and laser acupuncture in reducing the signs and symptoms of  
6 temporomandibular disorders (TMD). Six studies that evaluated the intensity of pain and  
7 the level of mouth opening of the patients submitted to acupuncture were selected, and all  
8 showed improvement. However, similar results were also observed in the groups treated  
9 with occlusal splint and placebo acupuncture. Only 1 study evaluated laser acupuncture  
10 and showed a higher proportion of patients with remission of symptoms in the experimental  
11 group. Authors concluded that the traditional acupuncture seems to relieve the signs and  
12 symptoms of TMD, as well as laser acupuncture when associated with occlusal splint.  
13 However, more rigorous and high-quality clinical trials are needed.

14 De Francisco et al. (2024) performed a qualitative and quantitative analysis of the scientific  
15 literature regarding the use of acupuncture and laser acupuncture in the treatment of pain  
16 associated with temporomandibular disorders (TMDs). The aim of this article was to assess  
17 the clinical evidence for acupuncture and laser acupuncture therapies as treatment for  
18 temporomandibular joint disorder (TMD). This systematic review includes randomized  
19 clinical trials (RCTs) of acupuncture and laser acupuncture as a treatment for TMD  
20 compared to other treatments. A total of 11 RCTs met inclusion criteria. The findings show  
21 that acupuncture is short-term helpful for reducing the severity of TMD pain with muscle  
22 origin. Meta-analysis revealed that the acupuncture group and laser acupuncture group had  
23 a higher efficacy rate than the placebo control group, showing a high efficacy of  
24 acupuncture and laser acupuncture group in the treatment of temporomandibular. In  
25 conclusion, this systematic review demonstrated that the evidence for acupuncture as a  
26 symptomatic treatment of TMD is limited. Further rigorous studies are required to establish  
27 whether acupuncture has therapeutic value.

## 28 29 **PRACTITIONER SCOPE AND TRAINING**

30 Practitioners should practice only in the areas in which they are competent based on their  
31 education training and experience. Levels of education, experience, and proficiency may  
32 vary among individual practitioners. It is ethically and legally incumbent on a practitioner  
33 to determine where they have the knowledge and skills necessary to perform such services.  
34

35 It is best practice for the practitioner to appropriately render services to a patient only if  
36 they are trained, equally skilled, and adequately competent to deliver a service compared  
37 to others trained to perform the same procedure. If the service would be most competently  
38 delivered by another health care practitioner who has more skill and expert training, it  
39 would be best practice to refer the patient to the more expert practitioner.  
40

41 Best practice can be defined as a clinical, scientific, or professional technique, method, or  
42 process that is typically evidence-based and consensus driven and is recognized by a



1 majority of professionals in a particular field as more effective at delivering a particular  
 2 outcome than any other practice (Joint Commission International Accreditation Standards  
 3 for Hospitals, 2020).

4  
 5 Depending on the practitioner’s scope of practice, training, and experience, a member’s  
 6 condition and/or symptoms during examination or the course of treatment may indicate the  
 7 need for referral to another practitioner or even emergency care. In such cases it is prudent  
 8 for the practitioner to refer the member for appropriate co-management (e.g., to their  
 9 primary care physician) or if immediate emergency care is warranted, to contact 911 as  
 10 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice  
 11 guideline for information.

### 12 13 **References**

14 Ahmad SA, Hasan S, Saeed S, Khan A, Khan M. Low-level laser therapy in  
 15 temporomandibular joint disorders: a systematic review. *J Med Life*. 2021;14(2):148-  
 16 164

17  
 18 Aisaiti A, Zhou Y, Wen Y, et al. Effect of photobiomodulation therapy on painful  
 19 temporomandibular disorders. *Sci Rep*. 2021;11(1):9049. Published 2021 Apr 27

20  
 21 Al-Badawi, E. A., Mehta, N., Forgione, A. G., Lobo, S. L., & Zawawi, K. H. (2004).  
 22 Efficacy of pulsed radio frequency energy therapy in temporomandibular joint pain and  
 23 dysfunction. *Cranio*, 22(1), 10-20

24  
 25 Al-Moraissi EA, Alradom J, Aladashi O, Goddard G, Christidis N. Needling therapies in  
 26 the management of myofascial pain of the masticatory muscles: A network meta-  
 27 analysis of randomised clinical trials. *J Oral Rehabil*. 2020;47(7):910-922

28  
 29 Al-Moraissi EA, Conti PCR, Alyahya A, Alkebsi K, Elsharkawy A, Christidis N. The  
 30 hierarchy of different treatments for myogenous temporomandibular disorders: a  
 31 systematic review and network meta-analysis of randomized clinical trials [published  
 32 correction appears in *Oral Maxillofac Surg*. 2021 Nov 5;:]. *Oral Maxillofac Surg*.  
 33 2022;26(4):519-533

34  
 35 Alves BM, Macedo CR, Januzzi E, Grossmann E, Atallah ÁN, Peccin S. Mandibular  
 36 manipulation for the treatment of temporomandibular disorder. *J Craniofac Surg*. 2013  
 37 Mar;24(2):488-93

38  
 39 Armijo-Olivo S, Pitance L, Singh V, Neto F, Thie N, Michelotti A. Effectiveness of  
 40 Manual Therapy and Therapeutic Exercise for Temporomandibular Disorders:  
 41 Systematic Review and Meta-Analysis. *Phys Ther*. 2016 Jan;96(1):9-25

- 1 Asquini G, Pitance L, Michelotti A, Falla D. Effectiveness of manual therapy applied to  
2 craniomandibular structures in temporomandibular disorders: A systematic review. *J*  
3 *Oral Rehabil.* 2021;49(4):442-455.
- 4
- 5 Brantingham, J. W., Cassa, T. K., Bonnefin, D., Pribicevic, M., Robb, A., Pollard, H.,  
6 Korporaal, C. (2013). Manipulative and Multimodal Therapy for Upper Extremity and  
7 Temporomandibular Disorders: A Systematic Review. *Journal of manipulative and*  
8 *physiological therapeutics*, 36(3), 143-201
- 9
- 10 Bronfort, G., Haas, M., Evans, R., Leininger, B., & Triano, J. (2010). Effectiveness of  
11 manual therapies: the UK evidence report. *Chiropractic & Osteopathy*, 18(3), 1-33
- 12
- 13 Busse JW, Casassus R, Carrasco-Labra A, Durham J, Mock D, Zakrzewska JM, Palmer C,  
14 Samer CF, Coen M, Guevremont B, Hoppe T, Guyatt GH, Crandon HN, Yao L,  
15 Sadeghirad B, Vandvik PO, Siemieniuk RAC, Lytvyn L, Hunskaar BS, Agoritsas T.  
16 Management of chronic pain associated with temporomandibular disorders: a clinical  
17 practice guideline. *BMJ.* 2023 Dec 15;383:e076227
- 18
- 19 Butts R, Dunning J, Pavkovich R, Mettelle J, Mourad F. Conservative management of  
20 temporomandibular dysfunction: A literature review with implications for clinical  
21 practice guidelines (Narrative review part 2). *J Bodyw Mov Ther.* 2017 Jul;21(3):541-  
22 548
- 23
- 24 Calixtre LB, Moreira RF, Franchini GH, Albuquerque-Sendín F, Oliveira AB. Manual  
25 therapy for the management of pain and limited range of motion in subjects with signs  
26 and symptoms of temporomandibular disorder: a systematic review of randomized  
27 controlled trials. *J Oral Rehabil.* 2015 Nov;42(11):847-61
- 28
- 29 Calixtre LB, Oliveira AB, de Sena Rosa LR, Armijo-Olivo S, Visscher CM, Albuquerque-  
30 Sendín F. Effectiveness of mobilisation of the upper cervical region and craniocervical  
31 flexor training on orofacial pain, mandibular function and headache in women with  
32 TMD. A randomised, controlled trial. *J Oral Rehabil.* 2019 Feb;46(2):109-119
- 33
- 34 Carmeli, E., Sheklow, S. L., & Bloomenfeld, I. (2001). Comparative Study of  
35 Repositioning Splint Therapy and Passive Manual Range of Motion Techniques for  
36 Anterior Displaced Temporomandibular Discs with Unstable Excursive Reduction.  
37 *Physiotherapy*, 87(1), 26-36
- 38
- 39 Chen J, Huang Z, Ge M, Gao M. Efficacy of low-level laser therapy in the treatment of  
40 TMDs: a meta-analysis of 14 randomised controlled trials. *J Oral Rehabil.* 2015  
41 *Apr*;42(4):291-9

- 1 Cho, S. H., & Whang, W. W. (2010). Acupuncture for temporomandibular disorders: a  
2 systematic review. *J Orofac Pain*, 24(2), 152-162  
3
- 4 Chortis, A. G., Chorti, A. G., Forrester, G., & Georgoudis, G. (2006). Therapeutic exercise  
5 in the management of anterior disc displacement of the temporomandibular joint.  
6 *Physical Therapy Reviews*, 11(2), 117-123  
7
- 8 Cuccia, A. M., Caradonna, C., & Caradonna, D. (2011). Manual Therapy of the Mandibular  
9 Accessory Ligaments for the Management of Temporomandibular Joint Disorders.  
10 *JAOA: Journal of the American Osteopathic Association*, 111(2), 102-112  
11
- 12 de Castro-Carletti EM, Müggendorf F, Dennett L, Sobral de Oliveira-Souza AI, Mohamad  
13 N, Pertille A, Rodrigues-Bigaton D, Armijo-Olivo S. Effectiveness of electrotherapy  
14 for the treatment of orofacial pain: A systematic review and meta-analysis. *Clin  
15 Rehabil*. 2023 Jul;37(7):891-926  
16
- 17 de Oliveira-Souza AIS, Mohamad N, de Castro Carletti EM, Müggendorf F, Dennett L, de  
18 Oliveira DA, Armijo-Olivo S. What are the best parameters of low-level laser therapy  
19 to reduce pain intensity and improve mandibular function in orofacial pain? A  
20 systematic review and meta-analysis. *Disabil Rehabil*. 2023 Oct;45(20):3219-3237  
21
- 22 Delgado de la Serna P, Plaza-Manzano G, Cleland J, Fernández-de-Las-Peñas C, Martín-  
23 Casas P, Díaz-Arribas MJ. Effects of Cervico-Mandibular Manual Therapy in Patients  
24 with Temporomandibular Pain Disorders and Associated Somatic Tinnitus: A  
25 Randomized Clinical Trial. *Pain Med*. 2020 Mar 1;21(3):613-624  
26
- 27 Di Francesco F, Minervini G, Siurkel Y, Ciccì M, Lanza A. Efficacy of acupuncture and  
28 laser acupuncture in temporomandibular disorders: a systematic review and meta-  
29 analysis of randomized controlled trials. *BMC Oral Health*. 2024 Feb 3;24(1):174  
30
- 31 Dinsdale A, Costin B, Dharamdasani S, Page R, Purs N, Treleaven J. What conservative  
32 interventions improve bite function in those with temporomandibular disorders? A  
33 systematic review using self-reported and physical measures. *J Oral Rehabil*.  
34 2022;49(4):456-475  
35
- 36 Ekici Ö, Dündar Ü, Büyükbosna M. Effectiveness of high-intensity laser therapy in patients  
37 with myogenic temporomandibular joint disorder: A double-blind, placebo-controlled  
38 study. *J Stomatol Oral Maxillofac Surg*. 2022;123(3):e90-e96  
39
- 40 Fernandes AC, Duarte Moura DM, Da Silva LGD, De Almeida EO, Barbosa GAS.  
41 Acupuncture in Temporomandibular Disorder Myofascial Pain Treatment: A  
42 Systematic Review. *J Oral Facial Pain Headache*. 2017 Summer;31(3):225-232

- 1 Fernandez-Carnero, J., La Touche, R., Ortega-Santiago, R., Galan-del-Rio, F., Pesquera,  
2 J., Ge, H. Y., & Fernandez-de-Las-Penas, C. (2010). Short-term effects of dry needling  
3 of active myofascial trigger points in the masseter muscle in patients with  
4 temporomandibular disorders. *J Orofac Pain*, 24(1), 106-112
- 5
- 6 Fernández-de-Las-Peñas C, Von Piekartz H. Clinical Reasoning for the Examination and  
7 Physical Therapy Treatment of Temporomandibular Disorders (TMD): A Narrative  
8 Literature Review. *J Clin Med*. 2020 Nov 17;9(11):3686
- 9
- 10 Fertout A, Manière-Ezvan A, Lupi L, Ehrmann E. Management of temporomandibular  
11 disorders with transcutaneous electrical nerve stimulation: A systematic review.  
12 *Cranio*. 2022;40(3):217-228
- 13
- 14 Fisch G, Finke A, Ragonese J, Dugas L, Wrzosek M. Outcomes of physical therapy in  
15 patients with temporomandibular disorder: a retrospective review. *Br J Oral Maxillofac*  
16 *Surg*. 2020 Aug 20:S0266-4356(20)30459-9
- 17
- 18 Garrigós-Pedron M, La Touche R, Navarro-Desentre P, Gracia-Naya M, Segura-Ortí E.  
19 Effects of a Physical Therapy Protocol in Patients with Chronic Migraine and  
20 Temporomandibular Disorders: A Randomized, Single-Blinded, Clinical Trial. *J Oral*  
21 *Facial Pain Headache*. 2018 Spring;32(2):137-150
- 22
- 23 Gauer RL, Semidey MJ. Diagnosis and treatment of temporomandibular disorders. *Am*  
24 *Fam Physician*. 2015 Mar 15;91(6):378-86
- 25
- 26 Gębska M, Dalewski B, Pałka Ł, Kołodziej Ł. Evaluation of the efficacy of manual soft  
27 tissue therapy and therapeutic exercises in patients with pain and limited mobility TMJ:  
28 a randomized control trial (RCT). *Head Face Med*. 2023 Sep 8;19(1):42
- 29
- 30 George, J. W., Fennema, J., Maddox, A., Nessler, M., & Skaggs, C. D. (2007). The effect  
31 of cervical spine manual therapy on normal mouth opening in asymptomatic subjects.  
32 *J Chiropr Med*, 6(4), 141-145
- 33
- 34 Grace, E. G., Sarlani, E., & Reid, B. (2002). The use of an oral exercise device in the  
35 treatment of muscular TMD. *Cranio*, 20(3), 204-208
- 36
- 37 Huggins, T., Boras, A. L., Gleberzon, B. J., Popescu, M., & Bahry, L. A. (2012). Clinical  
38 effectiveness of the activator adjusting instrument in the management of  
39 musculoskeletal disorders: a systematic review of the literature. *J Can Chiropr Assoc*,  
40 56(1), 49-57

- 1 Idáñez-Robles AM, Obrero-Gaitán E, Lomas-Vega R, Osuna-Pérez MC, Cortés-Pérez I,  
2 Zagalaz-Anula N. Exercise therapy improves pain and mouth opening in  
3 temporomandibular disorders: A systematic review with meta-analysis. *Clin Rehabil.*  
4 2023 Apr;37(4):443-461
- 5
- 6 Ismail, F., Demling, A., Hessling, K., Fink, M., & Stiesch-Scholz, M. (2007). Short-term  
7 efficacy of physical therapy compared to splint therapy in treatment of arthrogenous  
8 TMD. *J Oral Rehabil*, 34(11), 807-813
- 9
- 10 Jung, A., Shin, B. C., Lee, M. S., Sim, H., & Ernst, E. (2011). Acupuncture for treating  
11 temporomandibular joint disorders: a systematic review and meta-analysis of  
12 randomized, sham-controlled trials. *J Dent*, 39(5), 341-350
- 13
- 14 Kalamir, A., Bonello, R., Graham, P., Vitiello, A. L., & Pollard, H. (2012). Intraoral  
15 Myofascial Therapy for Chronic Myogenous Temporomandibular Disorder: A  
16 Randomized Controlled Trial. *Journal of manipulative and physiological therapeutics*,  
17 35(1), 26-37
- 18
- 19 Kalamir, A., Graham, P., Vitiello, A., Bonello, R., & Pollard, H. (2013). Intra-oral  
20 myofascial therapy versus education and self-care in the treatment of chronic,  
21 myogenous temporomandibular disorder: a randomised, clinical trial. *Chiropractic &*  
22 *Manual Therapies*, 21(1), 17
- 23
- 24 Kalamir, A., Pollard, H., Vitiello, A., & Bonello, R. (2010). Intra-oral myofascial therapy  
25 for chronic myogenous temporomandibular disorders: a randomized, controlled pilot  
26 study. *J Man Manip Ther*, 18(3), 139-146
- 27
- 28 Kalladka M, Young A, Khan J. Myofascial pain in temporomandibular disorders: Updates  
29 on etiopathogenesis and management. *J Bodyw Mov Ther.* 2021;28:104-113
- 30
- 31 Komiyama, O., Kawara, M., Arai, M., Asano, T., & Kobayashi, K. (1999). Posture  
32 correction as part of behavioural therapy in treatment of myofascial pain with limited  
33 opening. *J Oral Rehabil*, 26(5), 428-435
- 34
- 35 Kraaijenga S, van der Molen L, van Tinteren H, Hilgers F, Smeele L. Treatment of  
36 myogenic temporomandibular disorder: a prospective randomized clinical trial,  
37 comparing a mechanical stretching device (TheraBite®) with standard physical therapy  
38 exercise. *Cranio.* 2014 Jul;32(3) :208-16
- 39
- 40 Kulesa-Mrowiecka M, Piech J, Gaździk TS. The Effectiveness of Physical Therapy in  
41 Patients with Generalized Joint Hypermobility and Concurrent Temporomandibular

- 1 Disorders-A Cross-Sectional Study. *J Clin Med.* 2021;10(17) :3808. Published 2021  
2 Aug 25
- 3
- 4 La Touche, R., Goddard, G., De-la-Hoz, J. L., Wang, K., Paris-Aleman, A., Angulo-Diaz-  
5 Parreno, S., Hernandez, M. (2010). Acupuncture in the treatment of pain in  
6 temporomandibular disorders: a systematic review and meta-analysis of randomized  
7 controlled trials. *Clin J Pain*, 26(6), 541-550
- 8
- 9 La Touche R, Boo-Mallo T, Zarzosa-Rodríguez J, Paris-Aleman A, Cuenca-Martínez F,  
10 Suso-Martí L. Manual therapy and exercise in temporomandibular joint disc  
11 displacement without reduction. A systematic review. *Cranio.* 2022;40(5):440-450
- 12
- 13 Li DTS, Leung YY. Temporomandibular Disorders: Current Concepts and Controversies  
14 in Diagnosis and Management. *Diagnostics (Basel).* 2021;11(3):459. Published 2021  
15 Mar 6
- 16
- 17 List T, Axelsson S. Management of TMD: evidence from systematic reviews and meta-  
18 analyses. *J Oral Rehabil.* 2010 May;37(6):430-51
- 19
- 20 Liu GF, Gao Z, Liu ZN, Yang M, Zhang S, Tan TP. Effects of Warm Needle Acupuncture  
21 on Temporomandibular Joint Disorders : A Systematic Review and Meta-Analysis of  
22 Randomized Controlled Trials. *Evid Based Complement Alternat Med.*  
23 2021;2021 :6868625. Published 2021 Nov 27
- 24
- 25 Madani A, Ahrari F, Fallahrastegar A, Daghestani N. A randomized clinical trial  
26 comparing the efficacy of low-level laser therapy (LLLT) and laser acupuncture  
27 therapy (LAT) in patients with temporomandibular disorders. *Lasers Med Sci.* 2020  
28 Feb;35(1) :181-192
- 29
- 30 Maizlin, Z. V., Nutiu, N., Dent, P. B., Vos, P. M., Fenton, D. M., Kirby, J. M., Clement, J.  
31 J. (2010). Displacement of the temporomandibular joint disk: correlation between  
32 clinical findings and MRI characteristics. *J Can Dent Assoc*, 76, a3
- 33
- 34 Magri LV, Bataglion C, Leite-Panissi CRA. Follow-up results of a randomized clinical  
35 trial for low-level laser therapy in painful TMD of muscular origins. *Cranio.*  
36 2021;39(6):502-509
- 37
- 38 Martins WR, Blasczyk JC, Aparecida Furlan de Oliveira M, Lagôa Gonçalves KF, Bonini-  
39 Rocha AC, Dugailly PM, de Oliveira RJ. Efficacy of musculoskeletal manual approach  
40 in the treatment of temporomandibular joint disorder: A systematic review with meta-  
41 analysis. *Man Ther.* 2016 Feb;21:10-7

- 1 Matheson EM, Fermo JD, Blackwelder RS. Temporomandibular Disorders: Rapid  
2 Evidence Review. *Am Fam Physician*. 2023;107(1):52-58  
3
- 4 McNeely, M. L., Armijo Olivo, S., & Magee, D. J. (2006). A systematic review of the  
5 effectiveness of physical therapy interventions for temporomandibular disorders. *Phys  
6 Ther*, 86(5), 710-725  
7
- 8 Minakuchi, H., Kuboki, T., Matsuka, Y., Maekawa, K., Yatani, H., & Yamashita, A.  
9 (2001). Randomized controlled evaluation of non-surgical treatments for  
10 temporomandibular joint anterior disk displacement without reduction. *J Dent Res*,  
11 80(3), 924-928  
12
- 13 Moraes Ada R, Sanches ML, Ribeiro EC, Guimarães AS. Therapeutic exercises for the  
14 control of temporomandibular disorders. *Dental Press J Orthod*. 2013 Sep-  
15 Oct;18(5):134-9  
16
- 17 Nahian A, ÜNAL M, Mathew Jr J. Osteopathic Manipulative Treatment: Facial Muscle  
18 Energy, Direct MFR, and BLT Procedure – for TMJ Dysfunction. In: *StatPearls*.  
19 Treasure Island (FL): StatPearls Publishing; September 17, 2021  
20
- 21 Park EY, Cho JH, Lee SH, Kim KW, Ha IH, Lee YJ. Is acupuncture an effective treatment  
22 for temporomandibular disorder?: A systematic review and meta-analysis of  
23 randomized controlled trials. *Medicine (Baltimore)*. 2023 Sep 22;102(38):e34950  
24
- 25 Peixoto KO, Abrantes PS, De Carvalho IHG, De Almeida EO, Barbosa GAS.  
26 Temporomandibular disorders and the use of traditional and laser acupuncture: a  
27 systematic review. *Cranio*. 2023 Nov;41(6):501-507\  
28
- 29 Rashid, A., Matthews, N. S., & Cowgill, H. (2013). Physiotherapy in the management of  
30 disorders of the temporomandibular joint—perceived effectiveness and access to  
31 services: a national United Kingdom survey. *The British journal of oral &  
32 maxillofacial surgery*, 51(1), 52–57. .org/10.1016/j.bjoms.2012.03.009  
33
- 34 Reynolds B, Puentedura EJ, Kolber MJ, Cleland JA. Effectiveness of Cervical Spine High-  
35 Velocity, Low-Amplitude Thrust Added to Behavioral Education, Soft Tissue  
36 Mobilization, and Exercise for People With Temporomandibular Disorder With  
37 Myalgia: A Randomized Clinical Trial. *J Orthop Sports Phys Ther*. 2020  
38 Aug;50(8):455-465  
39
- 40 Serrano-Muñoz D, Beltran-Alacreu H, Martín-Caro Álvarez D, Fernández-Pérez JJ,  
41 Aceituno-Gómez J, Arroyo-Fernández R, Avendaño-Coy J. Effectiveness of Different  
42 Electrical Stimulation Modalities for Pain and Masticatory Function in

- 1 Temporomandibular Disorders: A Systematic Review and Meta-Analysis. *J Pain*. 2023  
2 Jun;24(6):946-956  
3
- 4 Shaffer SM, Brismée JM, Sizer PS, Courtney CA. Temporomandibular disorders. Part 2:  
5 conservative management. *J Man Manip Ther*. 2014 Feb;22(1):13-23  
6
- 7 Shaffer SM, Naze GS. Evaluation and management of temporomandibular disorders. Part  
8 1: an orthopedic physical therapy update on examination and clinical reasoning. *J Man  
9 Manip Ther*. 2023 Jun;31(3):133-142  
10
- 11 Shaffer SM, Naze GS. Evaluation and management of temporomandibular disorders. Part  
12 2: an orthopaedic physical therapy update on examination and clinical reasoning. *J Man  
13 Manip Ther*. 2023 Jun;31(3):143-152  
14
- 15 Sharma, S., Gupta, D. S., Pal, U. S., & Jurel, S. K. (2011). Etiological factors of  
16 temporomandibular joint disorders. *Natl J Maxillofac Surg*, 2(2), 116-119  
17
- 18 Shen, Y. F., Younger, J., Goddard, G., & Mackey, S. (2009). Randomized clinical trial of  
19 acupuncture for myofascial pain of the jaw muscles. *J Orofac Pain*, 23(4), 353-359  
20
- 21 Shimada A, Ishigaki S, Matsuka Y, Komiyama O, Torisu T, Oono Y, Sato H, Naganawa  
22 T, Mine A, Yamazaki Y, Okura K, Sakuma Y, Sasaki K. Effects of exercise therapy on  
23 painful temporomandibular disorders. *J Oral Rehabil*. 2019 May;46(5):475-481  
24
- 25 Shousha T, Alayat M, Moustafa I. Effects of low-level laser therapy versus soft occlusive  
26 splints on mouth opening and surface electromyography in females with  
27 temporomandibular dysfunction: A randomized-controlled study. *PLoS One*.  
28 2021;16(10):e0258063. Published 2021 Oct 1  
29
- 30 Smith, P., Moss crop, D., Davies, S., Sloan, P., & Al-Ani, Z. (2007). The efficacy of  
31 acupuncture in the treatment of temporomandibular joint myofascial pain: a  
32 randomised controlled trial. *J Dent*, 35(3), 259-267  
33
- 34 Tanhan A, Ozer AY, Polat MG. Efficacy of different combinations of physiotherapy  
35 techniques compared to exercise and patient education in temporomandibular  
36 disorders: A randomized controlled study. *Cranio*. 2023 Jul;41(4):389-401  
37
- 38 T.O., S. (2004). Physiotherapy in the management of TMD: A review of the literature part  
39 1 *International Journal of Therapy and Rehabilitation*, 11(12), 574-582  
40
- 41 Tough, E. A., White, A. R., Cummings, T. M., Richards, S. H., & Campbell, J. L. (2009).  
42 Acupuncture and dry needling in the management of myofascial trigger point pain: a



- 1 systematic review and meta-analysis of randomized controlled trials. *Eur J Pain*, 13(1),  
2 3-10
- 3
- 4 Tran C, Ghahreman K, Huppa C, Gallagher JE. Management of temporomandibular  
5 disorders: a rapid review of systematic reviews and guidelines. *Int J Oral Maxillofac*  
6 *Surg*. 2022;51(9):1211-1225
- 7
- 8 Treacy, K. (1999). Awareness/relaxation training and transcutaneous electrical neural  
9 stimulation in the treatment of bruxism. *J Oral Rehabil*, 26(4), 280-287
- 10
- 11 Turp, J. C. (2011). Limited evidence that acupuncture is effective for treating  
12 temporomandibular disorders. *Evid Based Dent*, 12(3), 89
- 13
- 14 Urbański P, Trybulec B, Pihut M. The Application of Manual Techniques in Masticatory  
15 Muscles Relaxation as Adjunctive Therapy in the Treatment of Temporomandibular  
16 Joint Disorders. *Int J Environ Res Public Health*. 2021;18(24):12970. Published 2021  
17 Dec 8
- 18
- 19 Van der Meer HA, Calixtre LB, Engelbert RHH, Visscher CM, Nijhuis-van der Sanden  
20 MW, Speksnijder CM. Effects of physical therapy for temporomandibular disorders on  
21 headache pain intensity: A systematic review [published correction appears in  
22 *Musculoskelet Sci Pract*. 2021 Jun;53:102344]. *Musculoskelet Sci Pract*.  
23 2020;50:102277
- 24
- 25 Vier C, Almeida MB, Neves ML, Santos ARSD, Bracht MA. The effectiveness of dry  
26 needling for patients with orofacial pain associated with temporomandibular  
27 dysfunction: a systematic review and meta-analysis. *Braz J Phys Ther*. 2019 Jan -  
28 Feb;23(1):3-11
- 29
- 30 Wieckiewicz M, Boening K, Wiland P, Shiau YY, Paradowska-Stolarz A. Reported  
31 concepts for the treatment modalities and pain management of temporomandibular  
32 disorders. *J Headache Pain*. 2015;16:106
- 33
- 34 Wieselmann-Penkner, K., Janda, M., Lorenzoni, M., & Polansky, R. (2001). A comparison  
35 of the muscular relaxation effect of TENS and EMG-biofeedback in patients with  
36 bruxism. *J Oral Rehabil*, 28(9), 849-853
- 37
- 38 Wright, E. F., Domenech, M. A., & Fischer, J. R., Jr. (2000). Usefulness of posture training  
39 for patients with temporomandibular disorders. *J Am Dent Assoc*, 131(2), 202-210
- 40
- 41 Wright, E. F., & North, S. L. (2009). Management and treatment of temporomandibular  
42 disorders: a clinical perspective. *J Man Manip Ther*, 17(4), 247-254

- 1 Yao L, Sadeghirad B, Li M, Li J, Wang Q, Crandon HN, Martin G, Morgan R, Florez ID,  
2 Hunskaar BS, Wells J, Moradi S, Zhu Y, Ahmed MM, Gao Y, Cao L, Yang K, Tian J,  
3 Li J, Zhong L, Couban RJ, Guyatt GH, Agoritsas T, Busse JW. Management of chronic  
4 pain secondary to temporomandibular disorders: a systematic review and network  
5 meta-analysis of randomised trials. *BMJ*. 2023 Dec 15;383:e076226  
6
- 7 Zhang L, Xu L, Wu D, Yu C, Fan S, Cai B. Effectiveness of exercise therapy versus  
8 occlusal splint therapy for the treatment of painful temporomandibular disorders: a  
9 systematic review and meta-analysis. *Ann Palliat Med*. 2021;10(6):6122-6132  
10
- 11 Zhang Y, Qian Y, Huo K, Liu J, Huang X, Bao J. Efficacy of laser therapy for  
12 temporomandibular disorders: A systematic review and meta-analysis. *Complement  
13 Ther Med*. 2023 Jun;74:102945