

1 **Clinical Practice Guideline:** **Physical Activity**

2
3 **Date of Implementation:** **June 19, 2014**

4
5 **Product:** **Specialty**

6
7
8 **GUIDELINES**

9 Among portal of entry practitioners, screening at risk and/or symptomatic patients for level
10 of physical activity is considered best practices. Providing a direct intervention (e.g.,
11 lifestyle and/or dietary changes) for patients for whom the screening results indicated a
12 need for intervention, will depend upon the practitioner’s education, training, experience,
13 and scope of practice. In the absence of such a direct intervention, providing a referral
14 intervention (e.g., to the patient’s medical physician) is necessary for adult patients
15 identified with low levels of physical activity or when other concerns arise regarding the
16 patient’s physical activity history. The screening described in this policy may be outside
17 the education, training, experience, or scope of some practitioner types. In the context of
18 best practices for these practitioners, a level of awareness that risk factors and/or
19 signs/symptoms of a physical activity issue are present is required and a subsequent referral
20 for appropriate evaluation is necessary and within the purview of all.

21
22 **INTRODUCTION**

23 Inadequate physical activity is epidemic in the U.S. Sedentary behavior and physical
24 inactivity are among the leading modifiable risk factors worldwide for cardiovascular
25 disease and all-cause mortality (Zhang and Liu, 2024; Mokdad, 2004; Danaei et al., 2009;
26 Johnson et al., 2014; Lavie et al., 2019). The Centers for Disease Control and Prevention
27 (CDC) analyzed the data from a 2020 national survey of American adults (National Health
28 Interview Survey) and concluded most are not meeting the federal physical activity
29 recommendations. Only 24% meet the minimum guidelines for aerobic exercise and
30 muscle-strengthening activity. However, 47% met the aerobic activity recommendations.

31
32 Sedentary lifestyle contributes to risk of cardiovascular disease, hypertension, type 2
33 diabetes, obesity, and osteoporosis. Conversely, regular physical activity has been shown
34 to help prevent these conditions. Furthermore, some evidence also indicates that regular
35 physical activity contributes to a reduction in all-cause morbidity and to an increase in
36 cognition, brain function, and lifespan (CDC, 2022).

37
38 **ASSESSING PHYSICAL ACTIVITY STATUS**

39 Generally, physical activity is assessed by objective findings and/or self-reported activities
40 (e.g., questionnaires or a patient diary). Objective, direct measurement methods include
41 instruments such as pedometers or motion detectors which can be used by individuals to
42 record activities over a specified period of time or point of service monitored exercise tests.

1 Among self-reported activities, questionnaires are the most feasible method for clinical
 2 practice. The patient is questioned on recall of recent or usual activities or in sedentary
 3 behaviors, usually over a specific period of time.

4
 5 National surveys conducted by government agencies usually question respondents about
 6 compliance with national standards such as those of the CDC discussed earlier or Healthy
 7 People 2020. No standards have been set for how physical activity assessment should be
 8 conducted in clinical practice; in most trials of physical activity counseling in clinical
 9 practice, self-reported measures were used.

10
 11 Recommendations for physical activity, from the first national physical activity guidelines
 12 put forth by the U.S. Department of Health and Human Services (DHHS, 2nd ed) in 2018,
 13 continue to apply today. These guidelines for adults are summarized below:

- 14 • **Adults**
 - 15 ○ Adults should move more and sit less throughout the day. Some physical
 16 activity is better than none. Adults who sit less and do any amount of moderate-
 17 to-vigorous physical activity gain some health benefits.
 - 18 ○ For substantial health benefits, adults should do at least 150 minutes (2 hours
 19 and 30 minutes) to 300 minutes (5 hours) a week of moderate-intensity, or 75
 20 minutes (1 hour and 15 minutes) to 150 minutes (2 hours and 30 minutes) a
 21 week of vigorous-intensity aerobic physical activity, or an equivalent
 22 combination of moderate- and vigorous-intensity aerobic activity. Preferably,
 23 aerobic activity should be spread throughout the week.
 - 24 ○ Additional health benefits are gained by engaging in physical activity beyond
 25 the equivalent of 300 minutes (5 hours) of moderate-intensity physical activity
 26 a week.
 - 27 ○ Adults should also do muscle-strengthening activities of moderate or greater
 28 intensity and that involve all major muscle groups on 2 or more days a week,
 29 as these activities provide additional health benefits.
- 30
 31 • **Children and Adolescents**
 - 32 ○ It is important to provide young people opportunities and encouragement to
 33 participate in physical activities that are appropriate for their age, that are
 34 enjoyable, and that offer variety.
 - 35 ○ Children and adolescents ages 6 through 17 years should do 60 minutes (1 hour)
 36 or more of moderate-to-vigorous physical activity daily:
 - 37 ■ **Aerobic:** Most of the 60 minutes or more per day should be either moderate
 38 or vigorous-intensity aerobic physical activity and should include vigorous
 39 intensity physical activity on at least 3 days a week.
 - 40 ■ **Muscle-strengthening:** As part of their 60 minutes or more of daily physical
 41 activity, children and adolescents should include muscle-strengthening

1 physical activity on at least 3 days a week. Muscle-strengthening includes
 2 activities like climbing or doing push-ups.

- 3 ■ Bone-strengthening: As part of their 60 minutes or more of daily physical
 4 activity, children and adolescents should include bone-strengthening
 5 physical activity on at least 3 days a week. Bone-strengthening includes
 6 activities such as jumping or running.

- 7
- 8 ● Older Adults

- 9 ○ As part of their weekly physical activity, older adults should do
 10 multicomponent physical activity that includes balance training as well as
 11 aerobic and muscle strengthening activities.
- 12 ○ Older adults should determine their level of effort for physical activity relative
 13 to their level of fitness.
- 14 ○ Older adults with chronic conditions should understand whether and how their
 15 conditions affect their ability to do regular physical activity safely.
- 16 ○ When older adults cannot do 150 minutes of moderate-intensity aerobic activity
 17 a week because of chronic conditions, they should be as physically active as
 18 their abilities and conditions allow.

19

20 For additional and more extensive health benefits, adults should increase their aerobic
 21 physical activity to 300 minutes (5 hours) a week of moderate-intensity, or 150 minutes a
 22 week of vigorous-intensity aerobic physical activity, or an equivalent combination of
 23 moderate- and vigorous-intensity activity. Additional health benefits are gained by
 24 engaging in physical activity beyond this amount.

25

26 The following table summarizes aerobic activity with respect to health benefits.

27

Classification of Total Weekly Aerobic Activity		
Level	Moderate intensity activity* (minutes/week)	Health Benefits
Inactive	No activity beyond baseline	None
Low	> baseline but < 150 minutes	Some
Medium	150 - 300 minutes	Substantial
High	> 300 minutes	Additional

28

29 *Moderate intensity exercise increases heart and breathing rates, but the individual can
 30 carry on a conversation. Brisk walking is the most common example of moderate intensity.

1 Current guidelines highlight the following:

- 2 • High volumes of moderate-to-vigorous physical activity appear to remove the
- 3 excess risk of all-cause mortality that is associated with high volumes of sitting.
- 4 • Very low time spent sitting reduces, but does not eliminate, the risk of no moderate-
- 5 to-vigorous physical activity.
- 6 • Given the high levels of sitting and low levels of physical activity in the population,
- 7 most people would benefit from both increasing moderate-to-vigorous physical
- 8 activity and reducing time spent sitting.

9
10 Specifically, regular physical activity helps improve health by:

- 11 • Reducing the risk of developing diabetes, hypertension, or dying from heart disease
- 12 • Reducing blood pressure among people who have hypertension
- 13 • Helping to control weight and build/maintain healthy bones, muscles, and joints
- 14 • Increasing the likelihood of living longer
- 15 • Reducing the risk of developing several cancers
- 16 • Improving older adults' ability to perform daily activities and prevent falls
- 17 • Improving mental health and reducing feelings of depression or anxiety

18
19 Other key findings from the 2018 guidelines includes:

20 **For people who are inactive**, that is, people who do not do any moderate- or vigorous-

21 intensity physical activity beyond basic movement from daily life activities:

- 22 • Reducing sedentary behavior has health benefits. It reduces the risk of all-cause
- 23 mortality, cardiovascular disease incidence and mortality, and the incidence of type
- 24 2 diabetes and some cancers. A good first step is to replace sedentary behavior with
- 25 light-intensity physical activity. Previously, evidence that light intensity physical
- 26 activity could provide health benefits was not sufficient to support a
- 27 recommendation.
- 28 • No matter how much time they spend in sedentary behavior or light-intensity
- 29 activity, inactive people can reduce their health risks by gradually increasing their
- 30 moderate-intensity physical activity.

31
32 **For people who are insufficiently active**, that is, people who do some moderate- or

33 vigorous-intensity physical activity, but who do not yet meet the key guidelines target

34 range (150 to 300 minutes a week of moderate-intensity physical activity for adults):

- 35 • Even small increases in moderate-intensity physical activity provide health
- 36 benefits. There is no threshold that must be exceeded before benefits begin to occur.
- 37 • Greater benefits can be achieved by reducing sedentary behavior, increasing
- 38 moderate-intensity physical activity, or a combination of both.
- 39 • For any given increase in moderate-to-vigorous physical activity, the relative gain
- 40 in benefits is greater for insufficiently active people than for people who are already
- 41 meeting the key guidelines.

1 **For people who are active**, that is, people who already meet the key guidelines (150 to
2 300 minutes a week of moderate-intensity physical activity for adults):

- 3 • Although those within the target range already have substantial benefits from their
4 current volume of physical activity, more benefits can be gained by doing additional
5 moderate-to-vigorous physical activity or reducing sedentary behavior.

6
7 **For people who are highly active**, that is, people who do more than the equivalent of 300
8 minutes a week of moderate-intensity physical activity:

- 9 • These people should maintain or increase their activity level by doing a variety of
10 activities.

11 **INTERVENTIONS**

12 Although the correlation among healthful diet, physical activity, and the incidence of
13 cardiovascular disease is strong, existing evidence indicates that the health benefit of
14 initiating behavioral counseling in the primary care setting to promote a healthful diet and
15 physical activity is minimal (Moyer, 2012). The USPSTF (2022) recommends selectively
16 offering or providing behavioral counseling interventions to patients based on clinician
17 professional judgment and patient preferences. Behavioral counseling interventions may
18 promote physical activity, healthy diet, reducing sedentary time, or some combination
19 thereof. Common dietary counseling advice promotes increased consumption of fruits,
20 vegetables, and fiber; reduced consumption of saturated fats, sodium, and sugar-sweetened
21 beverages; or both. Patient-tailored approaches to enhance skills with reading food labels,
22 preparing healthy meals, and recognizing appropriate caloric intake and portion size are
23 often used. Physical activity counseling often encourages patients to gradually increase
24 aerobic activity (walking is often emphasized) to achieve at least 150 minutes (2 hours and
25 30 minutes) per week of equivalent moderate-intensity activity. Reducing sedentary time
26 aims to limit time spent engaging in low-energy behaviors while awake such as sitting or
27 reclining while watching television or using a computer.

28
29
30 Primary care clinicians can deliver in-person behavioral counseling interventions or refer
31 patients to other settings. Interventions can be delivered individually, in a group, or both,
32 with or without follow-up (telephone calls or emails), or delivered remotely through a
33 combination of print materials, telephone calls, technology-based activities, or some
34 combination thereof. Typical counseling techniques include behavioral change techniques
35 such as goal setting, problem solving, and self-monitoring; approaches including
36 motivational interviewing principles or portions of the “5 A’s” model (assess, advise,
37 agree, assist, and arrange) are common. In determining whether behavioral counseling
38 interventions are appropriate, patients and clinicians should consider the following.

- 39 • Persons who are interested and ready to make behavioral changes may be most
40 likely to benefit from behavioral counseling.
- 41 • Higher-intensity counseling interventions may vary in availability and feasibility in
42 clinical settings.

- 1 • Adoption of healthy behavior advice may be increased by tailoring behavioral
2 counseling to consider patient motivations and goals, activity level and ability,
3 circumstances, preferences, and overall health status, as well as availability of
4 healthy eating establishments, grocery stores, parks, sidewalks, bicycle trails,
5 safe/pleasant walking paths close to home or workplace; traffic; public
6 transportation; crime; and pollution levels.

7
8 Encouraging patients to increase their baseline activity is sensible for several reasons:

- 9 • Increasing baseline activity burns calories, which can help in maintaining a healthy
10 body weight.
11 • Some baseline activities are weight-bearing and may improve bone health.
12 • Short episodes of activity are appropriate for people who were inactive and have
13 started to gradually increase their level of activity, and for older adults whose
14 activity may be limited by chronic conditions.
15 • There are reasons other than health to encourage more baseline activity. For
16 example, walking short distances instead of driving can help reduce traffic
17 congestion and the resulting air pollution.
18 • Encouraging baseline activities helps build a culture where physical activity in
19 general is the social norm.
20

21 The USPSTF found sufficient evidence that behavioral counseling interventions for a
22 healthy diet, physical activity, or both were associated with modest increases in physical
23 activity levels and some improvements in dietary health behaviors. On average,
24 participants of physical activity interventions (87 trials) increased physical activity by
25 approximately 33 minutes per week and had higher odds of meeting physical activity
26 recommendations after 6 to 12 months compared with participants in the control group.
27 Participants of healthy diet interventions (45 trials) increased fruit and vegetable intake and
28 fiber intake and decreased saturated fat intake compared with participants in the control
29 group. The USPSTF found little evidence of the effectiveness of sedentary behavior
30 interventions. Overall, sedentary behavior interventions did not demonstrate statistically
31 significant differences in sedentary behaviors in participants compared with control
32 groups.
33

34 The USPSTF found sufficient evidence that behavioral counseling interventions for a
35 healthy diet, physical activity, or both were associated with lower blood pressure, low-
36 density lipoprotein cholesterol (LDL-C), and adiposity measures (BMI, weight, and waist
37 circumference) after 6 to 12 months. Diet and physical activity interventions were
38 associated with lower systolic blood pressure, diastolic blood pressure, LDL-C level, and
39 adiposity-related outcomes such as weight, BMI, and waist circumference. Generally,
40 high-intensity interventions (>360 minutes) were associated with greater changes in
41 intermediate outcomes, specifically lower LDL-C and adiposity measures.

1 Because of the extremely important health benefits of physical activity, many Federal
 2 agencies and organizations recommend that all healthcare providers counsel individuals
 3 about physical activity. These include Healthy People 2030 (USDHHS), the CDC, and the
 4 American Academy of Family Physicians.

5
 6 **SCREENING RECOMMENDATIONS**
 7

USPSTF Recommendation

Grade C Recommendation: *For adults 18 years or older without known cardiovascular disease risk factors, the USPSTF recommends that clinicians individualize the decision to offer or refer adults without cardiovascular disease risk factors to behavioral counseling interventions to promote a healthy diet and physical activity.*

Definitions:

Grade B Recommendation: The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.

Grade C Recommendation: The USPSTF recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small.

Grade D Recommendation: The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits.

8
 9 This recommendation applies to adults 18 years or older without known CVD risk factors,
 10 which include hypertension or elevated blood pressure, dyslipidemia, impaired fasting
 11 glucose, or glucose tolerance, or mixed or multiple risk factors such as metabolic syndrome
 12 or an estimated 10-year CVD risk of 7.5% or greater. While obesity is a risk factor for
 13 CVD, a separate recommendation statement addresses individuals with a body mass index
 14 (BMI) of 30 or greater (calculated as weight in kilograms divided by the square of height
 15 in meters). Behavioral interventions to reduce CVD risk in adults with known modifiable
 16 risk factors (i.e., hypertension or dyslipidemia) and weight management interventions are
 17 addressed in separate USPSTF recommendations.

18
 19 A comprehensive review of the USPSTF rating process can be found in the *Preventive*
 20 *Care Services (CPG 140 – S)* clinical practice guidelines or at the USPSTF website
 21 <http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions>.

1 **Clinical Management Interventions**

2 The American Heart Association recommends that practitioners use cognitive-behavioral
3 counseling interventions to promote healthful diet and physical activity that combine 2 or
4 more of the following strategies (Artinian et al., 2010):

6 *Cognitive-behavioral strategies for promoting behavior change*

- 7 • Design interventions to target dietary and physical activity behaviors with specific,
8 proximal goal setting.
- 9 • Provide feedback on progress toward goals.
- 10 • Provide strategies for self-monitoring.
- 11 • Establish a plan for frequency and duration of follow-up contacts (e.g., in-person,
12 oral, written, electronic) in accordance with individual needs to assess and reinforce
13 progress toward goal achievement.
- 14 • Utilize motivational interviewing strategies, particularly when an individual is
15 resistant or ambivalent about dietary and physical activity behavior change.
- 16 • Incorporate strategies to build self-efficacy into the intervention.
- 17 • Use a combination of two of the above strategies (e.g., goal setting, feedback, self-
18 monitoring, follow-up, motivational interviewing, and self-efficacy) in an
19 intervention.

20
21 Additionally, health care practitioners may offer healthy diet and physical activity
22 interventions by referring the patient to community-based organizations. Strong links
23 between the practitioner and community-based resources may improve the delivery of
24 these services.

26 **PRACTITIONER SCOPE AND TRAINING**

27 Practitioners should practice only in the areas in which they are competent based on their
28 education, training, and experience. Levels of education, experience, and proficiency may
29 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
30 to determine where they have the knowledge and skills necessary to perform such services
31 and whether the services are within their scope of practice.

32
33 It is best practice for the practitioner to appropriately render services to a patient only if
34 they are trained, equally skilled, and adequately competent to deliver a service compared
35 to others trained to perform the same procedure. If the service would be most competently
36 delivered by another health care practitioner who has more skill and training, it would be
37 best practice to refer the patient to the more expert practitioner.

38
39 Best practice can be defined as a clinical, scientific, or professional technique, method, or
40 process that is typically evidence-based and consensus driven and is recognized by a
41 majority of professionals in a particular field as more effective at delivering a particular

1 outcome than any other practice (Joint Commission International Accreditation Standards
2 for Hospitals, 2020).

3
4 Depending on the practitioner’s scope of practice, training, and experience, a patient’s
5 condition and/or symptoms during examination or the course of treatment may indicate the
6 need for referral to another practitioner or even emergency care. In such cases it is prudent
7 for the practitioner to refer the patient for appropriate co-management (e.g., to their primary
8 care physician) or if immediate emergency care is warranted, to contact 911 as appropriate.
9 See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice guideline for
10 information.

11 **Practitioner Resources**

- 12 • Centers for Disease Controls & Prevention (CDC) Physical Activity Resources
13 <http://www.cdc.gov/physicalactivity/>
- 14 • US Dept of Health & Human Services – Healthy People 2030
15 [https://health.gov/healthypeople/objectives-and-data/browse-objectives/physical-](https://health.gov/healthypeople/objectives-and-data/browse-objectives/physical-activity)
16 [activity](https://health.gov/healthypeople/objectives-and-data/browse-objectives/physical-activity)
17

18 **Member Resources**

- 19 • US Department of Health and Human Services: Physical Activity Guidelines
20 <http://www.health.gov/paguidelines/>
- 21 • American Heart Association’s Getting Active: [https://www.heart.org/en/healthy-](https://www.heart.org/en/healthy-living/fitness/getting-active)
22 [living/fitness/getting-active](https://www.heart.org/en/healthy-living/fitness/getting-active)
23
- 24 • NIH’s Physical Activity Tools and Resources:
25 [http://www.nhlbi.nih.gov/health/public/heart/obesity/wecan/tools-](http://www.nhlbi.nih.gov/health/public/heart/obesity/wecan/tools-resources/physical-activity.htm)
26 [resources/physical-activity.htm](http://www.nhlbi.nih.gov/health/public/heart/obesity/wecan/tools-resources/physical-activity.htm)
- 27 • President’s Council on Fitness, Sports & Nutrition [http://www.fitness.gov/be-](http://www.fitness.gov/be-active/)
28 [active/](http://www.fitness.gov/be-active/)
- 29 • US Health & Human Services – Girls Health: Fitness
30 <http://www.girlshealth.gov/fitness/>
- 31 • National Institute on Aging (NIA)’s Exercise & Physical Activity: Your Everyday
32 Guide [https://www.nia.nih.gov/health/publication/exercise-physical-](https://www.nia.nih.gov/health/publication/exercise-physical-activity/introduction)
33 [activity/introduction](https://www.nia.nih.gov/health/publication/exercise-physical-activity/introduction)
- 34 • NIH’s Your Guide to Physical Activity and Your Heart
35 http://www.nhlbi.nih.gov/health/public/heart/obesity/phy_active.pdf
36

37 ***References***

38 Albright CL, Cohen S, Gibbons L, et al. Incorporating physical activity advice into primary
39 care: physician-delivered advice within the activity counseling trial. *Am J Prev Med.*
40 2000;18(3):225-234. doi:10.1016/s0749-3797(99)00155-5

- 1 Artinian NT, Fletcher GF, Mozaffarian D, et al. Interventions to promote physical activity
2 and dietary lifestyle changes for cardiovascular risk factor reduction in adults: a
3 scientific statement from the American Heart Association. *Circulation*.
4 2010;122(4):406-441. doi:10.1161/CIR.0b013e3181e8edf1
- 5
6 Centers for Disease Control & Prevention. Exercise or Physical Activity. Retrieved on
7 March 26, 2024 from <https://www.cdc.gov/nchs/fastats/exercise.htm>
- 8
9 Centers for Disease Control & Prevention. The Benefits of Physical Activity. (2023).
10 Retrieved on March 26, 2024 from
11 http://www.cdc.gov/physicalactivity/everyone/health/index.html?s_cid=cs_284
- 12
13 Centers for Disease Control & Prevention. Physical Activity: Why It Matters. Retrieved on
14 March 26, 2024 from <https://www.cdc.gov/physicalactivity/about-physical-activity/why-it-matters.html>
- 15
16 Danaei G, Ding EL, Mozaffarian D, et al. The preventable causes of death in the United
17 States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors
18 [published correction appears in *PLoS Med*. 2011 Jan;8(1). doi:
19 10.1371/annotation/0ef47acd-9dcc-4296-a897-872d182cde57]. *PLoS Med*.
20 2009;6(4):e1000058. doi:10.1371/journal.pmed.1000058
- 21
22
23 Eden KB, Orleans CT, Mulrow CD, Pender NJ, Teutsch SM. Does counseling by clinicians
24 improve physical activity? A summary of the evidence for the U.S. Preventive Services
25 Task Force. *Ann Intern Med*. 2002;137(3):208-215. doi:10.7326/0003-4819-137-3-
26 200208060-00015
- 27
28 Joint Commission International. (2020). Joint Commission International Accreditation
29 Standards for Hospitals (7th ed.): Joint Commission Resources
- 30
31 Johnson NB, Hayes LD, Brown K, Hoo EC, Ethier KA; Centers for Disease Control and
32 Prevention (CDC). CDC National Health Report: leading causes of morbidity and
33 mortality and associated behavioral risk and protective factors--United States, 2005-
34 2013. *MMWR Suppl*. 2014;63(4):3-27
- 35
36 Lavie CJ, Ozemek C, Carbone S, Katzmarzyk PT, Blair SN. Sedentary Behavior, Exercise,
37 and Cardiovascular Health. *Circ Res*. 2019;124(5):799-815.
38 doi:10.1161/CIRCRESAHA.118.312669
- 39
40 Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United
41 States, 2000 [published correction appears in *JAMA*. 2005 Jan 19;293(3):298]

1 [published correction appears in JAMA. 2005 Jan 19;293(3):293-4]. JAMA.
2 2004;291(10):1238-1245. doi:10.1001/jama.291.10.1238

3
4 Moyer VA; U.S. Preventive Services Task Force. Behavioral counseling interventions to
5 promote a healthful diet and physical activity for cardiovascular disease prevention in
6 adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern*
7 *Med.* 2012;157(5):367-371. doi:10.7326/0003-4819-157-5-201209040-00486

8
9 Office of Disease Prevention and Health Promotion (ODPHP). Physical Activity
10 Guidelines for Americans. Second Edition. Retrieved on March 26, 2024 from:
11 <https://health.gov/paguidelines/second-edition/>

12
13 Patnode CD, Redmond N, Iacocca MO, Henninger M. Behavioral Counseling
14 Interventions to Promote a Healthy Diet and Physical Activity for Cardiovascular
15 Disease Prevention in Adults Without Known Cardiovascular Disease Risk Factors:
16 Updated Evidence Report and Systematic Review for the US Preventive Services Task
17 Force. *JAMA.* 2022;328(4):375-388. doi:10.1001/jama.2022.7408

18
19 U.S. Department of Health and Human Services. Healthy People 2030. Retrieved March
20 26, 2024 from: <https://health.gov/healthypeople>

21
22 U.S. Department of Health and Human Services. Healthy People 2030: Physical Activity.
23 Retrieved March 26, 2024 from [https://health.gov/healthypeople/objectives-and-](https://health.gov/healthypeople/objectives-and-data/browse-objectives/physical-activity)
24 [data/browse-objectives/physical-activity](https://health.gov/healthypeople/objectives-and-data/browse-objectives/physical-activity)

25
26 USPSTF Recommendation Statement: Healthy Diet and Physical Activity for
27 Cardiovascular Disease Prevention in Adults Without Cardiovascular Disease Risk
28 Factors: Behavioral Counseling Interventions. (2022). Update in Progress. Retrieved
29 from March 26, 2024 from
30 [https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/healthy-](https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/healthy-lifestyle-and-physical-activity-for-cvd-prevention-adults-without-known-risk-factors-behavioral-counseling)
31 [lifestyle-and-physical-activity-for-cvd-prevention-adults-without-known-risk-factors-](https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/healthy-lifestyle-and-physical-activity-for-cvd-prevention-adults-without-known-risk-factors-behavioral-counseling)
32 [behavioral-counseling.](https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/healthy-lifestyle-and-physical-activity-for-cvd-prevention-adults-without-known-risk-factors-behavioral-counseling)

33
34 Zhang Y, Liu X. Effects of physical activity and sedentary behaviors on cardiovascular
35 disease and the risk of all-cause mortality in overweight or obese middle-aged and older
36 adults. *Front Public Health.* 2024 Feb 12;12:1302783. doi:
37 10.3389/fpubh.2024.1302783. PMID: 38410660; PMCID: PMC10894908