

1 **Clinical Practice Guideline: Ostectomy for Tailor’s Bunion**

2

3 **Date of Implementation: June 18, 2015**

4

5 **Product: Specialty**

6

7

8 **GUIDELINES**

9 American Specialty Health – Specialty (ASH) considers services consisting of CPT® Code
10 28110 to be medically necessary for the treatment of Tailor’s bunion **upon meeting ALL**
11 **of the following criteria:**

- 12 1. When supported by a diagnosis of other acquired deformities of foot (Tailor’s
13 bunion) (M21.621- M21.622)
- 14 2. Symptomatic bunion (5th metatarsal) **indicated by 1 or more of the following:**
- 15 • Ulceration at fifth metatarsophalangeal joint
 - 16 • Difficulty walking because of pain at fifth metatarsophalangeal joint
 - 17 • Inability to accommodate or modify footwear to control pain
- 18 3. Failure of **at least 2 of the following** non-operative treatments
- 19 • Padding
 - 20 • Orthotics
 - 21 • Shoe modification
 - 22 • Activity modification
 - 23 • Injection

24

25 **CPT® Codes and Descriptions**

| CPT® Code | CPT® Code Description |
|------------------|--|
| 28110 | Ostectomy, partial excision, fifth metatarsal head (bunionette) (separate procedure) |

26

27 **BACKGROUND**

28 Tailor’s bunion (also called bunionette) involves deformity of the fifth
29 metatarsophalangeal (MTP) joint, typically involves deformity with lateral prominence of
30 the fifth metatarsal head. Structural causes of this deformity may include a prominent
31 lateral condyle, a plantarflexed fifth metatarsal, a splay foot deformity, lateral bowing of
32 the fifth metatarsal, a combination of these deformities, or hypertrophy of the soft tissues
33 over the lateral aspect of the metatarsal head.

34

35 The patient with a tailor’s bunion may or may not have pain related to the deformity. The
36 most common presentation of bunionette deformity is pain associated with lateral pressure
37 with footwear. There also may be a history of localized swelling and/or callus formation.

1 The clinical examination of a patient with a tailor’s bunion will reveal a lateral or plantar-
2 lateral prominence of the fifth metatarsal head. Tenderness on palpation of the lateral
3 and/or plantar-lateral fifth metatarsal head may be associated with an overlying adventitial
4 bursa or hyperkeratotic lesion.

5
6 Radiographic findings may be helpful in the assessment of the exact nature of the deformity
7 and contributory structural pathology. Radiographic evaluation includes anteroposterior,
8 oblique, and lateral views of the weight-bearing foot. Common measurements obtained
9 using the AP view include metatarsal head width, IMA, and fifth metatarsophalangeal
10 angle (MTPA). Average normal IMA has historically been considered to be 6.5° , with
11 angles $>8^\circ$ defined as abnormal. Fifth MTPA is measured by the degree of divergence of
12 the fifth toe from the long axis of the metatarsal shaft. The MTPA measures on average
13 10.2° to 14° in normal feet but 16° in feet with symptomatic bunionette deformities (Shi et
14 al., 2018).

15
16 Conservative care is the first line of treatment for foot and toe deformity. Conservative
17 treatment for tailor’s bunion may include wide shoes, custom accommodative orthotics,
18 nonsteroidal anti-inflammatory medications, and barrier pads to alleviate pain. Skincare
19 with callous shaving of keratosis also can temporarily relieve pressure. The use of localized
20 injections to address concomitant metatarsophalangeal joint (MTPJ) synovitis has been
21 studied, with moderate, short-term, and sustained improvement in less than one-third of the
22 patients at 2-year follow-up (Shi et al., 2018; Morawe & Schmieschek, 2018).

23
24 Surgical treatment is indicated for patients experiencing pain due to bunionette deformity
25 who have failed nonsurgical care and patients who are not candidates for nonsurgical care.
26 The goal of surgical treatment is to decrease the prominence of the fifth metatarsal laterally.
27 Selection of the surgical procedure is based on the physical evaluation and radiographic
28 assessment. Surgical correction to alleviate the pain at the bone prominence varies from
29 exostectomy to differing types of osteotomies. Resection of the fifth metatarsal head, which
30 entails removal of a portion of the bone, for treatment of tailor’s bunion is generally
31 indicated for salvage conditions or in the presence of unreconstructable deformities (Ajis
32 et al., 2005; Shi et al., 2018).

33
34 Excision of the fifth metatarsal head, resection of the distal half of the metatarsal, and fifth
35 ray resection have all been used to treat a bunionette deformity but are not appropriate in
36 the initial treatment of symptomatic conditions. Therefore, these procedures are used as a
37 salvage procedures for infection, ulceration, or severe deformity. Situations in which these
38 procedures may be appropriate include severe osteopenia, extensive degenerative joint
39 changes, chronic ulceration, previous failed surgery, or poor medical health (DiDomenico
40 et al., 2013).

1 PRACTITIONER SCOPE AND TRAINING

2 Practitioners should practice only in the areas in which they are competent based on their
3 education, training, and experience. Levels of education, experience, and proficiency may
4 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
5 to determine where they have the knowledge and skills necessary to perform such services
6 and whether the services are within their scope of practice.

7
8 It is best practice for the practitioner to appropriately render services to a member only if
9 they are trained, equally skilled, and adequately competent to deliver a service compared
10 to others trained to perform the same procedure. If the service would be most competently
11 delivered by another health care practitioner who has more skill and training, it would be
12 best practice to refer the member to the more expert practitioner.

13
14 Best practice can be defined as a clinical, scientific, or professional technique, method, or
15 process that is typically evidence-based and consensus driven and is recognized by a
16 majority of professionals in a particular field as more effective at delivering a particular
17 outcome than any other practice (Joint Commission International Accreditation Standards
18 for Hospitals, 2020).

19
20 Depending on the practitioner’s scope of practice, training, and experience, a member’s
21 condition and/or symptoms during examination or the course of treatment may indicate the
22 need for referral to another practitioner or even emergency care. In such cases it is prudent
23 for the practitioner to refer the member for appropriate co-management (e.g., to their
24 primary care physician) or if immediate emergency care is warranted, to contact 911 as
25 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice
26 guideline for information.

27 28 **References**

29 Ajis, A., Koti, M., & Maffulli, N. (2005). Tailor's bunion: a review. *The Journal of foot*
30 *and ankle surgery*, 44(3), 236–245. <https://doi.org/10.1053/j.jfas.2005.02.005>

31
32 American College of Foot and Ankle Surgeons (ACFAS) Cosmetic Surgery Position
33 Statement (2020). Retrieved February 12, 2024 from: [https://www.acfas.org/policy-](https://www.acfas.org/policy-advocacy/policy-position-statements/acfas-position-statement-on-cosmetic-surgery)
34 [advocacy/policy-position-statements/acfas-position-statement-on-cosmetic-surgery](https://www.acfas.org/policy-advocacy/policy-position-statements/acfas-position-statement-on-cosmetic-surgery)

35
36 American Medical Association. (current year). *Current Procedural Terminology (CPT)*
37 *Current year (rev. ed.)*. Chicago: AMA

38
39 Cohen, B. E., & Nicholson, C. W. (2007). Bunionette deformity. *The Journal of the*
40 *American Academy of Orthopaedic Surgeons*, 15(5), 300–307.
41 <https://doi.org/10.5435/00124635-200705000-00008>

- 1 DiDomenico, L., Baze, E., & Gatalyak, N. (2013). Revisiting the tailor's bunion and
2 adductovarus deformity of the fifth digit. *Clinics in Podiatric Medicine and Surgery*,
3 30(3), 397-422. doi: 10.1016/j.cpm.2013.04.004
4
- 5 Gould, J. S., & Florence, M. N. (2013). *The Handbook of Foot and Ankle Surgery: An*
6 *Intellectual Approach to Complex Problems*: Jaypee Brothers, Medical Publishers
7
- 8 Joint Commission International. (2020). Joint Commission International Accreditation
9 Standards for Hospitals (7th ed.): Joint Commission Resources
10
- 11 Morawe, G. A., & Schmieschek, M. H. T. (2018). Minimally invasive bunionette
12 correction. Minimal-invasive Bunionette-Korrektur. *Operative Orthopädie und*
13 *Traumatologie*, 30(3), 184–194. <https://doi.org/10.1007/s00064-018-0542-z>
14
- 15 Roukis, T. S. (2005). The tailor's bunionette deformity: a field guide to surgical correction.
16 *Clinics in Podiatric Medicine and Surgery*, 22(2), 223-vi. doi:
17 10.1016/j.cpm.2004.10.004
18
- 19 Shi, G. G., Humayun, A., Whalen, J. L., & Kitaoka, H. B. (2018). Management of
20 Bunionette Deformity. *The Journal of the American Academy of Orthopaedic*
21 *Surgeons*, 26(19), e396–e404. <https://doi.org/10.5435/JAAOS-D-17-00345>
22
- 23 Thomas, J. L., Blich, E. L.^{4th}, Chaney, D. M., Dinucci, K. A., Eickmeier, K., Rubin, L.
24 G., & Vanore, J. V. (2009). Diagnosis and Treatment of Forefoot Disorders. Section 4.
25 Tailor's Bunion. *The Journal of Foot and Ankle Surgery*, 48(2), 257-263