

1 **Clinical Practice Guideline:** **J Codes - Podiatry**

2

3 **Date of Implementation:** **November 19, 2015**

4

5 **Product:** **Specialty**

6

7

8 **GUIDELINES**

9

10

11

12

13

14

15

16

17

18

19

20

21

A. Injections, as listed in the table below, will be approved provided they are reasonable, medically necessary, and related to the diagnosis and treatment of an illness or injury or are for purposes of immunization for which medical necessity is established. If the procedure has established frequency of injections within its guideline, billing of appropriate and related J code will follow this guideline as well.

When excessive injections appear, representing a departure from accepted standards of medical practice, the entire charge for injection given in excess of these standards will be excluded. If an injection is determined to fall outside of what is medically reasonable or necessary, the entire charge (i.e., for both the drug and its administration) will be excluded from payment.

HCPCS Codes and Descriptions

HCPCS Code	HCPCS Code Description	Drug Type
J0120	Injection, tetracycline, up to 250 mg	Antibiotic
J0171	Injection, adrenalin, epinephrine, 0.1 mg	Alpha- and Beta-adrenergic Agonists (sympathomimetic agents)
J0290	Injection, ampicillin sodium, 500 mg	Antibiotic
J0295	Injection, ampicillin sodium/sulbactam sodium, per 1.5 g	Antibiotic
J0456	Injection, azithromycin, 500 mg	Antibiotic
J0461	Injection, atropine sulfate, 0.01 mg	Anticholinergic
J0475	Injection, baclofen, 10 mg	Muscle relaxant and antispastic
J0558	Injection, penicillin G benzathine and penicillin G procaine, 100,000 units	Antibiotic

HCPCS Code	HCPCS Code Description	Drug Type
J0561	Injection, penicillin G benzathine, 100,000 units	Antibiotic
J0670	Injection, mepivacaine HCl, per 10 ml	Anesthetic
J0688	Injection, cefazolin sodium (Hikma), not therapeutically equivalent to J0690, 500 mg	Antibiotic
J0689	Injection, cefazolin sodium (Baxter), not therapeutically equivalent to J0690, 500 mg	Antibiotic
J0690	Injection, cefazolin sodium, 500 mg	Antibiotic
J0696	Injection, ceftriaxone sodium, per 250 mg	Antibiotic
J0702	Injection, betamethasone acetate 3 mg and betamethasone sodium phosphate 3 mg	Corticosteroid
J1020	Injection, methylprednisolone acetate, 20 mg	Corticosteroid
J1030	Injection, methylprednisolone acetate, 40 mg	Corticosteroid
J1040	Injection, methylprednisolone acetate, 80 mg	Corticosteroid
J1094	Injection, dexamethasone acetate, 1 mg	Corticosteroid
J1100	Injection, dexamethasone sodium phosphate, 1 mg	Corticosteroid
J1170	Injection, hydromorphone, up to 4 mg	Narcotic (Opioid) Analgesic
J1885	Injection, ketorolac tromethamine, per 15 mg	NSAID
J1940	Injection, furosemide, up to 20 mg	Diuretic
J2001	Injection, lidocaine HCl for intravenous infusion, 10 mg	Anesthetic
J2250	Injection, midazolam HCl, per 1 mg	Anesthetic
J2920	Injection, methylprednisolone sodium succinate, up to 40 mg	Corticosteroid
J3010	Injection, fentanyl citrate, 0.1 mg	Narcotic (Opioid) Analgesic
J3301	Injection, triamcinolone acetonide, not otherwise specified, 10 mg	Corticosteroid

HCPCS Code	HCPCS Code Description	Drug Type
J3302	Injection, triamcinolone diacetate, per 5 mg	Corticosteroid
J3303	Injection, triamcinolone hexacetonide, per 5 mg	Corticosteroid
J9040	Injection, bleomycin sulfate, 15 units	Antitumor Antibiotic and Antiviral

1
2
3
4
5

B. Dexamethasone, as listed in the table below, will be approved provided it is reasonable, medically necessary, and related to the diagnosis and treatment of an illness or injury for which medical necessity is established.

HCPCS Code	HCPCS Code Description	Drug Type
J8540	Dexamethasone, oral, 0.25 mg	Corticosteroid

6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

BACKGROUND

Antibiotics

Skin, soft tissue and bone infections present in varying ways with regards to presentation, etiology, and severity. These infections range from mild, such as pyoderma, to more severe infections that are life threatening (i.e., necrotizing fasciitis). The affected area will present with erythema, edema, warmth, pain and/or tenderness. Mild infections can impair function if in areas such as the feet. Comorbidities can also impact the severity of infections and impact the plan of care. Oral antibiotics are typically the treatment of choice in mild cases, while injections of antibiotics are considered for more severe cases. Selection of type of antibiotic is determined by the type of microbe present. Risk factors may influence the etiology, course of disease and response to treatment. Patient-related risk factors include critical illness, elderly age, compromised immune system, liver and kidney disease and vascular insufficiency. Other risk factors include mechanism of injury and exposure (Ki and Rotstein, 2008).

Anti-inflammatories (NSAID and corticosteroid)

The ankle and foot are sites for many types of injuries and inflammatory conditions that respond well to diagnostic and therapeutic injections. Examples of areas include plantar fascia, ankle joint, tarsal tunnel, interdigital space, and the first metatarsophalangeal (MTP) joint. Separate clinical policy guidelines address medical necessity of procedures for conditions affecting these areas. Any restrictions on number of injections will also apply to the NSAID or corticosteroid J codes listed in the above table. Common complaints for inflammatory conditions of the foot and ankle include pain and disability with limitation

29

1 of motion, tenderness, warmth, swelling, crepitus and/or deformity. Gait dysfunction is
2 often a result of these symptoms as well (Tallia and Cardone, 2003).

3 Anesthetics

4 Injections of local anesthetics are used for diagnostic purposes and surgical procedures.
5 Local or regional anesthetics render part of the body insensitive to pain without affecting
6 a person's wakefulness. They are also reversible and will wear off at some point in time
7 post procedurally, with the time depending upon the type and administration of anesthetic.
8 When used locally or regionally, an injection or series of injections occur at the toe, ankle,
9 or knee. Local anesthetics are less risky for the patient than general anesthesia. Practitioners
10 must ensure that patients receiving these injections are medically fit to undergo the
11 procedure, have home settings that allow them to return to home the same day, and be
12 physically able and willing to comply with the post-procedure regimen. Examples of
13 anesthetic injections include lidocaine injection with a corticosteroid injection or a
14 peripheral nerve block (hallux or ankle) for treatment of an ingrown toenail or for midfoot
15 surgery. Popliteal blocks may also be used for foot and ankle surgeries.

17 Miscellaneous

18 Epinephrine injection is used along with emergency medical treatment to treat life-
19 threatening allergic reactions, including anaphylaxis, caused by insect bites or stings,
20 foods, medications, latex, and other causes. Epinephrine is in a class of medications called
21 alpha- and beta-adrenergic agonists (sympathomimetic agents). It works by relaxing the
22 muscles in the airways and constricting the blood vessels.

23
24 Gablofen (baclofen injection) is a muscle relaxant and antispastic (gamma-aminobutyric
25 acid [GABA] ergic agonist). It is indicated in the management of severe spasticity of
26 cerebral or spinal origin in adult and pediatric patients ages 4 years and above. Baclofen
27 injection is used by patients who do not respond to or who cannot tolerate the side effects
28 of baclofen taken by mouth. Patients should first respond to a screening dose of intrathecal
29 baclofen prior to consideration for long term infusion via an implantable pump. If spasticity
30 is due to traumatic brain injury, consideration of Gablofen should be delayed for at least
31 one year. Gablofen should not be used for intravenous, intramuscular, subcutaneous, or
32 epidural administration.

33
34 Atropine is an anticholinergic. It works by blocking the effects of a chemical in the body
35 (acetylcholine) that acts in the nervous system, stomach, intestines, certain glands (e.g.,
36 salivary gland), urinary tract, and other tissues. It is used to decrease the production of saliva
37 and secretions of the airway prior to surgery. It is also used to treat spasms in the stomach,
38 intestines, and other organs. It may also be used to counteract the effects of certain other
39 medicines or for other conditions as determined by a doctor.
40

1 Hydromorphone and fentanyl citrate injections are used to relieve severe pain. They both
 2 are in a class of medications called narcotic analgesics. These medications work by
 3 changing the way the brain and nervous system respond to pain. They can be injected into
 4 muscle or veins. These medications may be used for in-office procedures that are invasive
 5 in nature and given their mechanism and potency; should be monitored closely by
 6 physicians.

8 **PRACTITIONER SCOPE AND TRAINING**

9 Practitioners should practice only in the areas in which they are competent based on their
 10 education, training, and experience. Levels of education, experience, and proficiency may
 11 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
 12 to determine where they have the knowledge and skills necessary to perform such services
 13 and whether the services are within their scope of practice.

14
 15 It is best practice for the practitioner to appropriately render services to a member only if
 16 they are trained, equally skilled, and adequately competent to deliver a service compared
 17 to others trained to perform the same procedure. If the service would be most competently
 18 delivered by another health care practitioner who has more skill and training, it would be
 19 best practice to refer the member to the more expert practitioner.

20
 21 Best practice can be defined as a clinical, scientific, or professional technique, method, or
 22 process that is typically evidence-based and consensus driven and is recognized by a
 23 majority of professionals in a particular field as more effective at delivering a particular
 24 outcome than any other practice (Joint Commission International Accreditation Standards
 25 for Hospitals, 2020).

26
 27 Depending on the practitioner’s scope of practice, training, and experience, a member’s
 28 condition and/or symptoms during examination or the course of treatment may indicate the
 29 need for referral to another practitioner or even emergency care. In such cases it is prudent
 30 for the practitioner to refer the member for appropriate co-management (e.g., to their
 31 primary care physician) or if immediate emergency care is warranted, to contact 911 as
 32 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practice
 33 guideline for information.

34 **References**

35
 36 American College of Foot and Ankle Surgeons (ACFAS) Cosmetic Surgery Position
 37 Statement (2020). Retrieved on February 12, 2024 from: <https://www.acfas.org/policy-advocacy/policy-position-statements/acfas-position-statement-on-cosmetic-surgery>

38
 39
 40 American Medical Association (current year). HCPCS Level II. American Medical
 41 Association

- 1 Joint Commission International. (2020). Joint Commission International Accreditation
2 Standards for Hospitals (7th ed.): Joint Commission Resources
3
4 Ki, V., & Rotstein, C. (2008). Bacterial skin and soft tissue infections in adults: A review
5 of their epidemiology, pathogenesis, diagnosis, treatment and site of care. *The*
6 *Canadian Journal of Infectious Diseases & Medical Microbiology*, 19(2), 173-184
7
8 Tallia, A. F., & Cardone, D. A. (2003). Diagnostic and Therapeutic Injection of the Ankle
9 and Foot. *American Family Physician*, 68(7), 1356-1363