

1 **Clinical Practice Guideline:** **Mechanical Traction (Provided in a Clinic Setting)**

2

3 **Date of Implementation:** **June 16, 2016**

4

5 **Product:** **Specialty**

6

7

Related Policies:
 CPG 83: Axial Decompression Therapy
 CPG 121: Passive Physiotherapy (Therapeutic) Modalities
 CPG 135: Physical Therapy Medical Policy/Guidelines
 CPG 155: Occupational Therapy Medical Policy/Guidelines
 CPG 265: Home Traction Therapy

8

9

10

11

12

13 **Table of Contents**

14 GUIDELINES 1

15 Cervical Spine 1

16 Lumbar Spine 2

17 Thoracic Spine..... 2

18 GENERAL MEDICAL NECESSITY CRITERIA..... 2

19 DESCRIPTION/BACKGROUND 3

20 Contraindications and Precautions 4

21 EVIDENCE REVIEW..... 5

22 Cervical 5

23 Lumbar 8

24 PRACTITIONER SCOPE AND TRAINING12

25 *References*13

26

27 **GUIDELINES**

28 **Cervical Spine**

29

American Specialty Health – Specialty (ASH) considers use of cervical mechanical traction as medically necessary for patients who meet **ALL of the following** criteria:

30

31

- Failure of other evidence-based therapeutic procedures to significantly improve symptoms after 3 weeks.

32

33

- Only used in combination with other evidence-based treatments including therapeutic exercise. The therapeutic exercise(s) should not cause aggravation or peripheralization of symptoms.

34

35

- 1 • Cervical radiculopathy should be supported by the exam findings including
2 provocative testing such as positive shoulder abduction, positive upper limb tension
3 test A, and/or positive neck distraction test.

4
5 ASH considers cervical mechanical traction as unproven because there is insufficient
6 evidence for treatment of other conditions or when the above criteria are not met.

7
8 **Lumbar Spine**

9 ASH considers use of lumbar mechanical traction as medically necessary for patients
10 who meet **ALL of the following** criteria:

- 11 • Failure of other evidence-based therapeutic procedures to significantly improve
12 symptoms after 3 weeks.
- 13 • Patient has sciatica or signs of nerve root compression and either peripheralization
14 with extension movements or a positive crossed straight leg raise test.
- 15 • Only used in combination with other evidence-based treatments including
16 therapeutic exercise with extension movements. The therapeutic exercise(s) should
17 not cause aggravation or peripheralization of symptoms.

18
19 ASH considers lumbar mechanical traction as unproven because there is insufficient
20 evidence for treatment of other conditions or when the above criteria are not met. These
21 guidelines are NOT relevant to axial or spinal decompression therapy.

22
23 **Thoracic Spine**

24 ASH considers mechanical traction applied to the thoracic spine as unproven because
25 there is insufficient evidence for treatment of thoracic conditions or other spinal
26 conditions beyond those outlined in this guideline.

27
28 ASH considers mechanical traction for spinal conditions not specified in this guideline
29 as unproven due to a lack of supporting evidence.

30
31 **GENERAL MEDICAL NECESSITY CRITERIA**

- 32 • This therapy service is considered medically necessary when the judgment,
33 knowledge, and skills of a qualified practitioner of therapy services (as defined by
34 the scope of practice in each state) are necessary to safely and effectively furnish
35 this therapy service because of the complexity and sophistication of the plan of care
36 and the medical condition(s) of the patient, with the goal of improving an
37 impairment or functional limitation.
- 38 • The patient’s condition has the potential to improve or is improving in response to
39 this therapy service.
- 40 • The patient has not achieved maximum improvement from care.

- 1 • There is an expectation that the patient’s anticipated improvement is attainable in a
2 reasonable and predictable period of time and will result in a clinically significant
3 level of functional improvement through the use of this therapy service.
- 4 • The patient’s treatment is individualized and there is documentation outlining
5 quantifiable, attainable treatment goals with the use of this therapy service and the
6 patient’s overall plan of care.
- 7 • This therapy service is intended to improve, adapt or restore functions which have
8 been impaired or lost as a result of illness, injury, loss of a body part, or congenital
9 abnormality.
- 10 • The use of this therapy service (e.g., dosage, frequency) corresponds with the
11 current nature, status, and severity of the patient’s condition(s).
- 12 • The use of this therapy service is decreased as the patient displays improvement
13 and the plan of care transitions into other skilled treatment procedures that can
14 safely and effectively restore, adapt or improve the patient’s impaired function(s).
- 15 • The use of this therapy service is safe and effective for the patient’s condition, and
16 the patient is able to properly provide the necessary feedback for its safe
17 application.
- 18 • The use of this therapy service is not redundant with other therapy services used on
19 the same body part during the same session and is not duplicative with another
20 practitioner’s treatment plan.

21
22 **CPT® Codes and Descriptions**

CPT® Code	CPT® Code Description
97012	Application of a modality to 1 or more areas; traction, mechanical

23
24 **DESCRIPTION/BACKGROUND**

25 Traction is a therapeutic method used to relieve pain by stretching and separating the
26 vertebrae to help to relieve direct nerve pressure and stress on the vertebral discs. Cervical
27 traction is a common nonsurgical treatment for a herniated disc in the neck that relieves
28 pain by opening up the cervical foramen to reduce pressure on compressed nerve roots
29 exiting the spinal canal. Traction can either be applied manually or by spinal traction
30 devices. This guideline focuses on various mechanical traction devices that provide
31 continuous or intermittent forces to the spine. It has been proposed that cervical traction
32 results in an expansion of the intervertebral spaces, an increase joint mobility, and a
33 stretching muscles and ligaments adjacent to the vertebral bodies, which will improve
34 clinical outcomes in those with neck pain. After 2 minutes of sustained traction, the
35 intervertebral spaces begin to widen. Forces between 20 and 50 pounds are frequently used
36 to achieve intervertebral separation. Continuous or static traction can be applied in a steady
37 amount for specific time periods. Intermittent or cyclical traction involves traction being

1 applied and released multiple times during one treatment session. Duration of cervical
2 traction can range from a few minutes to 20 to 30 minutes, one to three times weekly.

3
4 Traction is used for treatment of low back pain (LBP) as well and it is provided in
5 combination with other treatment modalities, as is cervical traction. Lumbar traction uses
6 a harness (with Velcro strapping) that is put around the lower rib cage and around the iliac
7 crest. Duration and level of force exerted through this harness can be varied in a continuous
8 or intermittent mode. The exact mechanism through which traction might be effective is
9 still unclear. It has been suggested that spinal elongation, through decreasing lordosis and
10 increasing intervertebral space, inhibits pain (nociceptive) impulses, improves mobility,
11 decreases mechanical stress, reduces muscle spasm or spinal nerve root compression (due
12 to osteophytes), releases luxation of a disc or capsule from the zygapophyseal joint, and
13 releases adhesions around the zygapophyseal joint and the annulus fibrosus. So far, the
14 proposed mechanisms have not been supported by sufficient empirical information.

15 16 **Contraindications and Precautions**

17 Contraindications for mechanical traction include:

- 18 • Where motion is contraindicated
- 19 • Acute injury or inflammation
- 20 • Joint hypermobility or instability
- 21 • Peripheralization of symptoms with traction
- 22 • Uncontrolled hypertension
- 23 • Congenital spinal deformity
- 24 • Fractures
- 25 • Impaired mentation

26
27 Precautions for mechanical traction include:

- 28 • Structural diseases or conditions affecting the tissues in the area to be treated (e.g.,
29 tumor, infection, osteoporosis, RA, prolonged systemic steroid use, local radiation
30 therapy)
- 31 • When pressure of the belts may be hazardous (e.g., with pregnancy, hiatal hernia,
32 vascular compromise, osteoporosis)
- 33 • Cardiovascular disease
- 34 • Displaced annular fragment
- 35 • Medial disc protrusion
- 36 • Cord compression
- 37 • When severe pain fully resolves with traction
- 38 • Claustrophobia or other psychological aversion to traction
- 39 • Inability to tolerate prone or supine position
- 40 • Disorientation

1 Additional precautions for *cervical* traction:

- 2 • TMJ problems
- 3 • Dentures

4 **EVIDENCE REVIEW**

5 **Cervical**

7 Although traction has been used as a treatment for neck pain for decades, its effectiveness
 8 is unproven. Large, well designed, randomized controlled trials are needed that evaluate
 9 the effect of cervical traction as an adjunct treatment in both chronic and acute neck pain
 10 syndromes. Regardless, cervical traction remains a common treatment modality in the
 11 treatment of neck pain and radiculopathy. Borman et al. (2008) evaluated cervical traction
 12 for the treatment of chronic neck pain. Patients received standard care (hot pack, ultrasound
 13 and exercise) or cervical traction + standard care. The main outcome measures of the
 14 treatment were pain intensity by visual analog scale (VAS), disability by neck disability
 15 index (NDI), and quality of life assessed by Nottingham Health Profile (NHP) Both groups
 16 improved significantly in pain intensity and the scores of NDI and physical status of NHP
 17 at the end of the therapies ($p < 0.05$). Authors concluded that there was no specific effect of
 18 traction over standard physical therapy interventions in patients with chronic neck pain.
 19 Young et al. (2009) conducted a randomized controlled trial (RCT) on 81 patients with
 20 cervical radiculopathy to examine the effects of manual therapy and exercise, with or
 21 without the addition of cervical traction, on pain, function, and disability. Patients were
 22 randomly assigned to 1 of 2 groups: a group that received manual therapy, exercise, and
 23 intermittent cervical traction and a group that received manual therapy, exercise, and sham
 24 intermittent cervical traction. Patients were treated, on average, 2 times per week for an
 25 average of 4.2 weeks. Results demonstrated there were no significant differences between
 26 the groups for any of the primary or secondary outcome measures at 2 weeks or 4 weeks.
 27 Authors concluded that the addition of mechanical cervical traction to a multimodal
 28 treatment program of manual therapy and exercise adds no significant additional benefit to
 29 pain, function, or disability in patients with cervical radiculopathy.

31 Chiu et al. (2011) investigated the efficacy of intermittent cervical traction in the treatment
 32 of chronic neck pain over a 12-week period in an RCT of 79 patients The experimental
 33 group received intermittent cervical traction, and the control group received infrared
 34 irradiation alone twice a week over a period of six weeks. The authors concluded that there
 35 were no significant differences between the two groups. Graham et al. (2013) completed a
 36 systematic review on physical modalities for acute to chronic neck pain. Of 103 reviews
 37 eligible, 20 were included and 83 were excluded. Moderate evidence of benefit in the short
 38 term was noted for intermittent traction over placebo for chronic neck pain. No benefit was
 39 noted for continuous traction over placebo for whiplash associated disorder (WAD).
 40 Moderate evidence of no benefit for continuous traction was noted, as it was no better than
 41 placebo for acute whiplash associated disorder, chronic myofascial neck pain or subacute

1 to chronic neck pain. Improved design and long term follow up were suggested for future
2 research.

3
4 Raney et al. (2009) sought to determine a clinical prediction rule (CPR) to identify those
5 patients that were likely to benefit from cervical traction and exercise. Patients were
6 randomly selected into the following groups: exercise only, exercise with mechanical
7 traction, or exercise with over-door traction for patients with cervical radiculopathy. Sixty-
8 eight patients (38 female) were included in data analysis of which 30 had a successful
9 outcome. A CPR with five variables was identified: (1) patient reported peripheralization
10 with lower cervical spine (C4-7) mobility testing; (2) positive shoulder abduction test; (3)
11 age > or =55; (4) positive upper limb tension test A; and (5) positive neck distraction test.
12 Having at least three out of five predictors present resulted in a +LR equal to 4.81 (95% CI
13 = 2.17-11.4), increasing the likelihood of success with cervical traction from 44 to 79.2%.
14 If at least four out of five variables were present, the +LR was equal to 23.1 (2.5-227.9),
15 increasing the post-test probability of having improvement with cervical traction to 94.8%.
16 This preliminary CPR provides the ability to a priori identify patients with neck pain likely
17 to experience a dramatic response with cervical traction and exercise. Before the rule can
18 be implemented in routine clinical practice, future studies are necessary to validate the rule.

19
20 In 2014, Fritz et al. examined the effectiveness of cervical traction in addition to exercise
21 for specific subgroups of patients with neck pain. Patients with neck pain and signs of
22 radiculopathy were randomized to 4 weeks of treatment with exercise, exercise with
23 mechanical traction, or exercise with over-door traction. Secondary outcomes favored
24 mechanical traction at several time points. The validity of the subgrouping rule was
25 supported on the Neck Disability Index at the 6-month time point only. Authors concluded
26 that adding mechanical traction to exercise for patients with cervical radiculopathy resulted
27 in lower disability and pain, particularly at long-term follow-ups.

28
29 Yang et al. (2017) performed a comprehensive search of current literature and conduct a
30 meta-analysis of randomized controlled trials (RCTs) to assess the neck pain relieving
31 effect of intermittent cervical traction (ICT). The meta-analysis included seven RCTs. The
32 results indicated that patients who received ICT for neck pain had significantly lower pain
33 scores than those receiving placebos did immediately after treatment. The pain scores
34 during the follow-up period and the neck disability index scores immediately after
35 treatment and during the follow-up period did not differ significantly. Authors concluded
36 that ICT may have a short-term neck pain-relieving effect. Some risks of bias were noted
37 in the included studies, reducing the evidence level of this meta-analysis. According to

38
39 Blanpied et al. (2017), for patients with chronic neck pain with mobility deficits, clinicians
40 should provide a multimodal approach that may include intermittent mechanical/manual
41 traction. They also report that for patients with chronic neck pain with radiating pain,
42 clinicians should provide mechanical intermittent cervical traction, combined with other

1 interventions such as stretching and strengthening exercise plus cervical and thoracic
2 mobilization/manipulation. However, Bier et al. (2018) states that the physical therapist is
3 advised not to use traction. Romeo et al. (2018) conducted a review and meta-analysis of
4 randomized controlled trials (RCTs) on the effect of cervical traction combined with other
5 physical therapy procedures versus physical therapy procedures alone on pain and
6 disability on patients with cervical radiculopathy (CR). Five studies met the inclusion
7 criteria. Mechanical traction had a significant effect on pain at short- and intermediate-
8 terms and significant effects on disability at intermediate term. Manual traction had
9 significant effects on pain at short- term. Authors conclude that the current literature lends
10 some support to the use of the mechanical and manual traction for CR in addition to other
11 physical therapy procedures for pain reduction but yielding lesser effects on
12 function/disability.

13
14 Colombo et al. (2020) investigated the effectiveness of traction therapy in reducing pain
15 for patients with cervical radicular syndrome (CRS) by performing a systematic review
16 with meta-analysis. A total of seven studies (589 patients), one with low risk of bias, were
17 evaluated. An overall estimate of treatment modalities showed low evidence that adding
18 traction to other treatments is statistically compared to other treatments alone. The
19 subgroup analyses were still statistically significant only for mechanical and continuous
20 modalities. Authors concluded that overall analysis showed that, compared to controls,
21 reduction in pain intensity after traction therapy was achieved in patients with cervical
22 radiculopathy. However, the quality of evidence was generally low and none of these
23 effects were clinically meaningful.

24
25 Jellad et al. (2024) sought to make a preliminary estimate of efficacy of adding mechanical
26 intermittent cervical traction (MICT) to conventional rehabilitation on cervicogenic
27 headache (CGH) in patients with cervical radiculopathy (CR). A total of 36 CR patients
28 with CGH were randomly allocated to 3 equally sized groups (A, B and C). The treatment
29 consisted of twelve sessions of conventional rehabilitation (4 weeks) combined with MICT
30 (2 kg for group A, 8 kg for group B and 12 kg for group C). Primary outcomes were CGH
31 intensity (visual analog scale) and frequency (days per week). Secondary outcomes were
32 radicular pain intensity (visual analog scale), cervical range of motion (cervical range of
33 motion instrument), proprioception (cervical range of motion instrument) and muscle
34 strength (MicroFET2 dynamometer), handgrip strength (handheld dynamometer), function
35 (Neck Disability Index), kinesiophobia (Tampa Scale for Kinesiophobia), anxiety and
36 depression (Hospital Anxiety and Depression questionnaire), and quality of life (World
37 Health Organization Quality of Life). Patients were assessed at baseline, one, three and six
38 months after the beginning of treatment. At one-, three- and six-month follow-ups, Group
39 C exhibited the highest improvement in CGH intensity and frequency compared to the
40 other groups. Both groups C and B showed a significant improvement in radicular pain
41 compared to group A at one month follow-up. The improvement in group C was
42 significantly better in terms of function and anxiety at three months and quality of life at

1 six months. Authors concluded that the blend of conventional rehabilitation alongside 12
 2 kg MICT seems to be efficacious in diminishing both the intensity and frequency of CGH
 3 in patients with CR. These advantages appear to last for up to six months following the
 4 treatment period, potentially leading to decreased CGH severity and occurrence rates,
 5 heightened functionality, reduced anxiety levels, and an overall enhancement in quality of
 6 life. These findings are preliminary and require confirmation in larger trials.

7 8 **Lumbar**

9 According to the Philadelphia Panel Evidence-Based Clinical Practice Guidelines on
 10 Selected Rehabilitation Interventions for Low Back Pain publication (2001), mechanical
 11 traction for chronic LBP was not effective or beneficial for pain, function, patient global
 12 assessment, and return to work. This was based on four (4) RCTs of mechanical traction
 13 versus placebo or no treatment and rated as level I (good evidence). A larger Cochrane
 14 Collaboration systematic review by Clarke et al. (2009) determined similar results (25
 15 RCTs). Available studies in this review involved mixed groups of acute, sub-acute and
 16 chronic patients with LBP with and without sciatica and were all consistent, indicating that
 17 continuous or intermittent traction as a single treatment for LBP is not likely effective for
 18 these patients. Traction for patients with sciatica cannot be judged effective at present
 19 either, due to inconsistent results and methodological problems in most studies (Clarke et
 20 al., 2009). An updated Cochrane review published in 2013 by Wegner et al. indicated that
 21 traction, either alone or in combination with other treatments, has little or no impact on
 22 pain intensity, functional status, global improvement and return to work among people with
 23 LBP (with or without sciatica). The effects shown by the included studies were small and
 24 not clinically relevant. These conclusions were applicable to both manual and mechanical
 25 traction.

26
 27 One study attempted to determine which subcategory of patients with LBP would most
 28 benefit from mechanical traction. Fritz et al. (2007) determined that patients with sciatica,
 29 signs of nerve root compression, and either peripheralization with extension movements or
 30 a positive crossed straight leg raise test were most likely to benefit from a combined
 31 traction and extension-oriented physical therapy intervention. The authors reported
 32 improvements in both disability (Oswestry Disability Questionnaire) and fear-avoidance
 33 beliefs (Fear Avoidance Belief Questionnaire) in the combined traction/extension-oriented
 34 approach group at two weeks compared to the group that received just an extension-
 35 oriented approach. This study provides some initial evidence for the use of traction for the
 36 subgroup of patients mentioned above. The primary limitation to this study is the type of
 37 traction table used is not one that is typically found in most clinical settings. The authors
 38 used a mechanical traction table allowing for modifications of a subject's position in
 39 flexion/extension, rotation or side-bending (3-dimensional ActiveTrac table, The Saunders
 40 Group, Inc.). The following parameters were utilized: static traction for a maximum of 12
 41 minutes (10 minutes at desired intensity and one minute ramp up/down) at 40% - 60% of
 42 the patient's body weight for a maximum of 12 sessions during a 6-week period (four

1 sessions/week during the first two weeks then one session/week during weeks three
 2 through six). Thackeray et al. (2016) examined the effectiveness of mechanical traction in
 3 patients with lumbar nerve root compression and within a predefined subgroup. One
 4 hundred twenty patients with low back pain with nerve root compression were recruited
 5 from physical therapy clinics. Using predefined subgrouping criteria, patients were
 6 stratified at baseline and randomized to receive an extension-oriented treatment approach
 7 with or without the addition of mechanical traction. During a 6-week period, patients
 8 received up to 12 treatment visits. Primary outcomes of pain and disability were collected
 9 at 6 weeks, 6 months, and 1 year by assessors blinded to group allocation. No significant
 10 differences in disability or pain outcomes were noted between treatment groups at any time
 11 point, nor was any interaction found between subgroup status and treatment. Authors
 12 concluded that patients with lumbar nerve root compression presenting for physical therapy
 13 can expect significant changes in disability and pain over a 6-week treatment period. There
 14 is no evidence that mechanical lumbar traction in combination with an extension-oriented
 15 treatment is superior to extension-oriented exercises alone in the management of these
 16 patients or within a predefined subgroup of patients.

17
 18 The North American Spine Society's clinical practice guideline on "Diagnosis and
 19 treatment of degenerative lumbar spinal stenosis" (2011) noted that there is insufficient
 20 evidence to make a recommendation for or against traction, electrical stimulation or
 21 transcutaneous electrical nerve stimulation for the treatment of patients with lumbar spinal
 22 stenosis.

23
 24 According to the AHRQ publication on Non-Invasive Techniques for Low Back Pain
 25 (2016):

- 26 • For low back pain with or without radicular symptoms, a systematic review
 27 included 13 trials that found no clear differences with inconsistent effects of
 28 traction versus placebo, sham, or no treatment in pain, function, or other outcomes,
 29 though two trials reported favorable effects on pain in patients with radicular back
 30 pain (SOE: insufficient for pain and function).
- 31 • For low back pain with or without radicular symptoms, a systematic review
 32 included five trials that found no clear differences between traction versus
 33 physiotherapy versus physiotherapy alone.
- 34 • For low back pain with or without radicular symptoms, a systematic review
 35 included 15 trials of traction versus other interventions that found no clear
 36 difference between traction versus other active interventions in pain or function
 37 (SOE: low for pain and function).
- 38 • A systematic review included five trials that found no clear differences between
 39 different types of traction.
- 40 • Eleven trials of traction in a systematic review reported no adverse events or no
 41 difference in risk of adverse events versus placebo or other interventions. Three
 42 subsequent trials reported findings consistent with the systematic review.

1 According to the American College of Physician’s clinical practice guideline (2017) on
2 noninvasive treatments for acute, subacute, and chronic low back pain, evidence was
3 insufficient to determine the effectiveness of traction tables/devices. Foster et al. (2018)
4 summarizes that passive electrical or physical modalities, such as traction, interferential
5 therapy, short-wave diathermy, are generally ineffective and not recommended.

6
7 Bilgilisoy Filiz et al. (2018) compared the effects of mechanical lumbar traction either in
8 the supine or in the prone position with conventional physical therapy (PT) in patients with
9 chronic low back pain and lumbosacral nerve root involvement in terms of disability, pain,
10 and mobility. Participants (N = 125) were randomly assigned to receive 15 sessions of PT
11 with additional mechanical lumbar traction either in the supine position (supine traction
12 group) or in the prone position (prone traction group) or only PT without traction (PT only
13 group). Patients were assessed at baseline and at the end of the PT sessions in terms of
14 disability, pain, and mobility. Disability was assessed using the modified Oswestry
15 Disability Index; pain was assessed using a visual analog scale, and lumbar mobility was
16 assessed using the modified lumbar Schober test. One hundred eighteen patients completed
17 the trial. All groups improved significantly for all outcomes. In the between-group analysis,
18 improvements of Oswestry Disability Index and visual analog scale were found
19 significantly better in the prone traction group compared with the PT only group. Authors
20 concluded that the addition of traction in the prone position to other modalities resulted in
21 larger immediate improvements in terms of pain and disability, and the results suggest that
22 when using traction, prone traction might be first choice. Chou et al. (2018) states that
23 clinicians should not offer traction for neck and back pain given lack of effectiveness.

24
25 Kuligowski et al. (2019) completed a study that enrolled 37 people aged 22-35. The
26 subjects underwent radiological evaluation (MRI), which constituted the basis for
27 assigning them to one of two groups: a protrusion group (PRO) or an extrusion group
28 (EXT). During the experiment, the patient was in the supine position while the therapist
29 administered three-dimensional traction using a manual therapy belt. Authors concluded
30 the following: 1. The type of intervertebral disc damage determines the functional status of
31 young people with degenerative disc disease. 2. The study demonstrated and confirmed a
32 positive effect of traction on the functional status of subjects with lumbar disc herniation.
33 3. Traction techniques are safe and can be successfully used in the treatment of lumbar disc
34 herniation as noted on MRI.

35
36 Hirayama et al. (2019) sought to develop a clinical prediction rule (CPR) that predicts
37 treatment responses to mechanical lumbar traction (MLT) among patients with lumbar disc
38 herniation (LDH). The subjects included 103 patients diagnosed with LDH for which they
39 underwent conservative therapy. The subjects received MLT for 2 weeks, and the
40 application of any other medication was left at the discretion of the attending physician.
41 The patients whose ODI after 2 weeks of treatment improved by $\geq 50\%$ of that at the initial
42 evaluation were defined as responders. Of the 103 subjects, 24 were responders, and the

1 five predictors selected for the CPR were limited lumbar extension range of motion, low-
2 level fear-avoidance beliefs regarding work, no segmental hypomobility in the lumbar
3 spine, short duration of symptoms, and sudden onset of symptoms. For the patients with at
4 least three of the five predictors, the probability of their ODI greatly improving increased
5 from 23.3% to 48.7% compared with the patients without these predictors (positive
6 likelihood ratio, 3.13).

7
8 Cheng et al. (2020) evaluated the effectiveness of traction in improving low back pain,
9 functional outcome, and disk morphology in patients with herniated intervertebral disks.
10 Seven articles involving 403 participants were included for quantitative analysis.
11 Compared with the control group, the participants in the traction group showed
12 significantly greater improvements in pain and function in the short term, with standard
13 mean differences of 0.44 (95% confidence interval (CI): 0.11-0.77) and 0.42 (95% CI:
14 0.08-0.76), respectively. The standard mean differences were not significant to support the
15 long-term effects on pain and function, nor the effects on herniated disk size. Authors
16 concluded that compared with sham or no traction, lumbar traction exhibited significantly
17 more pain reduction and functional improvements in the short term, but not in the long
18 term. There is insufficient evidence to support the effect of lumbar traction on herniated
19 disk size reduction.

20
21 Vanti et al. (2020) evaluated the effects of different types of traction added to or compared
22 with conservative treatments on pain and disability for patients with lumbar radiculopathy
23 (LR) in a systematic review and meta-analysis. Eight studies met the inclusion criteria, and
24 5 were meta-analyzed. Meta-analyses of results from low-quality studies indicated that
25 supine mechanical traction added to physical therapist treatments had significant effects on
26 pain and disability. Analyses of results from high-quality studies of prone mechanical
27 traction added to physical therapist intervention for pain and disability were not significant.
28 These results were also evident at short-term follow-up (up to 3 months after intervention).
29 Authors concluded that the literature suggests that, for pain and disability in LR, there is
30 short-term effectiveness of supine mechanical traction when added to physical therapist
31 intervention.

32
33 George et al. (2021) state that physical therapists should not use mechanical traction for
34 patients with chronic LBP with leg pain, based on the lack of benefit when added to other
35 interventions in an updated clinical practice guideline.

36
37 Farrokhi et al. (2024) explored associations between the utilization of active, passive, and
38 manual therapy interventions for low back pain (LBP) with 1-year escalation-of-care
39 events, including opioid prescriptions, spinal injections, specialty care visits, and
40 hospitalizations. This was a retrospective cohort study of 4827 patients identified via the
41 Military Health System Data Repository who received physical therapist care for LBP in 4
42 outpatient clinics between January 1, 2015, and January 1, 2018. One-year escalation-of-

1 care events were evaluated based on type of physical therapist interventions (i.e., active,
2 passive, or manual therapy) received using adjusted odds ratios. Most patients (89.9%)
3 received active interventions. Patients with 10% higher proportion of visits that included
4 at least 1 passive intervention had a 3% to 6% higher likelihood of 1-year escalation-of-
5 care events. Similarly, with 10% higher proportion of passive to active interventions used
6 during the course of care, there was a 5% to 11% higher likelihood of 1-year escalation-of-
7 care events. When compared to patients who received active interventions only, the
8 likelihood of incurring 1-year escalation-of-care events was 50% to 220% higher for those
9 who received mechanical traction and 2 or more different passive interventions, but lower
10 by 50% for patients who received manual therapy. Authors concluded that greater use of
11 passive interventions for LBP was associated with elevated odds of 1-year escalation-of-
12 care events. In addition, the use of specific passive interventions such as mechanical
13 traction in conjunction with active interventions resulted in suboptimal escalation-of-care
14 events, while the use of manual therapy was associated with more favorable downstream
15 health care outcomes.

16 **PRACTITIONER SCOPE AND TRAINING**

17 Practitioners should practice only in the areas in which they are competent based on their
18 education, training, and experience. Levels of education, experience, and proficiency may
19 vary among individual practitioners. It is ethically and legally incumbent on a practitioner
20 to determine where they have the knowledge and skills necessary to perform such services
21 and whether the services are within their scope of practice.

22
23
24 It is best practice for the practitioner to appropriately render services to a member only if
25 they are trained, equally skilled, and adequately competent to deliver a service compared
26 to others trained to perform the same procedure. If the service would be most competently
27 delivered by another health care practitioner who has more skill and training, it would be
28 best practice to refer the member to the more expert practitioner.

29
30 Best practice can be defined as a clinical, scientific, or professional technique, method, or
31 process that is typically evidence-based and consensus driven and is recognized by a
32 majority of professionals in a particular field as more effective at delivering a particular
33 outcome than any other practice (Joint Commission International Accreditation Standards
34 for Hospitals, 2020).

35
36 Depending on the practitioner's scope of practice, training, and experience, a member's
37 condition and/or symptoms during examination or the course of treatment may indicate the
38 need for referral to another practitioner or even emergency care. In such cases it is prudent
39 for the practitioner to refer the member for appropriate co-management (e.g., to their
40 primary care physician) or if immediate emergency care is warranted, to contact 911 as
41 appropriate. See the *Managing Medical Emergencies (CPG 159 – S)* clinical practices
42 guideline for information.

1 **References**

2 American Medical Association. (current year). Current Procedural Terminology (CPT)
3 Current year (rev. ed.). Chicago: AMA

4
5 Bergman S. Management of musculoskeletal pain. *Best Pract Res Clin Rheumatol.*
6 2007;21(1):153-166. doi:10.1016/j.berh.2006.10.001

7
8 Bier JD, Scholten-Peeters WGM, Staal JB, et al. Clinical Practice Guideline for Physical
9 Therapy Assessment and Treatment in Patients With Nonspecific Neck Pain. *Phys*
10 *Ther.* 2018;98(3):162-171. doi:10.1093/ptj/pzx118

11
12 Bilgilişoy Filiz M, Kiliç Z, Uçkun A, Çakir T, Koldaş Doğan Ş, Toraman NF. Mechanical
13 Traction for Lumbar Radicular Pain: Supine or Prone? A Randomized Controlled
14 Trial. *Am J Phys Med Rehabil.* 2018;97(6):433-439.
15 doi:10.1097/PHM.0000000000000892

16
17 Blanpied PR, Gross AR, Elliott JM, et al. Neck Pain: Revision 2017. *J Orthop Sports Phys*
18 *Ther.* 2017;47(7):A1-A83. doi:10.2519/jospt.2017.0302

19
20 Borman P, Keskin D, Ekici B, Bodur H. The efficacy of intermittent cervical traction in
21 patients with chronic neck pain. *Clin Rheumatol.* 2008;27(10):1249-1253.
22 doi:10.1007/s10067-008-0895-z

23
24 Cameron MH. (2017). *Physical Agents in Rehabilitation: From Research to Practice.* (5th
25 ed.). Saunders

26
27 Centers for Medicare and Medicaid Services. Local Coverage Determination (LCD):
28 Outpatient Physical and Occupational Therapy Services (L33631) Retrieved on
29 November 23, 2024 from [https://www.cms.gov/medicare-coverage-](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)
30 [database/details/lcd-](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)
31 [details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advance](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)
32 [d&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)
33 [%7cSAD%7cFAQ&PolicyType=Final&s=---](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)
34 [%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)
35 [laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&b](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)
36 [c=IAAAABAAAA&](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33631&ver=51&NCDId=72&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCD%7cTA&ArticleType=Ed%7cKey%7cSAD%7cFAQ&PolicyType=Final&s=---%7c5%7c6%7c66%7c67%7c9%7c38%7c63%7c41%7c64%7c65%7c44&Keyword=laser+procedures&KeywordLookUp=Doc&KeywordSearchType=And&kq=true&bc=IAAAABAAAA&)

37
38 Cheng YH, Hsu CY, Lin YN. The effect of mechanical traction on low back pain in patients
39 with herniated intervertebral disks: a systemic review and meta-analysis. *Clin Rehabil.*
40 2020;34(1):13-22. doi:10.1177/0269215519872528

- 1 Chou R, Huffman LH; American Pain Society; American College of Physicians.
 2 Nonpharmacologic therapies for acute and chronic low back pain: a review of the
 3 evidence for an American Pain Society/American College of Physicians clinical
 4 practice guideline [published correction appears in *Ann Intern Med.* 2008 Feb
 5 5;148(3):247-8]. *Ann Intern Med.* 2007;147(7):492-504. doi:10.7326/0003-4819-147-
 6 7-200710020-00007
 7
- 8 Chou R, Deyo R, Friedly J, et al. *Noninvasive Treatments for Low Back Pain.* Rockville
 9 (MD): Agency for Healthcare Research and Quality (US); February 2016
 10
- 11 Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint
 12 clinical practice guideline from the American College of Physicians and the American
 13 Pain Society [published correction appears in *Ann Intern Med.* 2008 Feb 5;148(3):247-
 14 8]. *Ann Intern Med.* 2007;147(7):478-491. doi:10.7326/0003-4819-147-7-200710020-
 15 00006
 16
- 17 Chou R, Côté P, Randhawa K, et al. The Global Spine Care Initiative: applying evidence-
 18 based guidelines on the non-invasive management of back and neck pain to low- and
 19 middle-income communities. *Eur Spine J.* 2018;27(Suppl 6):851-860.
 20 doi:10.1007/s00586-017-5433-8
 21
- 22 Clarke J, van Tulder M, Blomberg S, de Vet H, van der Heijden G, Bronfort G. Traction
 23 for low back pain with or without sciatica: an updated systematic review within the
 24 framework of the Cochrane collaboration. *Spine (Phila Pa 1976).* 2006;31(14):1591-
 25 1599. doi:10.1097/01.brs.0000222043.09835.72
 26
- 27 Clarke JA, van Tulder MW, Blomberg SE, et al. Traction for low-back pain with or without
 28 sciatica. *Cochrane Database Syst Rev.* 2007;(2):CD003010. Published 2007 Apr 18.
 29 doi:10.1002/14651858.CD003010.pub4
 30
- 31 Colombo C, Salvioli S, Gianola S, Castellini G, Testa M. Traction Therapy for Cervical
 32 Radicular Syndrome is Statistically Significant but not Clinically Relevant for Pain
 33 Relief. A Systematic Literature Review with Meta-Analysis and Trial Sequential
 34 Analysis. *J Clin Med.* 2020;9(11):3389. Published 2020 Oct 22.
 35 doi:10.3390/jcm9113389
 36
- 37 Farrokhi S, Bechard L, Gorczynski S, et al. The Influence of Active, Passive, and Manual
 38 Therapy Interventions for Low Back Pain on Opioid Prescription and Health Care
 39 Utilization. *Phys Ther.* 2024;104(3):pzad173. doi:10.1093/ptj/pzad173

- 1 Foster NE, Anema JR, Cherkin D, et al. Prevention and treatment of low back pain:
2 evidence, challenges, and promising directions. *Lancet*. 2018;391(10137):2368-2383.
3 doi:10.1016/S0140-6736(18)30489-6
4
- 5 Fritz, J. M., Lindsay, W., Matheson, J. W., Brennan, G. P., Hunter, S. J., Moffit, S. D.,
6 Swalberg, A., & Rodriguez, B. (2007). Is there a subgroup of patients with low back
7 pain likely to benefit from mechanical traction? Results of a randomized clinical trial
8 and subgrouping analysis. *Spine*, 32(26), E793–E800.
9 <https://doi.org/10.1097/BRS.0b013e31815d001a>
10
- 11 Fritz JM, Thackeray A, Brennan GP, Childs JD. Exercise only, exercise with mechanical
12 traction, or exercise with over-door traction for patients with cervical radiculopathy,
13 with or without consideration of status on a previously described subgrouping rule: a
14 randomized clinical trial. *J Orthop Sports Phys Ther*. 2014;44(2):45-57.
15 doi:10.2519/jospt.2014.5065
16
- 17 George SZ, Fritz JM, Silfies SP, et al. Interventions for the Management of Acute and
18 Chronic Low Back Pain: Revision 2021. *J Orthop Sports Phys Ther*.
19 2021;51(11):CPG1-CPG60. doi:10.2519/jospt.2021.0304
20
- 21 Graham N, Gross AR, Carlesso LC, et al. An ICON Overview on Physical Modalities for
22 Neck Pain and Associated Disorders. *Open Orthop J*. 2013;7:440-460. Published 2013
23 Sep 20. doi:10.2174/1874325001307010440
24
- 25 Gross AR, Goldsmith C, Hoving JL, et al. Conservative management of mechanical neck
26 disorders: a systematic review. *J Rheumatol*. 2007;34(5):1083-1102
27
- 28 Guzman J, Haldeman S, Carroll LJ, et al. Clinical practice implications of the Bone and
29 Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: from
30 concepts and findings to recommendations. *Spine (Phila Pa 1976)*. 2008;33(4
31 Suppl):S199-S213. doi:10.1097/BRS.0b013e3181644641
32
- 33 Haldeman S, Carroll L, Cassidy JD, Schubert J, Nygren A; Bone and Joint Decade 2000-
34 2010 Task Force on Neck Pain and Its Associated Disorders. The Bone and Joint
35 Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: executive
36 summary. *Spine (Phila Pa 1976)*. 2008;33(4 Suppl):S5-S7.
37 doi:10.1097/BRS.0b013e3181643f40
38
- 39 Hirayama K, Tsushima E, Arihara H, Omi Y. Developing a clinical prediction rule to
40 identify patients with lumbar disc herniation who demonstrate short-term improvement
41 with mechanical lumbar traction. *Phys Ther Res*. 2019;22(1):9-16. Published 2019 Apr
42 20. doi:10.1298/ptr.E9973

- 1 Hurwitz EL, Carragee EJ, van der Velde G, et al. Treatment of neck pain: noninvasive
2 interventions: results of the Bone and Joint Decade 2000-2010 Task Force on Neck
3 Pain and Its Associated Disorders. *J Manipulative Physiol Ther.* 2009;32(2
4 Suppl):S141-S175. doi:10.1016/j.jmpt.2008.11.017
5
- 6 Jellad A, Kalai A, Chaabeni A, et al. Effect of cervical traction on cervicogenic headache
7 in patients with cervical radiculopathy: a preliminary randomized controlled trial. *BMC*
8 *Musculoskelet Disord.* 2024;25(1):842. Published 2024 Oct 24. doi:10.1186/s12891-
9 024-07930-z
10
- 11 Joint Commission International. (2020). Joint Commission International Accreditation
12 Standards for Hospitals (7th ed.): Joint Commission Resources
13
- 14 Kuligowski T, Dębiec-Bąk A, Skrzek A. Effectiveness of Traction in Young Patients
15 Representing Different Stages of Degenerative Disc Disease. *Ortop Traumatol*
16 *Rehabil.* 2019;21(3):187-195. doi:10.5604/01.3001.0013.2925
17
- 18 Mancuso T, Poole JL. The effect of paraffin and exercise on hand function in persons with
19 scleroderma: a series of single case studies. *J Hand Ther.* 2009;22(1):71-78.
20 doi:10.1016/j.jht.2008.06.009
21
- 22 Philadelphia Panel. Philadelphia Panel evidence-based clinical practice guidelines on
23 selected rehabilitation interventions for knee pain. *Phys Ther.* 2001;81(10):1675-1700
24
- 25 Philadelphia Panel. Philadelphia Panel evidence-based clinical practice guidelines on
26 selected rehabilitation interventions for shoulder pain. *Phys Ther.* 2001;81(10):1719-
27 1730
28
- 29 Philadelphia Panel. Philadelphia Panel evidence-based clinical practice guidelines on
30 selected rehabilitation interventions for low back pain. *Phys Ther.* 2001;81(10):1641-
31 1674
32
- 33 Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and
34 Chronic Low Back Pain: A Clinical Practice Guideline From the American College of
35 Physicians. *Ann Intern Med.* 2017;166(7):514-530. doi:10.7326/M16-2367
36
- 37 Raney NH, Petersen EJ, Smith TA, et al. Development of a clinical prediction rule to
38 identify patients with neck pain likely to benefit from cervical traction and
39 exercise. *Eur Spine J.* 2009;18(3):382-391. doi:10.1007/s00586-008-0859-7
40
- 41 Romeo A, Vanti C, Boldrini V, et al. Cervical Radiculopathy: Effectiveness of Adding
42 Traction to Physical Therapy-A Systematic Review and Meta-Analysis of Randomized

- 1 Controlled Trials [published correction appears in Phys Ther. 2018 Aug
2 1;98(8):727]. Phys Ther. 2018;98(4):231-242. doi:10.1093/physth/pzy001
3
- 4 Thackeray A, Fritz JM, Childs JD, Brennan GP. The Effectiveness of Mechanical Traction
5 Among Subgroups of Patients With Low Back Pain and Leg Pain: A Randomized
6 Trial. J Orthop Sports Phys Ther. 2016;46(3):144-154. doi:10.2519/jospt.2016.6238
7
- 8 Vanti C, Panizzolo A, Turone L, et al. Effectiveness of Mechanical Traction for Lumbar
9 Radiculopathy: A Systematic Review and Meta-Analysis. Phys Ther.
10 2021;101(3):pzaa231. doi:10.1093/ptj/pzaa231
11
- 12 Verhagen AP, Scholten-Peeters GG, de Bie RA, Bierma-Zeinstra SM. Conservative
13 treatments for whiplash. Cochrane Database Syst Rev. 2004;(1):CD003338.
14 doi:10.1002/14651858.CD003338.pub2
15
- 16 Waddell G. (1998). The clinical course of low back pain (pp103-117) The Back Pain
17 Revolution. Churchill Livingstone.
18
- 19 Walsh NE, Brooks P, Hazes JM, et al. Standards of care for acute and chronic
20 musculoskeletal pain: the Bone and Joint Decade (2000-2010). Arch Phys Med
21 Rehabil. 2008;89(9):1830-1845. doi:10.1016/j.apmr.2008.04.009
22
- 23 Watters WC 3rd, Bono CM, Gilbert TJ, et al. An evidence-based clinical guideline for the
24 diagnosis and treatment of degenerative lumbar spondylolisthesis. Spine J.
25 2009;9(7):609-614. doi:10.1016/j.spinee.2009.03.016
26
- 27 Wegner I, Widyahening IS, van Tulder MW, et al. Traction for low-back pain with or
28 without sciatica. Cochrane Database Syst Rev. 2013;2013(8):CD003010. Published
29 2013 Aug 19. doi:10.1002/14651858.CD003010.pub5
30
- 31 Whitman JM, Flynn TW, Childs JD, et al. A comparison between two physical therapy
32 treatment programs for patients with lumbar spinal stenosis: a randomized clinical
33 trial. Spine (Phila Pa 1976). 2006;31(22):2541-2549.
34 doi:10.1097/01.brs.0000241136.98159.8c
35
- 36 Yang JD, Tam KW, Huang TW, Huang SW, Liou TH, Chen HC. Intermittent Cervical
37 Traction for Treating Neck Pain: A Meta-analysis of Randomized Controlled
38 Trials. Spine (Phila Pa 1976). 2017;42(13):959-965.
39 doi:10.1097/BRS.0000000000001948
40
- 41 Young IA, Michener LA, Cleland JA, Aguilera AJ, Snyder AR. Manual therapy, exercise,
42 and traction for patients with cervical radiculopathy: a randomized clinical trial

1 [published correction appears in Phys Ther. 2009 Nov;89(11):1254-5] [published
2 correction appears in Phys Ther. 2010 May;90(5):825]. Phys Ther. 2009;89(7):632-
3 642. doi:10.2522/ptj.20080283